

APPENDIX A.

FOCAL FISH

This chapter was drafted in 2004 and contains minor revisions, namely corrections to viability status and assessment results for 2010.

APPENDIX A. FOCAL FISH

A.1. CHINOOK SALMON	A-11
A.1.1. Life History and Requirements	A-11
A.1.2. Distribution	A-19
A.1.3. Genetic Diversity.....	A-20
A.1.4. ESU Definition	A-23
A.1.5. Life History Differences.....	A-23
A.1.6. Abundance	A-25
A.1.7. Productivity.....	A-28
A.1.8. Hatchery Production	A-29
A.1.9. Fishery	A-37
A.1.10. Assessment of Current Status and Limiting Factors	A-70
A.2. COHO SALMON.....	A-85
A.2.1. Life History and Requirements	A-85
A.2.2. Distribution	A-92
A.2.3. Genetic Diversity.....	A-94
A.2.4. ESU Definition	A-94
A.2.5. Life History Diversity	A-95
A.2.6. Abundance	A-95
A.2.7. Productivity.....	A-96
A.2.8. Hatchery Production	A-97
A.2.9. Fishery	A-105
A.2.10. Assessments of Current Status and Limiting Factors	A-126
A.3. CHUM SALMON	A-138
A.3.1. Life History and Requirements	A-139
A.3.2. Distribution	A-146
A.3.3. Genetic Diversity.....	A-147
A.3.4. ESU Definition	A-148
A.3.5. Life History Diversity	A-149
A.3.6. Abundance	A-149
A.3.7. Productivity.....	A-153
A.3.8. Hatchery Production	A-153
A.3.9. Fishery	A-155
A.3.10. Assessment of Current Status and Limiting Factors	A-160
A.4. STEELHEAD	A-169
A.4.1. Life History and Requirements	A-170
A.4.2. Distribution	A-178
A.4.3. Genetic Diversity.....	A-180
A.4.4. DPS Definition	A-182
A.4.5. Life History Differences.....	A-182
A.4.6. Abundance	A-183
A.4.7. Productivity.....	A-189
A.4.8. Hatchery Production	A-190
A.4.9. Fishery	A-198
A.4.10. Assessment of Current Status and Limiting Factors	A-213

A.5. BULL TROUT	A-226
A.5.1. Life History and Requirements	A-226
A.5.2. Distribution	A-232
A.5.3. Genetic Diversity.....	A-233
A.5.4. ESU Definition	A-233
A.5.5. Life History Diversity	A-234
A.5.6. Abundance	A-234
A.5.7. Productivity.....	A-236
A.5.8. Fishery	A-236
A.5.9. Assessment of Current Status and Limiting Factors	A-237
A.6. REFERENCES	A-239

Tables

Table A-1.	Current (2003 brood) annual release goals of spring Chinook salmon juveniles (subyearling and yearling) into lower Columbia basins.	A-30
Table A-2.	Current (2003 brood) annual release goals of fall Chinook salmon juveniles (subyearling and yearling) into Washington lower Columbia basins.	A-32
Table A-3.	Example of lower Columbia spring Chinook harvest exploitation rates under current management.	A-40
Table A-4.	Sliding scale* of harvest impacts on wild upriver spring Chinook based on Snake River wild spring Chinook run size (adapted from the 2001-05 Interim Management Agreement).....	A-43
Table A-5.	Allocation of non-Indian upriver wild spring Chinook impacts based on Willamette hatchery and upriver wild spring Chinook abundance.	A-44
Table A-6.	Spring Chinook catch and released during the 2002 non-Indian commercial tangle net fishery in the lower Columbia River.	A-45
Table A-7.	Ex-vessel value (in thousands of dollars expressed in 2002 dollars) of in-river commercial harvest of Columbia River spring Chinook, 1988–2002.....	A-46
Table A-8.	Example of lower Columbia fall Chinook current harvest exploitation and distribution under current management.	A-53
Table A-9.	Management regime for Canadian Chinook salmon fisheries affected by the PST.	A-54
Table A-10.	PFMC pre-season adopted Chinook catch quotas (in thousands of fish) for ocean fisheries north of Cape Falcon and critical stocks driving management, 1983–2001.....	A-56
Table A-11.	Management criteria and projected key stock escapements (in thousands of fish) for Chinook salmon in PFMC-adopted ocean salmon fisheries, 2003*.....	A-60
Table A-12.	Tentatively adopted 2003 fishery management measures for PFMC fisheries to mitigate potential impacts on ESA-listed ESUs of lower Columbia River salmonids.	A-61
Table A-13.	Stock accountability of fall Chinook returning to Columbia River, 1980–2002.....	A-63
Table A-14.	Stock composition of adult fall Chinook landed in mainstem Columbia River fisheries, 2001.....	A-63
Table A-15.	Baseline status and recovery objectives for lower Columbia River Chinook populations (relative to listing period baseline).	A-75
Table A-16.	Productivity improvements and impact reduction benchmarks consistent with recovery of Washington lower Columbia River Chinook populations if reductions are distributed across factors in proportion to their baseline impact.	A-77
Table A-17.	Presumed reductions in wild population fitness as a result of natural hatchery spawners and survival as a result of interactions with other hatchery species for Washington lower Columbia River Chinook populations.	A-81
Table A-18.	Coho salmon CWT recoveries in various fisheries.	A-93
Table A-19.	Average annual releases of coho salmon juveniles (fry and smolts) from Washington lower Columbia hatchery facilities during release years 1987–91 (NMFS 1995).	A-97
Table A-20.	Current (2003 brood) annual release goals of coho salmon smolts from Washington lower Columbia hatchery facilities.....	A-98
Table A-21.	Example of lower Columbia coho harvest exploitation and distribution under current management (combined early and late stock).	A-108
Table A-22.	Harvest management matrix identifying allowable fishery impacts and ranges of resulting recruitment based on parental spawner abundance and marine survival (OCN work group revisions to original PFMC matrix).....	A-113
Table A-23.	Harvest management matrix for Lower Columbia Natural (LCN) coho with maximum allowable ocean fishery mortality rates.....	A-114

Table A-24.	Management criteria and projected key stock escapements (in thousands of fish) for coho salmon in PFMC-adopted ocean salmon fisheries, 2003.....	A-115
Table A-25.	Expected mark rates for Council-adopted ocean salmon fisheries with selective coho retention, 2003.....	A-116
Table A-26.	Harvest management matrix for Lower Columbia Natural (LCN) coho with maximum allowable freshwater fishery mortality rates.....	A-123
Table A-27.	Cumulative exploitation rates for LCN coho under the combined management protocols proposed for setting ocean and in-river fishery harvest rates.	A-124
Table A-28.	Angler trips and economic impact (in 2002 dollars) of the Buoy 10 recreational fishery, 1982–2002.....	A-124
Table A-29.	Estimated catch, mortality, and escapement of marked and unmarked Columbia River basin coho salmon, 2002.....	A-126
Table A-30.	Status and recovery goals for lower Columbia River coho populations (relative to listing baseline).	A-129
Table A-31.	Productivity improvements and impact reduction benchmarks consistent with recovery of Washington lower Columbia River coho populations if reductions are distributed across factors in proportion to their baseline impact.	A-131
Table A-32.	Presumed reductions in wild population fitness as a result of natural hatchery spawners and survival as a result of interactions with other hatchery species for Washington lower Columbia River coho populations.	A-134
Table A-33.	Current (2003 brood) chum juvenile hatchery release goals for Washington lower Columbia tributaries.....	A-155
Table A-34.	Example of current chum harvest.	A-157
Table A-35.	Preliminary 2002 chum salmon harvest in ocean fisheries managed under PST.	A-158
Table A-36.	Harvest related mortality estimates for ESA-listed chum salmon in Columbia River basin fisheries during August–December, 2002.....	A-160
Table A-37.	Status and recovery goals for lower Columbia River chum populations (relative to listing baseline).	A-162
Table A-38.	Productivity improvements and impact reduction benchmarks consistent with recovery of Washington lower Columbia River chum populations if reductions are distributed across factors in proportion to their baseline impact.	A-164
Table A-39.	Presumed reductions in wild population fitness as a result of natural hatchery spawners and survival as a result of interactions with other hatchery species for Washington lower Columbia River chum populations.	A-166
Table A-40.	Lower Columbia River steelhead stock status as determined by SASSI 2002.	A-184
Table A-41.	Escapement of winter steelhead in lower Colcumbia Washington tributaries.	A-188
Table A-42.	Current (2003 brood) summer steelhead smolt release goals.....	A-190
Table A-43.	Current (2003 brood) Winter steelhead smolt release goals.....	A-193
Table A-44.	Example of lower Columbia steelhead harvest exploitation and distribution under current management.	A-201
Table A-45.	Estimated take of ESA-listed steelhead in Washington tributary recreational fisheries... A-204	
Table A-46.	Wild and hatchery contribution to Group A and Group B Index summer steelhead returns to Bonneville Dam, 1984–2003.....	A-205
Table A-47.	Wild steelhead catch (in thousands of fish) in treaty Indian fisheries by index group, 1985–2001.....	A-206
Table A-48.	Harvest related mortality estimates for listed steelhead ESUs in proposed Columbia River basin fisheries* during August—December 2002.	A-207

Table A-49.	Estimated total harvest of steelhead in the 2002 proposed treaty Indian fall fisheries and incidental harvest by ESU.	A-207
Table A-50.	Summary of catch rate, condition, and survival for steelhead captured in various spring fisheries, 2000–02 ^a	A-212
Table A-51.	Estimated take of ESA-listed steelhead in Washington tributary recreational fisheries... A-213	A-213
Table A-52.	Recovery goals for lower Columbia River steelhead populations (relative to listing baseline).	A-217
Table A-53.	Productivity improvements and impact reduction benchmarks consistent with recovery of lower Washington Columbia River steelhead populations if reductions are distributed across factors in proportion to their baseline impact.	A-219
Table A-54.	Presumed reductions in wild population fitness as a result of natural hatchery spawners for Washington lower Columbia River steelhead populations.	A-222
Table A-55.	Life history and population characteristics of bull trout originating in Washington portions of the lower Columbia.	A-234
Table A-56.	Bull trout collected from the Yale tailrace (Lake Merwin) and transferred to the mouth of Cougar Creek (Yale Reservoir) or released back into Yale Reservoir (1995–2000).	A-235

Figures

Figure A-1.	Washington lower Columbia spring Chinook life cycle.....	A-12
Figure A-2.	Washington lower Columbia fall Chinook life cycle.....	A-12
Figure A-3.	Historical demographically independent fall Chinook salmon populations in the lower Columbia River ESU (Myers et al. 2002).	A-21
Figure A-4.	Historical demographically independent spring Chinook salmon populations in the lower Columbia River ESU (Meyers et al. 2002).	A-22
Figure A-5.	Total run size of spring Chinook to the Cowlitz, Kalama and Lewis rivers.....	A-25
Figure A-6.	Natural spawner fall Chinook abundance estimates for the Coweeman and EF Lewis rivers.....	A-27
Figure A-7.	Fall Chinook natural spawner abundance estimates for the NF Lewis River.....	A-27
Figure A-8.	Returns of lower Columbia River hatchery and wild fall Chinook stocks.	A-28
Figure A-9.	Hatchery returns of spring Chinook to the Washington lower Columbia River hatcheries.	A-31
Figure A-10.	Returns of adult spring Chinook (escapement and harvest) to Washington lower Columbia River tributaries, 1980–2000.	A-32
Figure A-11.	Hatchery releases of fall Chinook to the Grays, Elochoman, Cowlitz, and Toutle River basins, 1967–2002.....	A-33
Figure A-12.	Hatchery releases of fall Chinook to the Kalama, Lewis, Washougal, and Little White Salmon River basins, 1967–2002.	A-34
Figure A-13.	Hatchery returns of fall Chinook to Washington lower Columbia River hatcheries.....	A-35
Figure A-14.	Hatchery returns of fall Chinook to Columbia River subbasin, 1950–2002.....	A-36
Figure A-15.	Total commercial catch (excluding the Willamette River) of spring Chinook in the lower Columbia fisheries.	A-37
Figure A-16.	Harvest (sport and commercial) of spring Chinook in the lower Columbia fisheries. ...	A-38
Figure A-17.	Spring Chinook fishery exploitation over time. Harvest dominated by Columbia River commercial fisheries until 1950s.	A-39
Figure A-18.	Harvest of spring Chinook in the Columbia River mainstem from 1985–2002.	A-41
Figure A-19.	Total sport harvest of adult spring Chinook in the Cowlitz, Kalama, and Lewis rivers..	A-42
Figure A-20.	Tributary tribal harvest and tribal distributions of spring Chinook in the Wind and Little White Salmon rivers, 1982–2002.....	A-47
Figure A-21.	Total sport harvest of adult spring Chinook in lower Columbia tributaries and Lower Gorge Tributaries fisheries.	A-48
Figure A-22.	Commercial harvest of lower river wild and lower river hatchery stock fall Chinook in the Columbia River.	A-50
Figure A-23.	Sport harvest of lower river wild and lower river hatchery fall Chinook stocks in the Columbia River.	A-50
Figure A-24.	Total harvest of fall Chinook in the Columbia River from 1980–98.	A-51
Figure A-25.	Approximate fall Chinook fishery exploitation rate over time. Primarily Columbia River commercial harvest until ocean fishery expansion in 1950s.....	A-52
Figure A-26.	Chinook salmon harvest in PSC fisheries in Alaska and Canada, by area, 1987–2002. .	A-54
Figure A-27.	California, Oregon, Washington, and treaty Indian ocean commercial troll landings (in thousands of fish) by month, 1976–2002.	A-57
Figure A-28.	Total value and price per pound (in 2002 dollars) for ocean troll Chinook landings in California, Oregon, and Washington 1979–2002.....	A-58
Figure A-29.	California, Oregon, and Washington ocean recreational salmon effort (in thousands of angler trips) by boat type, 1979–2002.....	A-59
Figure A-30.	Commercial and subsistence harvest of fall Chinook in the Columbia River from 1970–2001.....	A-64

Figure A-31.	Number of adult Chinook landed during early fall, late fall, and treaty Indian mainstem Columbia River seasons.....	A-66
Figure A-32.	Number of days fished during early fall, late fall, and treaty Indian mainstem Columbia River commercial seasons, 1982–2001.....	A-67
Figure A-33.	Buoy 10 fishery recreational catch and combined Oregon and Washington angler catch of Chinook on the lower Columbia River, 1982–2000.	A-67
Figure A-34.	Fall Chinook tributary harvest rate in the NF Lewis, Grays, and Elochoman Rivers, 1988–99. Harvest rate equals sport catch divided by run size at tributary mouth.	A-68
Figure A-35.	Fall Chinook tributary harvest rate in the Kalama, Cowlitz, Toutle, and Washougal Rivers, 1988–99. Harvest rate equals sport catch divided by run size at tributary mouth.	A-69
Figure A-36.	Status of historical spring Chinook populations among lower Columbia River subbasins.	A-71
Figure A-37.	Status of historic bright late fall Chinook salmon populations among lower Columbia River subbasins. Extinction risks are based on viability scores.	A-73
Figure A-38.	Status of historical tule fall Chinook salmon populations among lower Columbia River subbasins. Extinction risks are based on viability scores.	A-74
Figure A-39.	Net effect and relative contribution of potentially manageable impact factors on Chinook salmon in Washington lower Columbia River subbasins.	A-76
Figure A-40.	Approximate spring Chinook fishery exploitation rates over time and allocation of current exploitation rates among fisheries.....	A-78
Figure A-41.	Approximate fall Chinook fishery exploitation rates over time and allocation of current exploitation rates among fisheries. Time series is for tule fall Chinook.	A-79
Figure A-42.	Current, optimal, and historical subbasin productivity and equilibrium numbers inferred for Chinook salmon from stream reach habitat conditions using EDT.	A-82
Figure A-43.	Assumed dam impacts on Washington lower Columbia Chinook populations.	A-83
Figure A-44.	Life cycle fo lower Columbia River coho salmon.	A-86
Figure A-45.	Distribution of historical coho salmon populations among Washington lower Columbia River subbasins.	A-93
Figure A-46.	Columbia River coho ocean and freshwater run size.....	A-96
Figure A-47.	Hatchery releases of coho salmon in the lower Columbia River by basin, 1967–2002. ...	A-100
Figure A-48.	Hatchery releases of coho salmon by brood year from the Little White Salmon NFH, 1980–2002.....	A-101
Figure A-49.	Columbia River adult coho escapement to hatcheries.	A-101
Figure A-50.	Hatchery returns of adult coho salmon in the Cowlitz River basin by hatchery and stock type, 1960-2002.	A-102
Figure A-51.	Hatchery returns of adult coho salmon in the Lewis River basin by stock type, 1950–2002.....	A-103
Figure A-52.	Hatchery returns of adult coho salmon in the Kalama River basin by stock type, 1951–2002.....	A-103
Figure A-53.	Hatchery returns of adult coho salmon in the Washougal River basin by stock type, 1956–2002.....	A-104
Figure A-54.	Hatchery returns of adult coho salmon in the Grays and Chinook River basins by stock type, 1961–2002.	A-104
Figure A-55.	Hatchery returns of adult coho salmon in the Elochoman River basin by stock type, 1954–2002.....	A-105
Figure A-56.	Coho ocean harvest rate based on Oregon Production Index.....	A-106
Figure A-57.	Approximate coho fishery exploitation rates over time.....	A-106
Figure A-58.	Columbia River Zone 1-5 commercial harvest of early and late stock coho.	A-107

Figure A-59.	Harvest of Oregon Production Index Coho in the ocean and Columbia River, 1970–2000.....	A-109
Figure A-60.	Commercial and subsistence harvest of coho salmon in the Columbia River from 1970–2001.....	A-110
Figure A-61.	Recreational harvest of coho salmon in the Lower Columbia River for 1970–2001...	A-111
Figure A-62.	Recreational harvest of Columbia River early/late stock adult coho in Columbia River estuary.....	A-111
Figure A-63.	Recreational harvest of Columbia River early/late stock coho in Washington tributaries of Columbia River.	A-112
Figure A-64.	Coho salmon ocean fisheries harvest and ocean escapement of the primary coho stock components within the OPI area, 1970–2002.	A-117
Figure A-65.	Exploitation rate of ocean fisheries on OPI area coho stocks and percent of total OPI area production accounted for by coho ocean escapement to the Columbia River.....	A-117
Figure A-66.	California, Oregon, Washington, and treaty Indian ocean commercial troll landings (in thousands of fish) by month, 1976–2002.	A-119
Figure A-67.	Total value and price per pound (in 2002 dollars) for ocean troll coho landings in California, Oregon, and Washington, 1971–2002.....	A-120
Figure A-68.	Ocean recreational coho catch 1979–2002.	A-121
Figure A-69.	California, Oregon, and Washington ocean recreational salmon effort (in thousands of angler trips) by boat type, 1979–2002.....	A-122
Figure A-70.	Current status of historical demographically-independent lower Columbia River coho salmon populations.	A-128
Figure A-71.	Net effect and relative contribution of potentially manageable impact factors on coho salmon in Washington lower Columbia River subbasins.	A-130
Figure A-72.	Approximate coho salmon fishery exploitation rates over time and allocation of current exploitation rates among fisheries.....	A-132
Figure A-73.	Current, optimal, and historical subbasin productivity and capacity inferred for coho salmon from stream reach habitat conditions using EDT.....	A-135
Figure A-74.	Life cycle of Columbia River Chum salmon.	A-139
Figure A-75.	Historical demographically independent chum salmon populations in the lower Columbia River ESU (Myers et al. 2002).....	A-146
Figure A-76.	Age composition of natural spawning chum salmon.....	A-148
Figure A-77.	Chum escapement fish/mile peak counts for combined Washington index tributaries.	A-150
Figure A-78.	Fish-per-mile index area counts in Grays River, Hamilton Creek, and Hardy Creek....	A-151
Figure A-79.	Distribution of Lower Columbia chum spawning populations in 2002.....	A-152
Figure A-80.	Total historical hatchery releases of chum salmon into the lower Columbia basins. .	A-154
Figure A-81.	Chum salmon hatchery returns in the lower Columbia River by hatchery, 1957–2002.....	A-155
Figure A-82.	Commercial landings of chum salmon in the Columbia River from 1938–2002.	A-156
Figure A-83.	Fishery exploitation rates over time. Significant Columbia commercial harvest until 1950s.	A-157
Figure A-84.	Average harvest of Chinook, coho, sturgeon, and chum in late fall (mid September to November), 1993-2001.	A-159
Figure A-85.	Late fall commercial harvest of chum salmon in the Columbia River, 1970–2002.	A-159
Figure A-86.	Current status of historical demographically-independent lower Columbia chum populations.....	A-161
Figure A-87.	Net effect and relative contribution of potentially manageable impact factors on chum salmon in Washington lower Columbia River subbasins.	A-163

Figure A-88.	Approximate chum fishery exploitation rates over time and allocation of current rates among fisheries.	A-165
Figure A-89.	Current, optimal, and historical subbasin productivity and equilibrium population size inferred for chum salmon from stream reach habitat conditions using EDT.	A-167
Figure A-90.	Summer steelhead life history.	A-171
Figure A-91.	Winter steelhead life history.	A-171
Figure A-92.	Historical demographically independent summer steelhead populations in the Lower Columbia River DPS (Myers et al. 2002).	A-179
Figure A-93.	Historical demographically independent winter steelhead populations in the Lower Columbia River DPS (Myers et al. 2002).	A-180
Figure A-94.	Counts of total wild and hatchery summer steelhead trapped at Kalama Falls Salmon Hatchery trap, 1976–2001.	A-185
Figure A-95.	Counts of total wild and hatchery winter steelhead trapped at the Kalama Falls Salmon Hatchery trap, 1976–2001.	A-187
Figure A-96.	Total winter steelhead escapement (Grays River, Skamokawa Creek, Abernathy Creek, Germany Creek, Coweeman River, NF Toutle River, and SF Toutle River combined).	A-187
Figure A-97.	Hatchery releases of summer steelhead in the lower Columbia River by basin, 1982–2002.	A-191
Figure A-98.	Hatchery returns of summer steelhead to the Cowlitz and Kalama River basins by hatchery, 1996–2002.	A-192
Figure A-99.	Hatchery returns of summer steelhead to the Lewis and Washougal River basins by hatchery, 1995–2002.	A-193
Figure A-100.	Hatchery releases of winter steelhead to the Grays, Elochoman, and lower tributaries basins, 1982–2002.	A-195
Figure A-101.	Hatchery releases of winter steelhead to the Cowlitz and Kalama River basins, 1982–2002.	A-195
Figure A-102.	Hatchery releases winter steelhead to the Lewis and Washougal River basins, 1981–2002.	A-196
Figure A-103.	Hatchery returns winter steelhead in the lower Columbia River by hatchery, 1995–2002.	A-197
Figure A-104.	Harvest of summer steelhead in the Columbia River from 1938–2002.	A-198
Figure A-105.	Summer steelhead sport catch harvest (combined wild and hatchery) in Washington lower Columbia basin tributaries.	A-199
Figure A-106.	Steelhead fishery exploitation rate over time. Columbia commercial harvest significant until prohibited in 1975.	A-201
Figure A-107.	Lower Columbia River sport catch by steelhead index group, 1969–2002.	A-203
Figure A-108.	Lower Columbia River sport catch by area, 1969-2000.	A-203
Figure A-109.	Wild steelhead harvest rates for summer and winter steelhead in the Kalama basin, 1976–2000.	A-204
Figure A-110.	Steelhead harvest in treaty Indian fisheries by season, 1970–2002.	A-206
Figure A-111.	Harvest of winter steelhead in the Columbia River from 1953–99.	A-209
Figure A-112.	Steelhead fishery exploitation rate over time. Columbia commercial harvest significant until prohibited in 1975.	A-210
Figure A-113.	Distribution of historical summer steelhead populations among lower Columbia River subbasins. Extinction risks are based on viability scores rather than modeled risks. .	A-215
Figure A-114.	Current status of historical demographically-independent lower Columbia winter steelhead populations.	A-216
Figure A-115.	Net effect and relative contribution of potentially manageable impact factors on steelhead in Washington lower Columbia River subbasins.	A-218

Figure A-116. Approximate steelhead fishery exploitation rates over time and allocation of current exploitation rates among fisheries..... A-220

Figure A-117. Current, optimal, and historical subbasin productivity and capacity inferred for steelhead from stream reach habitat conditions using EDT..... A-223

Figure A-118. Assumed dam impacts on Washington lower Columbia steelhead populations. A-224

Figure A-119. Bull trout life history..... A-227

Figure A-120. Distribution of historical bull trout populations among lower Columbia River subbasins. A-233

Figure A-121. Annual peak counts of bull trout spawners observed in Cougar Creek, 1979–2003. . A-235

Figure A-122. Spawning population estimate of bull trout in Swift Reservoir, 1994–2003 (source: Dan Rawding and John Weinheimer, WDFW). A-236



A.1. Chinook Salmon

Chinook salmon (*Oncorhynchus tshawytscha*), also commonly referred to as king, spring, tye, or quinnat salmon, is the largest of the Pacific salmon (Netboy 1958). The species distribution historically ranged from the Ventura River in California to Point Hope, Alaska in North America, and in northeastern Asia from Hokkaido, Japan to the Anadyr River in Russia (Healey 1991). Other Chinook salmon have been reported in the Mackenzie River area of northern Canada (McPhail and Lindsey 1970). Of the Pacific salmon, Chinook salmon exhibit the most diverse and complex life history strategies.

Chinook salmon generally follow one of two freshwater cycles: stream or ocean type. After emerging from the gravel, stream-type Chinook salmon reside in fresh water for a year or more before migrating to the ocean (Figure A-1). Ocean-type Chinook migrate to the ocean within their first year (Figure A-2). These two types of Chinook salmon have different life history traits, geographic distribution, and genetic characteristics. Ocean-type behavior and life history strategy is regarded as a response to limited carrying capacity of the freshwater environment of less productive streams, such as smaller watersheds, glacially scoured rivers, and systems with periodic flooding. Ocean-type Chinook salmon occur primarily in coastal waters south of the 55th parallel, in Puget Sound, in the lower reaches of the Fraser and Columbia Rivers as well as California's Central Valley (Gilbert 1913, Rich 1920, Healey 1983). Stream-type Chinook emigrate as juveniles during their second, or more rarely, third year. This extended freshwater residency is characteristic of Chinook that inhabit more productive watersheds where conditions are more stable, and water flows are not subject to dramatic changes. Since stream-type Chinook enter marine waters at a larger size, they are not as dependent on estuaries as ocean-type Chinook for juvenile growth. In addition, stream-type Chinook make more use of the open ocean environment far from coastal waters. Stream-type Chinook populations are generally more predominant in waters north of the 55th parallel and in headwaters of the Fraser and Columbia rivers (Healey 1991).

Chinook in the lower Columbia River are further classified as fall or spring Chinook depending on adult migration timing. Fall Chinook dominate in the Washington tributaries of the lower Columbia River, though several tributaries also support spring Chinook. Today, the once abundant natural runs of fall and spring Chinook have been largely replaced by hatchery production. Although large Chinook runs continue to return to many of their natal streams, they are mostly sustained by hatchery production with few sustained, naturally reproducing, native populations.

A.1.1. Life History and Requirements

Like other Pacific salmon, the life history of Chinook involves spawning, incubation, and emergence in freshwater, migration to the ocean, and subsequent initiation of maturation and return to fresh water.

Upstream Migration Timing

The entry timing of individual runs into freshwater has evolved over thousands of years. Adult migrations were historically synchronized to streamflow characteristics and water temperatures in a particular river system. Freshwater entry and spawning timing are generally related to local temperature and water flow regimes (Miller and Brannon 1982). Migration timing was cued to the local thermal regime so that adults would reach spawning sites and deposit eggs in time to ensure that fry emerged during the following spring at a time when river or estuary productivity was sufficient for juvenile survival and growth. During lower flows, waterfalls, sandbars and other barriers can restrict migration. After entering freshwater, most fall salmon have a limited time to migrate and spawn; in some cases, as little as 2 to 3 weeks and delays can result in pre-spawning mortality or spawning in a sub-optimum location.

Three different major runs of Chinook salmon presently return to Washington tributaries of the lower Columbia River. Adult spring Chinook return to the Columbia River at 4 to 5 years of age. They enter the lower Columbia River from March through June, well in advance of spawning in August and September (Figure A-1). Spring Chinook typically spawn near headwater areas where higher gradient habitat exists. Historically, fish migrations were synchronized with periods of high rainfall or snowmelt to provide access to upper reaches of most tributaries where fish would hold until spawning (Fulton 1968, Olsen et al. 1992, WDF et al. 1993). Since spring Chinook enter freshwater well before the time of spawning, survival until the spawning period is primarily a function of body fat reserves at the time of freshwater entry.

Fall Chinook begin returning to the lower Columbia River in early to mid-August (Figure A-2). One race of lower Columbia River Chinook salmon are often called tules (pronounced “toolies”) and are distinguished by their dark skin coloration, and advanced state of maturation at the time of freshwater entry. Tule fall Chinook salmon populations may have historically spawned from the mouth of the Columbia River to the Klickitat River. Tule fall Chinook return to the Columbia River at 3 to 4 years of age, although 5-year olds are common in some populations. They enter freshwater from August to September and spawning generally occurs from late September to November, with peak spawning activity in mid-October. Fall Chinook spawn in the Grays River from late September to mid-November, but do not spawn until late October or November in the Washougal River. A later returning component of the fall Chinook salmon run exists in the Lewis River (WDF et al. 1993, Kostow 1995, Marshall et al. 1995).

The other fall race, bright fall Chinook, return to the Lewis River and several Lower Gorge Tributaries and the mainstem Columbia River. Their dominant age class varies by population and brood year, but is typically age 4. They enter the Columbia River in August to October, but spawning occurs in November to January, with peak spawning in mid-November. Because of the longer time interval between freshwater entry and spawning, these fall Chinook salmon are less mature at freshwater entry than tule fall Chinook salmon and are therefore commonly termed lower river ‘brights’ (Marshall et al. 1995) or lower river wild. A naturally produced, bright fall Chinook run also exists in the area immediately downstream of Bonneville Dam, and in the Wind River basin. These fish likely originated from Bonneville and Little White Salmon bright fall Chinook hatchery programs and are not included in the Lower Columbia Chinook ESU.

Spawning

Successful spawning depends on sufficient clean gravel of the right size, in addition to the constant need of adequate flows and water quality. The driving force in redd site selection appears to be the presence of good subgravel flow; this need is likely greater in Chinook than the other species of Pacific salmon. Chinook salmon have the largest eggs and therefore the smallest surface-to-volume ratio of Pacific

salmon. As a result, their eggs are likely more sensitive to reduced dissolved oxygen levels and require a higher rate of irrigation.

Describing typical Chinook spawning habitat is problematic as research has documented a broad range of water depth and velocity characteristics. Chinook have been documented spawning in streams as small as 7-10 ft (2-3 m) wide and only a few centimeters deep and as large as mainstem large rivers such as the Columbia and Sacramento. In addition, velocity measurements at redd sites have ranged from 0.33 ft/sec to 5 ft/sec (10 cm/sec to 150 cm/sec). There is no agreement as to whether depth and velocity characteristics of redd site selection differ between stream- and ocean-type Chinook.

The reported depths at which Chinook eggs are buried in the gravel also varied among researchers. Briggs (1953) reported egg depths of 7.9-14 in (20-36 cm) (average 11 in [28 cm]) for two small California streams. Vronskiy (1972) observed eggs buried from 4 to 31 in (10-80 cm) in the Kamchatka River, although few eggs were buried below 19.7 in (50 cm). The depth at which eggs are buried at a particular spawning site is partly dependent on water flow. Depth of redd excavation is negatively correlated with water velocity in the spawning area (Vronskiy 1972, Neilson and Banford 1983). Presumably, the higher mound in the tailspill of redds in low velocity areas improves subgravel irrigation of the eggs.

Although the measurements are not comparable among studies, the size of redds also appears to vary considerably among Chinook populations. Chapman et al. (1986) measured redd size range of 22-482 ft² (2.1-44.8 m²) for Chinook spawning in the Hanford reach of the Columbia River.

Chinook salmon fecundity also varies within and among populations. Fecundity is correlated with size. However, size explained only 50% or less of the variation in fecundity between individuals within a population. There seems to be an unresolved trade-off between egg size and egg number; consequently, egg size varies more between Chinook individuals than is usual for fishes. Latitudinal differences in fecundity may partly reflect a racial difference between stream- and ocean-type Chinook rather than a latitudinal cline. For example, high fecundity populations near the northern limit of the Chinook's range are all stream-type Chinook while low fecundity populations in the south are mainly ocean-type Chinook. However, if the data are segregated into stream- and ocean-type life histories, there is still a latitudinal cline in fecundity within ocean-type Chinook. In the Columbia River, where fecundity data are available for both stream- and ocean-type Chinook, stream-type Chinook have a greater fecundity than ocean-type, although the difference is not statistically significant (Galbreath and Ridenhour 1966, Healey and Heard 1984).

Incubation and Emergence

Chinook eggs incubate throughout the autumn and winter months. In the lower Columbia River, spring Chinook fry emerge from the gravel from November through March; peak emergence time is likely December and January. Fall Chinook fry generally emerge from the gravel in April, depending on the time of egg deposition and incubation water temperature.

As with other salmonids, water temperature controls incubation time and affects survival. When incubation temperature is held constant, the upper and lower temperature limits for Chinook salmon at 50% pre-hatching mortality is 61°F (16°C) and 36-37°F (2.5-3°C), respectively (Alderdice and Velsen 1978). The time to 50% hatch ranged from 159 days at 37°F (3°C) to 32 days at 61°F (16°C). Development rate and survival were better at low temperatures when water temperature varied with ambient temperature compared to when water temperature was constantly low. Presumably, the better performance reflected greater low water temperature tolerance after initial egg development (Alderdice and Velsen 1978). A simple thermal sum model appears to be adequate for predicting time to hatching (development rate = $468.7/T$, where T is the average temperature in Celsius during

incubation). It is likely that lower Columbia Chinook spawning begins in some locations where water temperatures approach the upper thermal limit (61°F [16 °C]), however, time of exposure to this temperature is likely brief as temperatures are typically dropping during the time of Chinook spawning.

During incubation, clean, well-oxygenated water flow is critical. Eggs often do not survive in gravel choked with sediment (Shaw and Maga 1943, Wickett 1954, Shelton and Pollock 1966). Shaw and Maga (1943) observed that siltation resulted in the greatest mortality when it occurred early in the incubation period. In experimental stream channels, research has established a relationship between egg survival and both percolation rate and dissolved oxygen concentration: egg mortality increases with decreasing percolation rate and increases rapidly when dissolved oxygen concentration drops (Shelton 1955; Gangmark and Bakkala 1960).

Floods can have their greatest impact to salmon populations during incubation, as they can scour salmon eggs from the gravel or deposit sediment over spawning gravels (Wade 2002). Flooding has been documented as an important cause of high mortality of Chinook eggs (Gangmark and Broad 1955, Gangmark and Bakkala 1960).

Estimates of egg to emergent fry survival are problematic because some fish migrate downstream as fry whereas others rear for a variable length of time in the river before migrating downstream. In Fall Creek, California, Wales and Coots (1954) and Coots (1957) found 68-93% mortality from egg deposition to the emergent fry stage; average mortality was 85% and the high mortality estimates (93%) were associated with floods. Lower egg to fry mortality (40%) was observed in a controlled channel in Mill Creek, California (Gangmark and Bakkala 1960). Gravel conditions affect success of emergence. Shelton (1955) found that only 13% of hatched alevins emerged from fine gravel while 80-90% emergence was observed in coarse gravels. Success of emergence from fine gravels was influenced by egg deposition depth; eggs near the surface realized a greater success of emergence.

Dewatering can occur in regulated rivers where discharge is varied to satisfy domestic or industrial water needs but also occurs in natural systems. Becker et al. (1982, 1983) investigated the effects of dewatering on four different stages of Chinook egg development based on accumulated thermal units. Alevins were most sensitive to periodic short-term and single prolonged dewatering; alevin survival was less than 4% in periodic dewaterings of 1 hour or a single dewatering of 6 hours. Cleavage eggs and embryos were the least sensitive to dewatering; embryos apparently suffered no ill effects from daily dewaterings of up to 22 hours over a 20-day period. Because dewatered eggs and embryos remained damp during dewatering, they probably suffered no shortage of oxygen, although metabolic waste elimination may have been a problem.

Freshwater Rearing

Fall Chinook comprise most of the Chinook populations in the lower Columbia River and they exhibit similar life history strategies to those observed in other fall Chinook populations. Fry emergence is generally around April, depending on the time of egg deposition and water temperature. Fry spend 1–4 months in fresh water and emigrate in the summer as subyearlings. A few fall Chinook remain in fresh water until their second spring and emigrate as yearlings (Chapman et al. 1994, Waknitz et al. 1995). Although the timing of emergence and downstream migration differs among lower Columbia fall Chinook, there appears to be little divergence from the strategies of spring emergence and summer emigration. The earliest timing appears to be in the Wind River basin where fry emerge from January to March and emigrate in the spring. The early emigration timing for Wind River fall Chinook may be a function of distance to the estuary, as the Wind River is further from the Columbia River estuary than most other lower Columbia basins. Early and late emergence and late emigration timing occurs in the Lewis River basin; the timing on the Lewis is a function of both late and extended spawn timing of the Lewis bright fall Chinook stock, and warmer winter water temperatures for incubation than most

basins. Consequently, fry emerge from early spring to early summer and seaward emigration occurs in the early to late summer.

Lower Columbia spring Chinook exhibit juvenile life history characteristics similar to those observed in other spring Chinook populations. They have more of a tendency to spend one full year in fresh water and emigrate to sea in their second spring than do fall Chinook. However, some stocks migrate downstream from their natal tributaries in the fall and early winter into larger rivers, including the mainstem Columbia River, where they are believed to over-winter before emigration the next spring as yearling smolts.

Although there is some variation in timing, all populations of Chinook appear to display similar migratory behavior. At the time of emergence, there is an extensive downstream dispersal of fry, although some fry are able to take up residence at the spawning site. For populations that spawn close to tidewater, this downstream dispersal carries fry to estuarine nursery areas, whereas in other locations it serves to distribute the fry among suitable freshwater nursery areas (Healey 1991). After spring and fall Chinook fry leave their gravel nests, they generally move to suitable rearing habitat within side sloughs, side channels, spring-fed seep areas and along the outer edges of the stream. These quiet-water side margin and off-channel slough areas are vital for early juvenile habitat (Wade 2002). The presence of woody debris and overhead cover aid in food and nutrient inputs, and provide protection from predators primarily for the first 2 months of freshwater residence. As Chinook fry grow, some gradually move away from the quiet shallow areas to rear in deeper, faster areas of the stream (Lister and Walker 1966, Chapman and Bjornn 1969). This movement to faster water often coincides with summer low flows that can constrain salmonid production.

Later in the spring, there appears to be a second dispersal that carries some populations to the sea or simply redistributes fry within the river system, presumably to suitable summer rearing areas. For those populations that spend a full year in fresh water, there is a third late fall redistribution to suitable overwintering habitat, usually from the tributaries to the river mainstem (Healey 1991). On the other hand, some overwintering juveniles need habitat to sustain their growth and protect them from predators and winter flows. Wetlands, off-channel habitat, undercut banks, rootwads, and pools with overhead cover are important habitat components during this time. During the late spring and fall distributions, fry tend to shift to deeper water and move seaward. The redistributions may punctuate developmental stages as well as achieve more efficient use of freshwater nursery habitat. Fry redistributions may have adaptive value by shortening the length of spring migration for yearling smolts, especially for headwater spawning populations in larger rivers (Healey 1991).

Survival rates from fry to subyearling migrant or fry to yearling migrant are mostly unknown, except for data collected on the Sacramento River by the USFWS (unpublished). Based on the ocean returns of Chinook from the same brood year tagged as fry and smolts, survival from fry to smolt ranged from 3 - 34% for the 1980–82 year classes. These survival rates are similar to those for other Pacific salmon (Foerster and Ricker 1941, Hunter 1959, Parker 1965) so it is reasonable to assume that Chinook in other river systems have similar survival rates. Predators are usually implicated as the principal agent of mortality among fry and fingerling of Chinook and other species; heavy losses to predators have been documented (Foerster and Ricker 1941, Hunter 1959). However, on the Elochoman River, Patten (1971) observed 1-4% predation by sculpins of Chinook released from the Elokomin Hatchery during 1962 and 1963. In this instance, the release of Chinook fingerling occurred during a single night in 1962 and over three nights in 1963; thus, Chinook were only available to predators for a brief period.

Juvenile Migration

The timing of parr-to-smolt transition seems to depend on a number of environmental and genetic traits that maximize individual survival (Myers et al. 1998). Differences in the timing of smoltification

and emigration to the ocean may be affected by distance of migration to the marine environment, stream stability, stream flow and temperature regimes, stream and estuary productivity, and general weather regimes (Myers et al. 1998). Such environmental factors may be the reason why stream-type Chinook—which usually spawn further inland than ocean-type Chinook—appear unable to smolt as subyearlings. Ocean-type fish have been found to exhibit a faster growth rate relative to stream-type fish (Gilbert 1913, Carl and Healey 1984, Cheng et al. 1987).

Ocean-type juveniles enter salt water following one of three distinct strategies. Some fry migrate to the ocean soon after yolk resorption at 1-2 in (30-45 mm) in length (Lister et al. 1971, Healey 1991). In most river systems, however, fry migrate at 60–150 days post-hatching or as fingerling in the late summer or autumn of their first year. When environmental conditions are not conducive to subyearling emigration, ocean-type Chinook salmon may remain in fresh water for their entire first year.

Stream-type Chinook salmon migrate during their second or, more rarely, their third spring. The underlying biological bases for differences in juvenile life history appear to be both environmental and genetic (Randall et al. 1987). Distance of migration to the marine environment, stream stability, stream flow and temperature regimes, stream and estuary productivity, and general weather regimes have been implicated in the evolution and expression of specific emigration timing. Once stream-type Chinook salmon leave freshwater, they usually move quickly through the estuary, into coastal waters, and ultimately to the open ocean (Healey 1983, Healey 1991). Thus, they are often more dependent on freshwater, rather than estuarine, ecosystems.

The majority of fall-run Chinook salmon emigrate to the marine environment as subyearlings (Reimers and Loeffel 1967, Howell et al. 1985, Hymer et al. 1992a, Olsen et al. 1992, WDF et al. 1993). Most lower Columbia fall Chinook exhibit the ocean-type life history, emigrating to saltwater within their first year (Myers et al. 1998). A portion of returning adults whose scales indicate a yearling smolt migration may be the result of extended hatchery-rearing programs rather than of natural, volitional yearling emigration. It is also possible that modifications in the river environment may have altered the duration of freshwater residence (Myers et al. 1998).

In the lower Columbia basin, spring Chinook generally remain in the river for a full year. However, some stocks migrate downstream from their natal tributaries in the fall and early winter into larger rivers, including the mainstem Columbia River, where they are believed to over-winter before outmigration the next spring as yearling smolts. Cowlitz River spring-run Chinook clearly exhibit yearling smolt pattern as revealed by scale analysis of returning adults (Table 5 in Myers et al. 2003). However, the natural timing of lower Columbia spring-run Chinook salmon emigration is likely obscured by hatchery releases of spring-run Chinook salmon juveniles late in their first autumn or early in their second spring (Myers et al 1998, 2003). Age analysis based on scales from naturally spawning spring-run adults from the Kalama and Lewis rivers indicated a significant contribution to escapement by fish that entered saltwater as subyearlings (Hymer et al. 1992a).

Estuary Rearing and Growth

Ocean-type Chinook salmon reside in estuaries for longer periods as fry and fingerlings than do yearling, stream-type Chinook salmon smolts (Reimers 1973, Kjelson et al. 1982, Healey 1991). Rivers with well-developed estuaries, such as the Columbia, are able to sustain larger ocean-type populations than those without (Levy and Northcote 1982). Juvenile Chinook salmon growth in estuaries is often superior to river-based growth (Rich 1920a, Reimers 1971, Schluchter and Lichatowich 1977).

Since ocean-type Chinook salmon spend more time in the estuary, they are more susceptible to changes in the productivity of that environment than stream-type Chinook salmon. Estuaries may be ‘overgrazed’ when large numbers of ocean-type juveniles enter the estuary *en masse* (Reimers 1973,

Healey 1991). The potential also exists for large-scale hatchery releases of fry and fingerling ocean-type Chinook salmon to overwhelm the production capacity of estuaries (Lichatowich and McIntyre 1987). The loss of coastal wetlands to urban or agricultural development may more directly affect ocean-type populations than stream-type populations. For example, Thomas (1983) and Johnson et al. (2003b) have documented substantial loss of marsh and swamp habitat throughout the estuary and lower Columbia River mainstem; further, many researchers (Levy and Northcote 1982, Myers and Horton 1982, Simenstad et al. 1982, Levings et al. 1986, Bottom et al. 1984) have documented that small juvenile salmonids usually occupy shallow, protected habitats such as salt marshes, tidal creeks, and intertidal flats.

Diet of juvenile fall Chinook varies considerably based on fish size and location in the river, estuary, and nearshore habitats (e.g. Craddock et al. 1976, McConnell et al. 1978, Sibert and Kask 1978, Kjelson et al. 1982, Levy and Northcote 1982, McCabe et al. 1983, Bottom et al. 1984, Dawley et al. 1986, McCabe et al. 1986, Bottom and Jones 1990, Sherwood et al. 1990, Healey 1991, Brodeur 1992, Miller and Simenstad 1997, Simenstad and Cordell 2000).). For young Chinook in the lower Columbia River mainstem, Craddock et al. (1976) determined that diptera were the primary prey species during the winter and spring while zooplankton (primarily *Daphnia*) were the major prey item from July to October. Chironomids, *Daphnia*, amphipods (*Eogammarus* and *Corophium spp.*), *Neomysis*, small fish (juvenile herring, sticklebacks, other salmon), and crustacea larvae have all been identified as important food items in estuaries (Healey 1991). Bottom et al. (1984) and Bottom and Jones (1990) reported that young Chinook in the Columbia River estuary primarily ate amphipods (*Corophium*), cladocerans (*Daphnia*), and diptera, with *Corophium* dominant in winter and spring and *Daphnia* dominant in summer. Seasonal changes in diet are typical, however, it is unclear whether this is related to seasonal abundance of food items or a result of diet shifts as Chinook grow.

Growth in the estuary is correlated with food supply. As a result, growth rate varies between estuaries and between years within an estuary (Healey 1982, Neilson et al. 1985). Reported growth rate estimates range from 0.00275 in/d to 0.52 in/d (0.07 mm/d to 1.32 mm/d), although most estimates seem to fall near the range of 0.0197-0.295 in/d (0.5-0.75 mm/d) (Reimers 1971, Fedorenko et al. 1979, Healey 1980, Levy and Northcote 1981, Kjelson et al. 1982, Neilson et al. 1985, Levings et al. 1986). However, it is uncertain whether growth rate estimates are a measure of the true growth rate or are an artifact of sampling bias.

In the Columbia River estuary, subyearling Chinook salmon were captured in every month of the year and were distributed throughout freshwater, estuarine, and marine regions (Bottom et al. 1984). Reimers (1973), working in the Sixes River, Oregon, suggested that estuarine rearing is critical to fall Chinook survival. Subyearling Chinook were one of the most abundant species collected in the Columbia River estuary; Bottom et al. (1984) suggested that subyearling Chinook abundance was partially related to their slow migration through the estuary (i.e. subyearling Chinook were available for long periods of time in a variety of estuarine habitats). For example, subyearling Chinook tagged and released in April and May were captured in the estuary through October (Bottom et al. 1984). Subyearling Chinook moved through the estuary slower than other salmonids; in fact, migration rate appeared to decrease for about half the hatchery groups when they entered the estuary (Bottom et al. 1984). Generally, juvenile hatchery subyearling Chinook released further upstream in the basin migrated at a faster rate than juveniles released lower in the system (Bottom et al. 1984). Subyearling Chinook abundance was highest in the spring and summer months; during spring and summer, subyearling Chinook were most frequently associated with water column and nearshore habitats while in the winter, they were more frequently associated with nearshore, shoals, and bay habitats (Bottom et al. 1984). Subyearling Chinook represented 68% of the total catch of juvenile salmonids in the estuary (Bottom et al. 1984).

Recent sampling of juvenile salmonids in the Columbia River plume has started to illustrate patterns of habitat use by salmonids in the plume and nearshore ocean habitats (Fresh et al. 2003), although

limited years of data are currently available. First, juvenile salmon distance offshore appears to be positively related to river flow as measured at Bonneville Dam; generally, Chinook and coho salmon yearling were captured further offshore in the plume environment as river flow increased (Fresh et al. 2003). Second, preliminary evidence suggests that some juvenile salmonids (chum, steelhead, and yearling coho) may preferentially utilize the plume front compared to other areas in the plume or adjacent ocean habitats, although this did not appear to be the case for yearling Chinook salmon (Fresh et al. 2003). Although reasons for the apparent preference to the plume front are not clear, this area may be a more productive habitat than elsewhere in the plume and adjacent ocean.

Ocean Migrations

Ocean migrations of Chinook salmon extend well into the North Pacific Ocean. Chinook salmon tend to be widely distributed and run deeper (to 110 m) than other salmon species (Major et al. 1978). Most Chinook salmon remain at sea from 1 to 6 years (more commonly 2 to 4 years). Early maturing males returning to freshwater after 1 year at sea are commonly known as jacks. A small number of yearling males mature in fresh water or return after 2 or 3 months in salt water (Rutter 1904, Gilbert 1913, Rich 1920a, Mullan et al. 1992).

Ocean migratory pattern differences between and within ocean- and stream-type Chinook salmon stocks may be partly responsible for different fluctuations in abundance. They may also reflect long-term geographic and seasonal differences in marine productivity and estuary availability. In addition, differences in the ocean distribution of specific stocks may be indicative of resource partitioning and may be important to the success of the species as a whole. Current migratory patterns may have evolved as a balance between the relative benefits of accessing specific feeding grounds and the energy expenditure necessary to reach them. If the migratory pattern for each population is, in part, genetically based, then the efficiency with which subsequent generations reach and return from their traditional feeding grounds will be increased (NMFS 1998).

Actual oceanic migratory patterns are difficult to discern, especially in the vast marine areas where no fisheries are prosecuted and, hence, no tagged fish are recaptured. Coded-wire tag (CWT) data can help elucidate oceanic migrations, at least in areas where fisheries occur. Myers et al. (1998) stated that CWT recoveries of Chinook from the lower Columbia River ESU (ocean-type) generally indicate a northerly ocean migration route, but with little contribution to Alaskan fisheries. For several specific examples, CWT recovery indicates that: Grays River Hatchery fall Chinook are harvested primarily in southern British Columbia (51%), Columbia River (25%), and Washington ocean (12%) fisheries; Cowlitz River Salmon Hatchery fall Chinook are harvested primarily in Washington ocean (30%), British Columbia (21%), Alaska (15%), Cowlitz River (11%), and Columbia River (8%) fisheries; and Kalama Hatchery fall Chinook are harvested primarily in Alaska (38%), British Columbia (36%), Columbia River (14%), and Washington ocean (6%) fisheries. These three example stocks demonstrate that lower Columbia fall Chinook can range far to the north and that the distribution is rather variable among stocks.

While collecting samples for genetic analysis of oceanic mixed-stock harvest from 88 locations extending from British Columbia to northern California, Utter et al. (1987) found that Columbia River tule fall Chinook tended to be caught in the coastal waters of Washington, while upriver brights tended to be caught in Alaska and British Columbia commercial harvest.

A.1.2. Distribution

During the last 10,000 years, flow, water chemistry and physical features of specific habitats have shaped the characteristics of Chinook salmon populations in the lower Columbia basin (Miller 1965).

Since physical conditions varied between the different lower Columbia River tributaries, Chinook once returned to individual spawning sites over a longer period than they do today. Chinook returning to hatcheries were originally divided into race based on time of arrival at the hatchery. Fish arriving before July 31, were categorized as spring Chinook and after that date as fall Chinook (Senn, H. 1993). This method, however, ignored the entry time of summer Chinook adults. As a result, summer Chinook have been mixed with both spring and fall races.

Fall Chinook were predominant in the lower Columbia, with runs returning to the Cowlitz, Toutle, Coweeman, Lewis, Kalama, Chinook, Grays, Elochoman, Washougal, Big White Salmon and Little White Salmon rivers, as well as to some smaller Washington-side tributaries of the lower Columbia River (Figure A-3). Chinook populations in many of these tributaries began declining by the early 1900s because of overharvest and poor land use practices. The Big White Salmon River (Rkm 270) supported runs of Chinook salmon prior to the construction of Condit Dam (Rkm 4) in 1913 (Fulton 1968). Although some fall-run salmon spawning occurs below Condit Dam, there have been substantial introductions of non-native stocks (WDF et al. 1993), and the persistence of a discrete native stock is unlikely. Fall-run fish from the Big White Salmon River were used to establish the nearby Spring Creek National Fish Hatchery (NFH) in 1901 (Hymer et al. 1992a). Spring Creek NFH is one component of the extensive hatchery system in Washington and Oregon producing fall Chinook salmon (Howell et al. 1985). Among other fall-run populations, a later returning component of the fall Chinook salmon run exists in the Lewis and Sandy rivers (WDF et al. 1993, Kostow 1995, Marshall et al. 1995). Because of the longer time interval between freshwater entry and spawning, Lewis and Sandy river fall Chinook salmon are less mature at freshwater entry than tule fall Chinook salmon and are commonly termed lower river 'brights' (Marshall et al. 1995).

Historically in Washington, spring Chinook returned to the Cowlitz, Lewis, Kalama, and Big White Salmon rivers (Figure A-4). The Cowlitz, Kalama, Lewis, Clackamas, and Sandy rivers presently contain both spring and fall runs, while the Big White Salmon River historically contained both spring and fall runs but presently only contains fall-run fish (Fulton 1968, WDF et al. 1993). The Klickitat River probably contained only spring-run Chinook salmon due to falls that blocked access to fall-run Chinook salmon during autumn low flows (Fulton 1968). The spring run on the Big White Salmon River was extirpated following construction of Condit Dam (Fulton 1968), while a variety of factors may have caused the decline and extinction of spring-run Chinook salmon on the Hood River (Nehlsen et al. 1991, Kostow 1995). Dams have reduced or eliminated access to upriver spring Chinook spawning areas on the Cowlitz, Lewis, Clackamas, Sandy, and Big White Salmon rivers.

A.1.3. Genetic Diversity

Utter et al. (1989) examined allozyme variability at 25 polymorphic loci in samples from 86 Chinook populations extending from the Skeena River, British Columbia, to the Sacramento and San Joaquin Rivers, California. Their cluster analysis of genetic distances (Nei 1972) indicated the existence of nine genetically distinct regional groups of populations. Three groups were located in the Columbia River basin: lower Columbia River and its tributaries, populations above Bonneville Dam (except the Snake River), and the Snake River.

Schreck et al. (1986) examined allele frequency variability at 18 polymorphic loci to infer genetic relationships among 56 Columbia River Basin Chinook salmon populations. A hierarchical cluster analysis of genetic correlations between populations identified two major groups. The first contained spring Chinook salmon east of the Cascade Mountains and summer Chinook in the Salmon River. This group contained three subclusters:

1. wild and hatchery run spring Chinook salmon east of the Cascades,

2. spring run Chinook in Idaho, and
3. widely scattered groups of spring Chinook in the White Salmon River Hatchery, the Marion Forks Hatchery, and the Tucannon River.




Figure A-3. Historical demographically independent fall Chinook salmon populations in the lower Columbia River ESU (Myers et al. 2002).



Spring Chinook

Figure A-4. Historical demographically independent spring Chinook salmon populations in the lower Columbia River ESU (Meyers et al. 2002).

A second major group consisted of spring Chinook salmon west of the Cascade Crest, summer fish in the upper Columbia River, and all fall-run fish. Three subclusters also appeared in this group:

1. spring- and fall-run Chinook salmon in the Willamette River,
2. spring- and fall-run Chinook salmon below Bonneville Dam, and
3. summer- and fall-run Chinook salmon in the upper Columbia River.

Winans (1989) estimated levels of gene diversity with 33 loci for spring, summer, and fall run Chinook salmon at 28 localities in the Columbia River Basin. Fall-run Chinook tended to have significantly greater levels of gene diversity than both spring and summer Chinook salmon.

Waples et al. (1991) examined 21 polymorphic loci in samples from 44 populations of Columbia River Basin Chinook salmon. An unweighted pair group method with arithmetic mean (UPGMA) tree of Nei's (1978) genetic distances between samples showed three major clusters of Columbia River Basin Chinook salmon: 1) Snake River spring and summer Chinook salmon, and mid- and upper Columbia River spring Chinook salmon, 2) Willamette River spring Chinook salmon, and 3) mid- and upper Columbia River fall and summer Chinook salmon, Snake River fall Chinook salmon, and lower Columbia River fall and spring Chinook salmon.

In the NMFS status review, geneticists analyzed a set of allele frequencies for 31 loci in 55 samples from the Columbia and Snake rivers to depict population structure among these drainages. Samples in this analysis were separated into two distinct clusters: ocean-type populations and stream-type populations; except for a sample of spring Chinook salmon from the Klickitat River, which was

genetically intermediate between the two clusters. Results showed that additional genetic population structure was apparent within these two life history types. Within ocean-type Chinook salmon, samples of spring and fall Chinook salmon from the lower Columbia River were distinct from all inland samples. Furthermore, lower Columbia River spring-run fish were genetically more closely allied with nearby fall-run fish in the lower Columbia River than with spring-run fish in the Snake and upper Columbia rivers (Myers et al. 1998).

Taken together, the results of these studies indicate that the timing of Chinook salmon returns to natal rivers is not necessarily consistent with genetic subdivisions. For example, spring Chinook populations in the Snake, Willamette and lower, mid, and upper Columbia rivers were genetically distinct from each other, but had similar run timings. In addition, lower Columbia River tule fall Chinook fish and upper Columbia River bright fall Chinook have similar run timings, but were genetically distinct from one another. Conversely, spring and fall Chinook in the lower Columbia River have different run timing, but were genetically similar (NMFS 1998). The large genetic groupings seem to be driven by geographic isolation more than run timing. Utter et al. (1989) stated that their clustering or gene diversity analyses did not support the concept that Chinook salmon adult run times represented distinct 'races' with separate ancestries, rather that genetic divergence into temporally distinct runs tended to occur within an area from a common ancestry.

A.1.4. ESU Definition

The lower Columbia River Chinook salmon ESU includes all native populations from the mouth of the Columbia River to the Cascade Crest, excluding populations above Willamette Falls (Myers et al. 1998, 2003). Celilo Falls, which historically may have presented a migrational barrier to Chinook salmon under certain flow conditions, is the eastern boundary of the ESU. Exclusions from the ESU are stream-type spring Chinook found in the Klickitat River (mid-Columbia ESU) and the introduced Carson spring Chinook. Tule fall Chinook from the Wind and Little White Salmon rivers are included in the ESU, but introduced bright fall Chinook salmon populations in the Wind, White Salmon, and Klickitat rivers are not included. Information suggests that spring Chinook in the Clackamas and Sandy rivers are predominantly introduced Chinook from the Willamette River ESU and are probably not representative of spring Chinook historically found in these two rivers.

Chinook populations in this ESU are considered by NMFS to be ocean-type (Myers et al. 1998). However, some spring Chinook populations have a large proportion of yearling migrants. Data for naturally reproducing spring Chinook is limited and scale-based aging data, such as that collected by Hymer et al. (1992) may be biased by yearling hatchery releases. These populations exhibit a range of juvenile life history patterns that appear to depend on local environmental conditions. CWT recoveries for lower Columbia River ESU populations indicate a northerly migration route, but with little contribution to the Alaskan fishery. Populations in this ESU also tend to mature at ages 3 and 4, somewhat younger than populations from the coastal, upriver, and Willamette ESUs. Ecologically, the Lower Columbia River ESU crosses several ecoregions: Coastal, Willamette Valley, Cascades, and East Cascades (Myers et al. 1998).

A.1.5. Life History Differences

The obvious life history difference observed among Chinook in the lower Columbia River basin is the presence of spring- and fall-run Chinook. However, as described above, there is little evidence that spring and fall Chinook in the lower Columbia basin are genetically distinct runs. Both spring and fall Chinook in the region have been considered ocean-type Chinook (i.e. migrate to the ocean during their first summer as subyearlings). However, recent scale analysis of juvenile spring Chinook indicates that most lower Columbia spring Chinook emigrate as yearlings. This analysis is heavily biased by the

abundance of hatchery-released yearling spring Chinook; it is unlikely that native spring Chinook in the lower Columbia have adapted a stream-type life history.

Another difference among lower Columbia fall Chinook is the observed rate of straying among Chinook stocks in different regions. For example, fall Chinook in the Coastal Range tributaries (i.e. Chinook, Grays, and Elochoman basins) have a high rate of straying, perhaps because of the relatively short length of these tributaries and/or because Chinook mainly only use the lower rivers just above tidal influence. On the other hand, Chinook in the western Cascade Range tributaries (i.e. Cowlitz, Kalama, Lewis, and Washougal) exhibit a high degree of spawning site fidelity, potentially because fish returning to larger-sized basins normally have a higher degree of homing fidelity. Of the hatchery releases analyzed in this region, more than 90% of the freshwater recoveries occurred in their natal river basin.

Among spring Chinook populations in the lower Columbia River basin, there is little deviation in the life history strategies described above. There is little evidence documenting naturally produced juvenile spring Chinook stream residence time. Spring Chinook in the region may emigrate in the summer as subyearlings, however, documenting this is problematic when yearling hatchery spring Chinook dominate the emigration.

Although fall Chinook salmon populations are generally thought to be one widely mixed stock as a result of straying and egg transfers between hatcheries (Howell et al. 1985, WDF et al. 1993, Marshall et al. 1995), numerous life history differences can be observed among fall Chinook populations throughout the lower Columbia basin. Many of the differences in life history strategies can be attributed to the presence of wild fish maintaining the historical characteristics of a population. Deviations from the typical life history pattern (described above) are observed in Abernathy/Germany, Cowlitz, NF Lewis, EF Lewis, Lower Gorge Tributaries, and Wind River fall Chinook.

In Abernathy and Germany creeks, sexually mature 1-year old fall Chinook have been found. In the Cowlitz basin, spawning generally occurs from September to November, over a broader time period than most fall Chinook, and peak spawning activity does not occur until the first week in November, which is later than most fall Chinook. The NF Lewis River has sustained a healthy natural population of bright fall Chinook. These fish generally migrate from August through October, over a broader time period than other lower Columbia fall Chinook. NF Lewis River bright fall Chinook typically spawn from October through January, with peak activity in November. This spawn timing is substantially later than most other lower Columbia fall Chinook stocks. Also, the dominant age classes of NF Lewis River bright fall Chinook are 4- and 5- year olds. Furthermore, CWT data indicates that NF Lewis River bright fall Chinook have a more northerly ocean distribution than other fall Chinook from the region. On the EF Lewis River, fall Chinook spawning occurs in two distinct segments; the early segment spawns in October and the late segment spawns from November through January. It is possible that the late segment is related to the bright fall Chinook population on the NF Lewis River.

Dominant age classes of EF Lewis fall Chinook include 3-, 4-, and 5-year olds. In the Wind River, tule fall Chinook range from 2 to 4 years old, with 4-year olds predominating, while Wind River bright fall Chinook range from 2 to 6 year olds, with 5-year old spawners predominating.

Wind River bright fall Chinook likely originated from strays from Bonneville Hatchery and Little White Salmon NFH and are not indigenous to the Wind River. Some upriver bright fall Chinook, spawn from mid-October to late November in the mainstem Columbia below Bonneville Dam. This stock was discovered in 1994 and is considered to have originated from hatchery strays from the Bonneville Hatchery upriver bright fall Chinook program. These are not considered part of the Lower Columbia River Chinook salmon ESU.

A.1.6. Abundance

Spring Chinook

There is widespread agreement that natural production has been substantially reduced over the last century. Chinook salmon in the region have been strongly affected by losses and alterations of freshwater habitat (Bottom et al. 1985, WDF et al. 1993, Kostow 1995). Large runs of spring Chinook returned to the lower Columbia historically, most notably to the upper Cowlitz and upper Lewis basins. Both the Lewis and Cowlitz spring Chinook are identified as depressed by WDFW in SASSI (2002). For example, in 1946, WDF estimated spring Chinook escapement in the Cowlitz basin above the proposed Mayfield Dam site was 9,000 fish; when adjusted for harvest, this escapement represents a total spring Chinook run to the Cowlitz of 32,490 fish (most produced from the Cispus River). From 1962 to 1966, an average of 9,928 spring Chinook were counted annually at Mayfield Dam; from 1978 to 1985, only 3,894 spring Chinook were counted annually at the dam. Historically, spring Chinook were abundant in the upper Lewis basin, especially in the Muddy Fork and upper NF Lewis mainstem, with an estimate of at least 3,000 returning to spawn prior to the completion of Merwin Dam in 1932 (WDF 1951). The Merwin Dam was constructed downstream of the spring Chinook habitat, and by 1950 only a remnant population of spring Chinook (<100) remained. The spring Chinook run to the Kalama may have been significant historically, but by the early 1950s, only a remnant population of spring Chinook (<100) existed in the Kalama. Kalama spring Chinook spawning escapement has averaged 444 fish since 1980 and most spawners are considered first generation hatchery fish.

Spring Chinook continue to return to the Cowlitz, Lewis, Kalama, Wind and Little White Salmon rivers, however these runs are almost entirely from hatchery production. Total runs (i.e. escapement plus catch) to the Cowlitz, Lewis, and Kalama rivers have ranged from 3,000 to 36,900 during 1980–2002 (Figure A-5; WDF 1951).




Figure A-5. Total run size of spring Chinook to the Cowlitz, Kalama and Lewis rivers.

In the Lewis River, the naturally spawning spring Chinook population is considered healthy based on escapement trends (WDF/WDW 1993), but some research suggests that the native Lewis River spring Chinook run is extinct (Myers et al. 1998) and that most natural spawners are resulting from hatchery programs. The Cowlitz River now produces very few spring Chinook from natural spawning (average escapement of 338 fish since 1980), and these are generally considered hatchery strays (Hillson and

Tipping 2000, cited in Wade 2000). The Kalama River spring Chinook population is considered healthy, but shows signs of a severe short-term decline (WDF/WDW 1993). All naturally spawning of spring Chinook in the lower Little White Salmon River stopped after the filling of the Bonneville Pool. In addition, hatchery spring Chinook runs exist in the Little White Salmon and Wind rivers, however, spring Chinook were not historically present in these basins. Spring Chinook were historically present in the Big White Salmon River, but were extirpated after the construction of Condit Dam in 1917.

Overall, the number of naturally spawning spring Chinook runs in the Lower Columbia River ESU is very low. The Biological Recovery Team (BRT) established by NMFS to evaluate the status of Chinook was unable to identify any healthy native spring Chinook populations in the ESU. Based on expanded peak fish counts in index areas, the 5-year (1992–96) geometric mean of spring run natural spawning escapement to the Lower Columbia River ESU was 11,200 fish. CWT accounting indicates that approximately 68% of natural spawners are first generation hatchery strays. Long-term escapement trends for spring Chinook are positive or stable although short-term trends are negative. The BRT concluded that the pervasive influence of hatchery fish in almost every river in the ESU and the degradation of freshwater habitat suggested that many naturally spawning populations are not able to replace themselves (NMFS 1998).

Fall Chinook

Natural production of fall Chinook has also dropped far below historical levels. Historically, the Cowlitz River was the primary producer of fall Chinook in the lower Columbia River ESU; an estimated 100,000 adults once returned to the Cowlitz basin (WDF 1951). Although little historical information is available on tributary escapement, WDF and WDW estimated that the total Cowlitz run in 1948 was 63,612 fall Chinook, with approximately 14,000 fish spawning above the proposed Mayfield Dam site. From 1961 to 1966, an average of 8,535 fall Chinook were counted annually at Mayfield Dam. The natural spawning escapement goal of 3,000 fish was met in 2002, however, the escapement goal had not been previously met since the late 1980s. In the Coweeman basin, WDF estimated fall Chinook escapement in 1951 as 5,000 fish. Since 1964, Coweeman fall Chinook escapement has averaged 302 fish, although annual spikes in escapement have been observed periodically over the last 15 years (Figure A-6) (WDFW 2002). In the Kalama River basin, Chinook escapement in 1936 was estimated as 20,000 fish, although only 7,000 were allowed to spawn naturally and 13,000 were collected at the Fallert Creek Hatchery (operating since 1895). Fall Chinook spawning escapements in the Kalama have averaged 5,514 fish since 1964. However, most natural spawners are likely first generation hatchery fish.

On the NF Lewis River, annual fall Chinook spawning escapements have averaged 11,232 since 1964; most spawners in this basin are from natural production. The 5,700 fish escapement goal for NF Lewis fall Chinook is met and exceeded in most years (Figure A-7). Other basins in the lower Columbia River historically supported fall Chinook runs of a few thousand fish, including the Grays, Elochoman, and Washougal rivers; escapement to these basins is currently below historical levels and sustained primarily by hatchery fish (although hatchery fall Chinook are no longer released into the Grays River).




Figure A-6. Natural spawner fall Chinook abundance estimates for the Coweeman and EF Lewis rivers.




Figure A-7. Fall Chinook natural spawner abundance estimates for the NF Lewis River.

Today, fall Chinook continue to return to the Cowlitz, Lewis, Kalama, Washougal, Grays, Chinook, and Elochoman rivers, as well as to several smaller lower Columbia tributaries. Only three fall Chinook stocks, the North Lewis River, EF Lewis River, and Coweeman River fall runs, are considered to be of native origin and predominantly natural production. The Lewis River populations are considered healthy based on escapement trend (Wade 2000), however, recent analysis suggests that EF Lewis fall Chinook are depressed based on low spawner escapement levels. Coweeman fall Chinook are also considered depressed based on low spawner escapement levels.

Based on expanded peak fish counts in index areas, the 5-year (1991–95) geometric mean of fall run escapement to the Lower Columbia River ESU was 29,000 natural spawners and 37,000 hatchery spawners. Long-term escapement trends for fall Chinook are mixed, with most larger stocks positive. However, short-term trends are negative.

Lower river hatchery stock is a management unit representing hatchery and natural production of tule Chinook. Lower river wild stock is a management unit representing later-timed wild fall Chinook

production, primarily from the Lewis River. Figure A-8 displays total Columbia River returns (fishery harvest and spawning escapement combined) for these stocks from 1984 to 2002.




Figure A-8. Returns of lower Columbia River hatchery and wild fall Chinook stocks.

A.1.7. Productivity

Spring Chinook

Very little data are available to assess the productivity of spring Chinook in the lower Columbia River. In the absence of data, natural spring Chinook production is believed to be quite low. The Northwest Power and Conservation Council's (NPCC)¹ smolt density model was applied to many systems in the lower Columbia to estimate potential spring Chinook salmon smolt production (NPPC 1989). (The NPPC smolt density model produces optimistic smolt potential estimates compared to the EDT model.) In the Cowlitz basin, the model predicts potential spring Chinook production of 329,000 smolts below Mayfield Dam, 788,400 smolts for the Toutle system, and 1,600,000 smolts for the basin above Mayfield Dam. In the Kalama basin, the model predicts smolt production of 111,192 smolts in the Kalama below Kalama Falls and 465,160 smolts above Kalama Falls. Wind River smolt production was estimated as 157,533 smolts, while the Little White Salmon River can produce an estimated 32,350 smolts. Smolt production estimates were not available for the Lewis River basin. Based on the smolt density model, the lower Columbia basins with existing populations of spring Chinook (except the Lewis) could produce a total of 3,483,635 smolts. The vast majority of the lower Columbia production potential is in habitat upstream of the Cowlitz and Lewis hydro electric projects.

Fall Chinook

Most fall Chinook salmon populations in Washington tributaries to the lower Columbia River are thought to be one widely-mixed stock as a result of straying and egg transfers between hatcheries (Howell et al. 1985, WDF et al. 1993, Marshall et al. 1995). However, very few egg transfers have been

¹ The Northwest Power and Conservation Council (NPCC) was formerly known as the Northwest Power Planning Council (NWPPC)

made to Cowlitz and Kalama hatcheries, and the existing hatchery stocks are assumed to be similar to the original natural spawning populations in those rivers.

Cowlitz River fall Chinook natural spawners are a mixed stock of composite production. Their status was listed as healthy by SASSI in 1993, but current fall Chinook stocks are considered depressed by WDFW (Hillson and Tipping 2000 cited in Wade 2000). Mobrand Biometrics (1999) compared observed and estimated adult wild fall Chinook returns to the Cowlitz River from about 1920 to 1999. Their results show that production, once estimated at 100,000 adults, declined to ~18,000 fish in the 1950s, ~12,000 in the 1960s and recently declined to less than 2,000 fish. An ecosystem diagnosis and treatment (EDT) analysis attributed the extreme loss in major production to mainstem dams that barred fish passage to historical habitat in the upper basin. They also attributed losses in the lower Cowlitz downstream of the Toutle River to major human-caused changes to the river channel, such as dredging, diking, and straightening. The EDT analysis states that "*uncertainty exists with all of the run-size estimates discussed, and the results must be applied with caution; however, the pattern is troubling*" (Mobrand Biometrics 1999).

Natural spawning occurs in the Washougal, Grays, Chinook, and Elochoman drainages, but the majority of returning fall Chinook that spawn naturally are considered to be hatchery strays. There has been a concern regarding releases of Rogue River fall Chinook at Youngs Bay, which are released into the lower Columbia River to increase harvest opportunities, and their documented straying into many tributaries in the lower Columbia River (NMFS 1998). In recent years, ODFW addressed the concern by eliminating Rogue stock releases from Big Creek Hatchery where they tended to home poorly. Rogue stock release now are entirely within Youngs Bay, Oregon where an intensive gill-net fishery exists.

Today, the fall Chinook run in the Lewis River appears to be the only healthy naturally produced population in the lower Columbia River ESU (NMFS 1998). NF Lewis River fall Chinook represent about 80% to 85% of the wild fall Chinook escapement to the lower Columbia River (WDF 1990). In a recent stock status inventory (SASSI 2002), WDFW grouped the lower Columbia River fall Chinook populations into the following categories:

- Healthy - Elochoman, Abernathy, Toutle (Green), Kalama, NF Lewis, Washougal, and Wind (bright),
- Depressed - Grays, Skamokawa, Germany, Mill, Cowlitz, Coweeman, SF Toutle, and EF Lewis, and
- Critical - Wind (tule).

A.1.8. Hatchery Production

Spring Chinook

Spring Chinook salmon populations in the lower Columbia River have been heavily influenced by hatchery programs, which were developed to mitigate for lost spring Chinook production associated with dam construction and other habitat degradation (see Habitat section below). Present spring Chinook salmon populations in the lower Columbia River are primarily produced by hatchery programs. Total releases have changed over the years by basin. In the Lewis and Cowlitz River basins, annual releases have generally been less than 1.5 million smolts. Spring Chinook releases into the Wind River basin are primarily from the Carson National Fish Hatchery (NFH), while releases into the Little White Salmon River basin are primarily from the Little White Salmon NFH, although some releases from the Carson NFH have been made. The Carson NFH has often produced more spring Chinook adults than the Little White Salmon NFH, although production numbers have been similar in recent years. The current

(2003 brood) release goal of yearling spring Chinook in the lower Columbia Washington tributaries totals 5,137,000 (Table A-1).

Table A-1. Current (2003 brood) annual release goals of spring Chinook salmon juveniles (subyearling and yearling) into lower Columbia basins.

Basin	Hatchery	Release Goal	
		Yearling	Subyearling
L. Cowlitz	Cowlitz Salmon Hatchery	967,000	
U. Cowlitz	Cowlitz Trout Hatchery		300,000
Kalama	Kalama Falls/Fallert Creek Hatchery	500,000	
Lewis	Lewis/Speelyai Hatcheries	1,050,000	
Deep River	Cowlitz/Lewis Hatcheries	200,000	
Little White Salmon	LWS/Carson Hatcheries	1,000,000	
Wind	Carson/LWS Hatcheries	1,420,000	
<i>Lower Columbia Total</i>		<i>5,137,000</i>	<i>300,000</i>

The Cowlitz River spring Chinook stock has received only limited transfers of non-native stocks, but is strongly influenced by hatchery-derived fish (WDF et al. 1993). Stocks on the Lewis and Kalama rivers are a composite of the Cowlitz River spring Chinook stock and other lower Columbia and Willamette River spring Chinook salmon stocks (WDF et al. 1993). Numerically, most of the spring Chinook spawning naturally in lower Columbia River tributaries on the Washington side are now hatchery strays (Marshall et al. 1995). All Washington populations of spring Chinook salmon in the lower Columbia River are currently managed as populations of mixed origin (WDF et al. 1993).

Adult returns to the hatcheries below Bonneville Dam (Cowlitz-type spring Chinook) has ranged from a few hundred to nearly 25,000 during 1950–2000. The hatchery returns of upriver spring Chinook stock to the Little White Salmon and Carson hatcheries on the Wind and Little White Salmon rivers ranged from a few hundred to over 20,000 during 1950–2000. Since 1995, returns to the Little White Salmon and Carson hatcheries have exceeded the hatchery returns to Cowlitz, Kalama and Lewis hatcheries (Figure A-9).




Figure A-9. Hatchery returns of spring Chinook to the Washington lower Columbia River hatcheries.

Hatchery-produced spring Chinook provide significant harvest opportunity in the mainstem Columbia and in Cowlitz, Kalama, Lewis, Wind, and Little White Salmon tributary sport fisheries. Hatchery-produced spring Chinook are now adipose fin-clipped to provide selective harvest opportunity in the mainstem Columbia and in Cowlitz, Kalama, Lewis rivers and in the future in Wind and Little White Salmon tributary sport fisheries. Total adult spring Chinook returns to the tributaries below Bonneville Dam (Cowlitz-type spring Chinook) have ranged from 3,100 to 36,900 during 1980-2002 (Figure A-10). The adult returns of Carson-stock spring Chinook to the Wind and Little White Salmon rivers have ranged from about 1,200 to over 46,900 during 1980-2002 (Figure A-10). The adult returns to these five tributaries are believed to be nearly 100% from hatchery-produced smolts.

Cowlitz River spring Chinook are the largest component of the lower Columbia River hatchery spring Chinook stocks. Historically, total (i.e. adults and jacks) spring Chinook hatchery returns to the Cowlitz normally have been greater than 10,000 fish, with a peak return in 1987 of nearly 37,000. However, in recent years, hatchery returns to the Cowlitz have declined to a magnitude similar to spring Chinook returns in the Kalama and Lewis River basins. Meanwhile, hatchery returns of spring Chinook to the Wind and Little White Salmon rivers have increased in recent years. The adult production from the Little White Salmon NFH and Carson NFH has exceeded the adult production from Cowlitz, Kalama and Lewis hatcheries since 1995 (Figure A-10).




Figure A-10. Returns of adult spring Chinook (escapement and harvest) to Washington lower Columbia River tributaries, 1980–2000.

Fall Chinook

Currently, there are 10 hatcheries (WDFW, ODFW, and USFWS) that release fall Chinook salmon into the lower Columbia River ESU in Washington. The current (2003 brood) release goals for Washington hatcheries in the lower Columbia ESU total 35.7 million juveniles (Table A-2).

Table A-2. Current (2003 brood) annual release goals of fall Chinook salmon juveniles (subyearling and yearling) into Washington lower Columbia basins.

Basin	Brood Source	Annual Release Goal	
		Tule	URB
Little White Salmon	LWS/Priest Rapids Hatcheries		2,000,000
Washougal	Washougal Hatchery	4,000,000	
Kalama	Kalama Hatchery	5,000,000	
Toutle/Green	NF Toutle Hatchery	2,500,000	
Cowlitz	Cowlitz Salmon Hatchery	5,000,000	
Abernathy	Abernathy Hatchery	Program discontinued	
Elochoman	Elochoman Hatchery	2,000,000	
Grays	Grays River	Program discontinued	
Chinook	Sea Resources Hatchery	107,500	
Columbia (Bonneville Pool)	Spring Creek Hatchery	15,100,000	
Total		33,707,000	2,000,000

Within the ESU, however there are differences in degree of hatchery influence on the local stocks. The Cowlitz fall Chinook returns are predominately produced from the Cowlitz Salmon Hatchery, however there have been few transfers of outside stocks into the Cowlitz. The Kalama Hatchery stock has also

generally maintained eggs solely from Kalama-origin fish. The North Lewis has maintained a healthy wild component with minimal hatchery influence and Lewis River fall Chinook hatchery production was discontinued after 1985. The EF Lewis and Coweeman fall Chinook populations are at low levels but are not influenced by hatchery production.

Historical releases of most fall Chinook hatchery programs peaked in the late 1970s and 1980s, with over 10 million Chinook released annually in the Grays, Cowlitz, Kalama, Washougal, and Little White Salmon River basins (Figure A-11 and Figure A-12). The highest annual release of fall Chinook in a lower Columbia basin was over 30 million Chinook in the Little White Salmon River in 1978.

Throughout the range of fall Chinook salmon, stocks have often been transferred among watersheds, regions, states, and countries, either to initiate or maintain hatchery populations or naturally spawning populations. The transfer of non-native fish into some areas has shifted the genetic profiles of some hatchery and natural populations so that the affected population is genetically more similar to distant hatchery populations than to local populations (Howell et al. 1985, Kostow 1995, Marshall et al. 1995). However, most fall Chinook salmon releases into the Lower Columbia River ESU originated from stocks within the ESU, although some upriver stocks were propagated as described earlier. Because of extensive mixing of hatchery and wild populations, it is often difficult to determine the proportion of native and non-native hatchery fish released into a given watershed. Transplanted hatchery fish routinely acquire the name of the river system into which they have been released. The majority of fall run Chinook salmon populations in Washington tributaries of the lower Columbia are thought to be essentially one stock, widely mixed as a result of adult straying and egg transfers between hatcheries (Howell et al. 1985, Utter et al. 1989, WDF et al. 1993, Marshall et al. 1995).




Figure A-11. Hatchery releases of fall Chinook to the Grays, Elochoman, Cowlitz, and Toutle River basins, 1967–2002.

Fall Chinook Hatchery Releases in the Kalama, Lewis, Washougal, and Little White Salmon River Basins, 1967-2002




Figure A-12. Hatchery releases of fall Chinook to the Kalama, Lewis, Washougal, and Little White Salmon River basins, 1967–2002.

The majority of natural spawners in the Grays, Elochoman, Cowlitz, Kalama, and Washougal rivers has been of hatchery origin and strays from several lower Columbia River hatcheries are found in these basins (WDF et al. 1993, Marshall et al. 1995). As well, strays from Oregon’s Rogue River fall run Chinook program have been observed in the Elochoman River and Abernathy Creek (WDF et al. 1993, Marshall et al. 1995). However, the release location of this stock has been changed to address this problem. These select area brights are uniquely marked for monitoring and removal at hatchery traps.

Large numbers of upriver bright fall Chinook strays from the Little White Salmon NFH and Bonneville Hatchery programs have been found naturally spawning above Bonneville Dam in the Wind, White Salmon, and Klickitat rivers (WDF et al. 1993). Broodstock for this program was collected by intercepting various upriver bright stocks headed for spawning sites above The Dalles Dam.

Lower Columbia River fall Chinook salmon hatchery stocks continue to comprise the majority of all Chinook salmon in the Lower Columbia River ESU. However, influence of hatchery fish on natural spawning populations in the North Lewis, East Lewis, and Coweeman rivers is thought to be negligible. Returns to lower Columbia River hatchery facilities in Washington typically range from 10,000 to 40,000 adults (Figure A-13).




Figure A-13. Hatchery returns of fall Chinook to Washington lower Columbia River hatcheries.

Adult hatchery returns of lower Columbia hatchery fall Chinook vary annually and this number is affected by numerous factors, including hatchery juvenile releases, smolt-to-adult survival, ocean survival, and harvest rates.

- Cowlitz tule fall Chinook are the largest individual hatchery run with annual returns usually around 5,000 fish and many years with escapement over 10,000 fish (Figure A-14). Tule fall Chinook hatchery returns in the Cowlitz River basin peaked in the early 1970s and again in the late 1980s.
- Kalama and Washougal tule fall Chinook are the next largest hatchery runs in the lower Columbia River (Figure A-14). Kalama tule fall Chinook returns peaked in the early 1970s and the late 1990s, while Washougal tule fall Chinook returns peaked in the late 1980s and late 1990s.
- The Elochoman River has a substantial hatchery return of tule fall Chinook; returns in the Elochoman peaked in the late 1980s and late 1990s (Figure A-14).
- In the Lewis River basin, tule fall Chinook hatchery returns have been relatively low and constant over time (Figure A-14).




Figure A-14. Hatchery returns of fall Chinook to Columbia River subbasin, 1950–2002.

A.1.9. Fishery

Spring Chinook

Before 1976, over 50% of the mainstem Columbia River spring Chinook run was harvested, primarily in April and May. After 1977, target fisheries for upriver spring Chinook were eliminated and, as a result, lower Columbia River commercial fisheries ended by early March and sport fisheries closed before April. Consequently, harvest rates were reduced substantially. No lower Columbia fisheries during the April/May peak of the runs occurred again until 2001 when adipose fin-clipped hatchery adults returned, enabling fisheries to selectively retain marked hatchery fish and release unmarked wild fish. Commercial fisheries began using live capture methods in 2001, with gear changed from gillnet to tangle net web combined with on-board fish recovery boxes. These selective fishery capabilities in the lower Columbia spring Chinook fisheries have increased hatchery harvest opportunity substantially while minimizing harvest mortality on wild spring Chinook.

Lower Columbia River commercial harvest of spring Chinook ranged from 0 to 18,300 fish during 1985–2002; Washington-origin lower Columbia spring Chinook provided a small portion of the catch during the same period (harvest ranged from 0 to 2,200 for lower river stocks other than Willamette; Figure A-15).

The 1985–2002 lower Columbia total harvest of spring Chinook ranged from zero in 1995 to 32,800 in 2002. Fisheries harvest bottomed out during 1994–2000 when Columbia spring Chinook runs crashed, but increased in 2001 when runs improved and again in 2002 when runs continued to improve and selective fisheries were implemented. The mainstem Columbia sport harvest of spring Chinook has exceeded the commercial harvest in the two most recent years (Figure A-16).




Figure A-15. Total commercial catch (excluding the Willamette River) of spring Chinook in the lower Columbia fisheries.




Figure A-16. Harvest (sport and commercial) of spring Chinook in the lower Columbia fisheries.

Spring Chinook Harvest Over Time

Historically, commercial seasons for spring Chinook occurred in the lower Columbia River in winter and spring. The seasonal structure from 1909 to 1942 was fairly constant, with commercial fishing open 270 days each year. Before 1942, spring Chinook harvest rates typically were 50% or greater. However, lower Columbia stocks were harvested at a lower rate than upper river stocks because March and most of April—peak time for lower Columbia spring Chinook—was closed to fishing. Reductions in the commercial season began in 1943. The commercial spring season (late April–May) was first reduced and then in 1975 completely eliminated to protect depressed stocks of upper Columbia River wild spring Chinook. From 1975 to 2001, commercial fishing was closed by early March. In 2002, full fleet selective commercial fisheries were implemented in late February to late March enabling increased harvest of hatchery spring Chinook.

Sport harvest in the mainstem Columbia River was generally concentrated in April until 1975, when the spring sport fishery was closed. The sport fishery closed by mid- to late March until the coming of selective fisheries in 2001. During 2001–2003, the selective April–May sport fishery was significant for harvest of hatchery spring Chinook. As the mainstem Columbia fishery has been restricted, the tributary fisheries have increased in importance. Most harvest of lower Columbia spring Chinook now occurs in the tributary sport fisheries, chiefly in April and May.

Ocean harvest of spring Chinook was far less than the Columbia River harvest until the 1950s, when the ocean commercial fishery grew rapidly in response to reduced commercial opportunity in the coastal rivers and estuaries. The ocean harvest of spring Chinook peaked in the 1970s and, by the 1990s, was significantly reduced. Total harvest of wild spring Chinook significantly reduced after selective fisheries were implemented (Figure A-17).




Figure A-17. Spring Chinook fishery exploitation over time. Harvest dominated by Columbia River commercial fisheries until 1950s. Ocean harvest significant 1960–1990. Sport harvest increased in 1960s. Tributary sport harvest more significant after 1975. Selective harvest in Columbia River beginning in 2001.

Current Spring Chinook Harvest Distribution

Ocean Fisheries

Current harvest impacts to wild lower Columbia spring Chinook are reduced from historical impacts. The majority of harvest-related mortality of wild spring Chinook now occurs in ocean fisheries because they are not selective for hatchery marked fish (whereas most Columbia River fisheries are currently selective for hatchery fish as describe in the next section). Historically, most ocean harvest occurred in Canadian fisheries although Canadian Chinook fisheries have been substantially reduced in recent years. CWT recoveries from 1989-1994 brood year Cowlitz River Hatchery spring Chinook determined the following distribution: Cowlitz River sport (29%), British Columbia (29%), Washington coast (22%), Columbia River (6%), Oregon coast (5%) and Alaska (3%). In the same period, Lewis River Hatchery spring Chinook were distributed to: Lewis sport (69%), Alaska (11%), British Columbia (10%), Washington coast (5%), Columbia River (4%), and Oregon coast (1%). CWT data suggests that upriver spring Chinook are impacted far less by ocean fisheries than are other Columbia River Chinook stocks.

While lower Columbia spring and fall Chinook are both harvested in Pacific Ocean fisheries, spring Chinook are less subject to ocean fisheries harvest than are falls because of the differences in the patterns and timing of their migration. Although mature fish comprise the majority of the fall Chinook catch in the ocean, a significant portion of the spring Chinook catch can be immature fish. The impacts of the Washington ocean harvest typically depend on the abundance levels of Columbia fall Chinook; these drive Washington ocean Chinook quota levels. Additional details are located in Fall Chinook, PSC Fisheries, and PFMC Ocean Fisheries sections.

Future ocean harvest likely will remain similar to levels of recent years (~18%) because of PST abundance-based management agreements and the anticipation of further development of selective fisheries for Chinook (Table A-3). It is noted, however, that lower Columbia spring Chinook are not included directly as a stock to be considered in abundance-based management agreements with

Canada. Harvest impacts in ocean fisheries could be higher than 18% in years when Chinook abundance is high for key Canadian or U.S. fall Chinook stocks. Ocean harvest could potentially be reduced if selective Chinook fisheries were implemented through the PSC and PFMC processes but there are significant technical complexities in implementing selective ocean Chinook fisheries.

Table A-3. Example of lower Columbia spring Chinook harvest exploitation rates under current management.

Fishery	H*	W**	Comment
Alaska	4%	4%	PSC guidelines for Chinook
Canada	9%	9%	PSC abundance-based management
Washington/Oregon/California ocean	5%	5%	Quotas based on fall Chinook abundance
Columbia River	15%	2%	Selective commercial and sport fisheries
Tributary	20%	2%	Selective sport fisheries
<i>Total</i>	53%	22%	Total lower Columbia stocks (Cowlitz, Kalama, Lewis) Wind and Little White Salmon are upriver stock; ocean harvest is negligible, but total harvest may be similar to lower Columbia hatchery spring Chinook because Columbia harvest includes treaty Indian fishery upstream of Bonneville Dam

* H denotes hatchery fish exploitation rate. Columbia River fisheries managed for commercial/sport allocation and hatchery escapement.

** W denotes wild fish exploitation rate. Columbia River fisheries managed to meet ESA standards for wild Willamette and upriver spring Chinook.

In-river Commercial

In the Columbia River, spring Chinook are harvested in non-Indian winter commercial gillnet fisheries. From 1938 to 1973, approximately 55% of upriver spring Chinook runs were harvested in directed Columbia River commercial and sport fisheries. During 1975-2000 (excluding 1977), no lower river fisheries targeted upriver stocks and commercial fisheries focused on Willamette spring Chinook. Recent conservation measures to protect Willamette River spring Chinook required the release of wild Willamette spring Chinook in all freshwater fisheries. Additionally, since 2001 Columbia River sport and commercial fisheries have been able to retain adipose fin-clipped hatchery fish only and must release unmarked wild fish. As a result, a new tangle net commercial fishery was developed in Zones 1-5 that was selective for adipose fin-clipped hatchery spring Chinook. Multiple gear and education requirements were mandatory for all fishery participants. The new regulations were adopted to improve the survival rates of wild fish captured and released during the fishery. (Lower Columbia fishery impacts on wild spring Chinook now come primarily from catch and release handling mortality.) Although upriver wild spring Chinook are retained in treaty Indian fisheries, total impacts to upriver spring Chinook are constrained by ESA impact limits.

A 2001 management agreement negotiated between the *U.S. v. Oregon* parties (states, tribes, federal agencies) and NMFS concerning limitations on ESA-listed upriver and Snake River wild spring Chinook allowed for a 17% total impact rate on ESA-listed upriver spring Chinook and 2% of this impact was allocated to non-Indian fisheries. The 2% non-Indian allocation was further allocated among commercial and sport fisheries in the lower Columbia; at 1.02% for sport, and 0.68% for commercial. The remaining 0.3% was reserved for upper Columbia and Snake River non-Indian fisheries. Spring Chinook are harvested in Zone 6 Indian winter commercial fisheries although sturgeon are the primary target species for the winter fishery. Spring Chinook are harvested annually in both tribal commercial gillnet

and C&S Zone 6 spring fisheries. The focus for tribal spring fisheries is to attain at least 10,000 spring Chinook for ceremonial needs. Since 2001, increased Upper Columbia spring Chinook abundance has enabled significant tribal ceremonial and subsistence harvest as well as commercial harvest (Figure A-18).




Figure A-18. Harvest of spring Chinook in the Columbia River mainstem from 1985–2002.

In-river Sport

Spring Chinook are the focus of considerable recreational fishing effort in Columbia River estuary, mainstem, select area, and tributary fisheries. In recent years, harvest has been selective for adipose fin-clipped hatchery fish. The selective fishery strategy has enabled the mainstem sport fishery to extend into April and May for the first time since 1977. The major hatchery populations in the lower Columbia River contributing to these fisheries include Cowlitz, Kalama, Lewis, Wind, and Little White Salmon spring Chinook. The Wind and Little White Salmon tributary sport fisheries are not yet selective, but are expected to become selective in 2005 when all returning hatchery adults will be adipose fin-clipped.

Substantial spring Chinook sport fisheries have existed in some lower Columbia subbasins. Average annual spring Chinook sport harvest during the late 1970s and early 1980s was 6,410 in the Cowlitz River, 1,149 in the Kalama basin, and 5,504 in the Lewis River. Total annual sport harvest in the Cowlitz, Kalama, and Lewis rivers combined was about 6,000 to 15,000 for the years 1980–93, but has dropped to 3,200 or less since 1994 (Figure A-19). The reduction in sport harvest corresponds to reduction in spring Chinook runs to these rivers beginning in the mid-1990s.




Figure A-19. Total sport harvest of adult spring Chinook in the Cowlitz, Kalama, and Lewis rivers.

Sport harvest is substantial in the Wind and Little White Salmon (Drano Lake) and much larger than the Lewis, Cowlitz, or Kalama in recent years, with some years’ sport harvest exceeding 10,000 fish. Harvest in the Wind and Little White Salmon is shared between the sport fishery and subsistence and commercial harvest by the Yakama Nation.

Spring Chinook In-River Harvest Management Details

Annual spring Chinook fisheries in the mainstem Columbia are planned consistent with a 2001-2005 agreement between the state, federal, and tribal parties to the *U.S. v. Oregon* federal court case. The agreement establishes the total harvest impact limits for ESA-listed upriver origin wild spring Chinook and treaty Indian and non-Indian harvest sharing. The lower Columbia fisheries are also regulated consistent with ESA limits on Willamette wild spring Chinook and sport and commercial allocation of Willamette hatchery spring Chinook. Regulations are being developed to establish ESA limitations on lower Columbia River wild spring Chinook, however, this regulation development process has lagged behind similar processes that established ESA limitations on upriver and Willamette wild spring Chinook.

When entering the Columbia River, spring Chinook have unique migratory characteristics specific to their stocks. Upper and lower Columbia spring Chinook stocks enter the Columbia River at different times. Harvest managers make use of these differences to set different seasons for different stocks so that harvest rates can be adjusted. In both the mainstem Columbia sport and commercial fisheries, as well as the tributary sport fisheries, current Columbia River management employs selective fishing for marked hatchery spring Chinook.

Lower Columbia River spring Chinook stocks can be separated into two groups for in-river fisheries management; lower river spring Chinook (Cowlitz, Kalama, and Lewis river populations in Washington and Willamette River in Oregon), and upriver spring Chinook (Wind and Little White Salmon River populations).

Mainstem Columbia River harvest impacts on Willamette wild spring Chinook average 4.3%, while the Snake River wild limits for lower Columbia fisheries are 1.7% (ODFW and WDFW, 2001). The Willamette spring Chinook migration through the lower Columbia is earlier than lower Columbia River spring Chinook; Snake River spring Chinook are later timed. Therefore, a mid-range impact of approximately

3% is a reasonable expectation for lower Columbia River wild spring Chinook stocks in mainstem Columbia fisheries.

Select Area fisheries for spring Chinook were developed in the mid-1990s along the Oregon shore of the Columbia River, primarily in Youngs Bay. Spring Chinook smolts are released in off-channel areas outside of the normal migration corridor for populations of wild and hatchery spring Chinook and are harvested in subsequent years near the release sites. One site on the Washington side of the Columbia River (Deep River) has had limited success for spring Chinook select area fisheries. The existing Select Area fisheries likely harvest few spring Chinook destined for Washington tributaries of the lower Columbia River basin.

The *U.S. v. Oregon* agreement for spring Chinook management establishes a sliding scale of harvest impact limits for ESA-listed upriver origin wild spring Chinook based on the abundance of wild Snake River spring Chinook. The agreement also establishes treaty Indian and non-Indian harvest sharing (Table A-4). Fisheries that selectively harvest hatchery fish have dramatically reduced the impacts of the non-Indian fishery on wild fish. (treaty Indian fisheries are not limited to hatchery fish.) The lower Columbia fisheries are also regulated consistent with ESA limits on Willamette wild spring Chinook, 20% for 2001 and 15% for 2002 and beyond.

Table A-4. Sliding scale* of harvest impacts on wild upriver spring Chinook based on Snake River wild spring Chinook run size (adapted from the 2001-05 Interim Management Agreement).

Columbia River Mouth Run Size	Snake River Run Size**	Proposed Tribal Harvest Rate	Non-Indian Harvest Rate [§]	Total Harvest Rate	Non-Indian Wild Limited Rate
<25,000	<2,500	5%	<0.5%	<5.5%	<0.5%
25,000	2,500	5%	0.5%	5.5%	0.5%
30,000	3,000	5%	1%	6%	0.5%
40,000	4,000	6%	1%	7%	0.5%
50,000	5,000	7%	1.5%	8.5%	1%
75,000	7,500	7%	2%	9%	1.5%
100,000	10,000	8%	2%	10%	
<i>130,000</i>	<i>13,000</i>	9%	2%	11%	
200,000	20,000	10%	2%	12%	
250,000	25,000	11%	2%	13%	
300,000	30,000	12%	2%	14%	
350,000	35,000	13%	2%	15%	
400,000	40,000	14%	2%	16%	
450,000	45,000	15%	2%	17%	

Italics indicate 2003 pre-season projections; the spring Chinook run forecast at the river mouth is 145,400.

* This scale is applied if the Snake River wild spring Chinook run is $\geq 7.5\%$ of the total run. The limited harvest rate would be used if the Snake River wild forecast is less than 7.5% of the total run.

**If the Snake River wild spring Chinook forecast is less than 10,000, the total harvest rate is restricted to 9% or less. When wild fish harvest rate is restricted to 9% or less, non-Indian fisheries transfer 0.5% harvest rate to treaty Indian fisheries, however, non-Indian fisheries would never go below a 0.5% harvest rate.

[§] If the total forecast is <25,000 or the Snake River forecast is <2,500, the non-Indian harvest rate would be maintained as close to zero as possible while maintaining minimal fisheries targeting other harvestable species.

Non-Indian sport and commercial allocation is based on abundance of upriver wild spring Chinook as well as Willamette hatchery spring Chinook (Table A-5). The 2003 mainstem Columbia River spring Chinook allocation for non-Indian fisheries was guided by five major principles: 1) meet conservation

requirements for wild spring Chinook, including ESA-listed species, 2) manage spring Chinook harvest within the provisions of the *U.S. v. Oregon* management agreement, 3) meet hatchery escapement goals, 4) implement selective fisheries to focus sport and commercial harvest on hatchery fish, and 5) allocate 15% of the non-Indian upriver spring Chinook impacts to sport and non-treaty Indian fisheries upstream of McNary Dam and provide for a lower river fisheries management buffer.

Table A-5. Allocation of non-Indian upriver wild spring Chinook impacts based on Willamette hatchery and upriver wild spring Chinook abundance.

		Willamette Hatchery Fish Run Size		
		<40,000	40-75,000	>75,000
Upriver Fish Run Size* (Impacts)	30-<50,000 (0.85%)	Comm—10% (0.08)** Sport—90% (0.77) [§]	Comm—30% (0.25) Sport—70% (0.60)	Comm—25% (0.21) Sport—75% (0.64)
	50-<75,000 (1.25%)	Comm—40% (0.5) Sport—60% (0.75)	Comm—35% (0.44) Sport—65% (0.81)	Comm—30% (0.37) Sport—70% (0.88)
	>75,000 (1.7%)	Comm—50% (0.85) Sport—50% (0.85)	Comm—40% (0.68) Sport—60% (1.02)	Comm—35% (0.59) Sport—65% (1.11)

Italics indicate the 2003 estimated run sizes and allocation among non-Indian commercial and sport fisheries

* An upriver run size update along with an assessment of upriver impact needs and Willamette allocation will be conducted after mid-April.

** If the sport fishery impact allocation will be used before May 15 and the commercial fishery does not need its entire upriver impact allocation to attain the Willamette allocation or an equitable catch share, commercial impacts may be transferred to the sport fishery.

§ If the sport fishery does not need their entire upriver spring Chinook allocation to continue the fishery through May 15, the remaining sport impacts may be transferred to the commercial fishery for late spring commercial fishing opportunity.

Every year, after annual run size forecasts are available and public input has been received, the Columbia River Compact sets the structure of sport and commercial fisheries to meet allocation policies and fisheries objectives. Initial fishery planning is based on preseason run forecasts, but seasons are adjusted for smaller or larger runs based on dam counts and information about fishery catch rates. Fish run sizes and catches are monitored in-season so that catch does not exceed allowed guidelines.

Commercial harvest constraints resulting from low abundance of wild fish and ESA limitations to protect listed stocks provided much of the motivation for the development of a new fishery. Meanwhile, the recent hatchery practice of marking all hatchery releases with an adipose fin clip gave the fisheries the capability of selecting hatchery fish. Starting in 2000, modifications to gillnet gear (e.g. reducing mesh size to a maximum of 4½ inches) were tested to evaluate their effectiveness: could hatchery fish be retained and wild fish be released and survive? Gear testing indicated that, while the small mesh gill nets could not gill Chinook salmon, they could retain live Chinook salmon by tangling. This meant fish could be retained or released after determining whether they were of wild or hatchery origin.

A 2002 winter season demonstration involved a non-Indian commercial tangle net fishery using 5½ inch maximum mesh size and targeting hatchery spring Chinook salmon. Salmon catches increased throughout the duration of the fishery; Chinook adipose fin mark rate ranged from 42 to 72% and averaged 50% for the season. Chinook catches and impact rates are presented in Table A-6. The steelhead:Chinook ratio decreased over the period of the fishery. Early on, the ratio averaged 2.5:1; during the middle part of the fishery, the ratio averaged 0.9:1; and at the end of the fishery, the ratio averaged 0.4:1. Steelhead mark rate fluctuated between 20 and 50%; season average steelhead mark rate was 40%. A total of 21,600 steelhead were handled and it is possible that some steelhead were handled more than once. Immediate mortality rate for steelhead was estimated at 2%; Most of the

steelhead (84%) handled were released in condition 1 (vigorous, not bleeding). Some steelhead handled may have been summer steelhead, rather than winter steelhead.

Table A-6. Spring Chinook catch and released during the 2002 non-Indian commercial tangle net fishery in the lower Columbia River.

Fishing Period	Spring Chinook Kept				Spring Chinook Released			Upriver Impacts ^a
	Upriver	Willamette River	Other Lower River	Total	Upriver	Other Lower River	Total	
1/7–2/15	19	115	20	154	25	29	54	0.007%
2/25–3/1	175	311	52	538	317	97	414	0.015%
3/4–3/8	302	386	76	764	426	132	558	0.022%
3/10–3/15	1,037	897	205	2,139	1,690	475	2,165	0.082%
3/17–3/22	3,417	1,824	384	5,625	4,967	741	5,708	0.251%
3/24–3/25	1,489	955	190	2,634	2,623	422	3,045	0.123%
3/26–3/27	2,051	744	148	2,943	2,779	253	3,031	0.145%
<i>Season Totals</i>	<i>8,490</i>	<i>5,232</i>	<i>1,075</i>	<i>14,797</i>	<i>12,827</i>	<i>2,149</i>	<i>14,975</i>	<i>0.645%</i>

^a Upriver impacts were derived directly from WDFW fishery monitoring data; impacts are calculated based on the percent of upriver spring Chinook handled during the fishery, total spring Chinook catch for the fishery, upriver spring Chinook run size, and a long-term catch and release mortality factor.

After analysis of this 2002 fishery, objectives for the 2003 tangle net fishery were identified as:

1. provide commercial fishers with an opportunity to harvest their allocation of surplus Willamette hatchery spring Chinook,
2. manage the fishery to remain within ESA-related impact limits for listed upriver and Willamette River wild spring Chinook stocks,
3. improve steelhead condition at capture and reduce steelhead handling and mortality, and
4. maintain adequate spring Chinook catch rate to limit total fishing time.

The selective fishery management for spring Chinook commercial fisheries has increased opportunity and harvest volume compared to recent recent past The 2002 commercial spring Chinook fishery ex-vessel (value), increased from an average of \$686,000 during 1988-1997 to \$1,462,000 in 2002 (Table A-7).

Treaty Indian spring Chinook fisheries occur in the Wind River and in Drano Lake (Little White Salmon) following annual agreement with WDFW regarding sport and Indian catch allocation. The Yakama Nation sets regulations for subsistence fisheries in the Wind River and commercial fisheries in Drano Lake. Washington sets commercial regulations consistent with the tribal regulations. In recent years, the Columbia River Compact has adopted rules allowing Yakama tribal members to sell Drano Lake commercially-caught spring Chinook in Oregon. Yakama Tribes also collect surplus spring Chinook at Carson and Little White Salmon hatcheries for ceremonial and subsistence purposes. The tribal harvest and surplus distribution in these tributaries has increased in recent years in response to larger returns (Figure A-20).

Table A-7. Ex-vessel value (in thousands of dollars expressed in 2002 dollars) of in-river commercial harvest of Columbia River spring Chinook, 1988–2002.

		Oregon		Washington	
		Non-Indian Gill Net	Treaty Indian	Non-Indian Gill Net	Treaty Indian
1988–97	Price per Pound	3.87	3.38	4.43	4.20
	Ex-V. Value	433	2	245	6
1998	Price per Pound	2.75	0	0	4.29
	Ex-V. Value	98	0	0	*
1999	Price per Pound	2.97	0	2.98	4.23
	Ex-V. Value	84	0	*	*
2000	Price per Pound	2.79	2.91	5.01	1.97
	Ex-V. Value	236	2	16	52
2001	Price per Pound	2.67	1.39	3.84	1.28
	Ex-V. Value	594	34	135	283
2002	Price per Pound	2.95	1.21	4.23	1.18
	Ex-V. Value	932	17	295	218

* Less than \$500.

Significant spring Chinook sport fisheries have existed in the lower mainstem and many lower Columbia basins. Sport seasons are set by the Washington Fish and Wildlife Commission and managed and monitored in-season by WDFW. Sport harvest is substantial in the Wind and Little White Salmon (Drano Lake) rivers and is much larger than the Lewis, Cowlitz, or Kalama in recent years, with total sport harvest recently exceeding 10,000 fish (Figure A-21). Harvest in the Wind and Little White Salmon is shared between the sport fishery and subsistence and commercial harvest by the Yakama Nation.

Significant angler effort is expended during Columbia River recreational fisheries, creating significant economic impacts. Recreational fishing effort and angler satisfaction have increased in recent years compared to the 1990s because of hatchery-selective fishing opportunity. There is significant spring Chinook fishing effort in the mainstem Columbia below Bonneville Dam, and the Willamette, Cowlitz, Kalama, Lewis, Wind, and Little White Salmon rivers all support significant tributary sport fisheries.




Figure A-20. Tributary tribal harvest and tribal distributions of spring Chinook in the Wind and Little White Salmon rivers, 1982–2002.




Figure A-21. Total sport harvest of adult spring Chinook in lower Columbia tributaries and Lower Gorge Tributaries fisheries.

Fall Chinook Fishery

Columbia River fall Chinook are harvested in ocean commercial and recreational fisheries from Oregon to Alaska, as well as the Columbia River commercial gill net and sport fisheries. Lower Columbia tule fall Chinook are an important contributor to Washington ocean troll and sport fisheries as well as the Columbia River estuary sport fishery. In the past, harvest rates on fall-run stocks have been moderately high, with an average total exploitation rate of 65% (1982–1989 brood years) (PSMFC 1994). The average ocean exploitation rate for this period was 46%, while the freshwater harvest rate on the fall run has averaged 20%, ranging from 30% in 1991 to 2.4% in 1994.

Currently, ocean and mainstem Columbia River fisheries are managed for Snake and Coweeman River wild fall Chinook ESA harvest rate limits, consequently limiting harvest of other co-mingled Columbia River fall Chinook stocks. Unlike spring Chinook, hatchery fall Chinook are not marked so total harvest rate is the same for hatchery and wild fish. Ocean and mainstem Columbia River fisheries on tule stocks are limited to a 49% harvest rate because of the ESA harvest limits on Coweeman fall Chinook. Columbia River harvest of Snake River fall Chinook is limited to 31.29%, of which 8.25% is non-Indian harvest, and 23.04% is treaty Indian harvest. These ESA harvest limits on Snake River and Coweeman fall Chinook were established in consultation processes between state and tribal governments and NMFS. Coweeman fall Chinook were selected to represent lower Columbia tule fall Chinook stocks because they have not been influenced by hatchery production, and are considered a genetic legacy. These are maximum harvest rates and actual harvest is often less. Annual harvest varies depending on management response to annual Chinook abundance determined in PSC, PFMC, and Columbia River Compact forums. Considerable basin-specific data are available to address harvest effects on, and specific distribution of, distinct Columbia fall Chinook stocks.

Harvest of lower Columbia fall Chinook is managed within four separate, broad stock units:

- Lower River Hatchery (LRH) stock are an earlier spawning component and contain both hatchery and naturally produced fish returning to most of the Washington lower Columbia tributaries.

- Lower River Wild (LRW) stock is primarily produced from the Lewis River and is all naturally produced.
- Upriver Bright (URB) stock is primarily produced in the Columbia Basin upstream of the lower Columbia area, but there are non-listed URB natural spawners present in the mainstem Columbia immediately below Bonneville Dam and in the lower Wind River.
- Bonneville Pool Hatchery (BPH) stock are an earlier spawning hatchery component released at Spring Creek Hatchery upstream of Bonneville Dam with some natural spawning components in tributaries between Bonneville and The Dalles dams.

These three stocks have different migratory characteristics and there are management criteria specific to each stock. Columbia River fisheries are managed based on annual forecasts of abundance for each stock in aggregate. Tributary fisheries are managed based on the annual abundance of returns to the specific tributaries. The harvest of fall Chinook in the Columbia River is subject to *U.S. v. Oregon* Fall Management Agreements regarding Indian and non-Indian allocation, as well as agreements on the allocation of sport and commercial fishing and ESA requirements for listed fall Chinook. Additionally, annual agreements for allocation of harvest between sport and commercial and ocean and Columbia River fisheries are made during the North of Falcon process, a public process aimed at balancing harvest and fishery escapement objectives between ocean and freshwater users. The Columbia River Compact (Oregon and Washington joint regulatory forum) sets specific Columbia River commercial and sport seasons that meet the intent of the annual agreements.

Annual ocean harvest of Columbia River fall Chinook is developed through provisions of the Pacific Salmon Treaty and the Pacific Fishery Management Council process for fisheries off the coasts of Washington, Oregon, and California. Lower Columbia fall Chinook ocean harvest occurs primarily off the coasts of British Columbia and Washington.

Columbia River fall Chinook are an important contributor to ocean fisheries from Oregon to Alaska. The LRH component is the most southerly distributed and the abundance of this stock is a major consideration when setting Chinook harvest levels off the Washington coast. The LRH fish also contribute significantly to Canadian fisheries. LRW and URB components are more northerly distributed in the ocean.

The modern day commercial harvest of lower Columbia fall Chinook peaked during 1987–88 when record fall Chinook numbers returned to the Columbia River. Harvest of lower river hatchery stock (tules) was almost 180,000 adults and lower river wild stock was nearly 19,000 adults (Figure A-22). The commercial harvest of lower river fall Chinook reduced significantly after 1989 and remained low through the 1990s.

Columbia sport harvest of lower river Chinook peaked in 1987–89, with lower river hatchery harvest nearly 33,000 in 1987 and lower river wild harvest nearly 5,000 in 1989 (Figure A-23).

Commercial Catch Zones 1-5 of Lower River Fall Chinook Adults Entering the Columbia River, 1980-2002




Figure A-22. Commercial harvest of lower river wild and lower river hatchery stock fall Chinook in the Columbia River.

Sport Catch of Lower River Wild Fall Chinook Adults Entering the Columbia River, 1980-2002




Figure A-23. Sport harvest of lower river wild and lower river hatchery fall Chinook stocks in the Columbia River.

Fall Chinook Harvest Over Time

Lower Columbia fall Chinook were historically harvested in Columbia River fall fisheries from August to October. Before 1949, Columbia River commercial seasons were open daily during the fall, except for a

closed period from August 25 to September 10. Most harvest was from Columbia River commercial fishing until the 1950s, when ocean fisheries increased in response to reduced Columbia River and coastal estuary commercial fisheries. Ocean harvest peaked in the 1970s, but in the 1990s reduced significantly in response to declines in the abundance of Columbia River tule Chinook. Columbia River mainstem sport fisheries for fall Chinook began increasing in the 1980s, and now the annual mainstem sport harvest of fall Chinook is similar to the commercial fishery. Fall Chinook tributary fisheries advanced in popularity in the 1960s. Most tribal Chinook harvest occurred in a dip net fishery at Celilo Falls, with tribal commercial landings of salmon ranging from 0.8 to 3.5 million pounds annually during 1938–1956. The Celilo fishery ended in 1957 with the inundation of the falls by The Dalles Dam. Commercial fishing in Zone 6 (Bonneville to McNary dams) was closed by state law during 1957–1967. It reopened exclusively for treaty Indian commercial fishing in 1968 following federal court decisions regarding treaty Indian fishing rights. Since 1980, URB fall Chinook have been the primary fall Chinook harvested in the Columbia River, however the harvest of LRH stock has also been very large in some years. The largest harvest of fall Chinook occurred in 1987 (Figure A-24), when a record 872,000 fall Chinook adults returned to the Columbia River.

In general, the approximate fall Chinook fishery exploitation rate over time held steady around 70-80% until the 1990s when fisheries were reduced as a result of ESA-driven management changes (Figure A-25).




Figure A-24. Total harvest of fall Chinook in the Columbia River from 1980–98.

Approximate Fishery Exploitation Rates Over Time Fall Chinook




Figure A-25. Approximate fall Chinook fishery exploitation rate over time. Primarily Columbia River commercial harvest until ocean fishery expansion in 1950s. Northern migration with Canada and Alaskan interception significant in some years. Commercial harvest primarily in September. LRH component important to Washington ocean fisheries. Mainstem Columbia sport harvest increased in 1990s. Tributary sport harvest focus is September.

Current Fall Chinook Harvest Rates and Distribution

The current harvest of lower Columbia fall Chinook is significantly reduced from past harvest levels. Reductions in the Columbia River harvest actually began by the 1950s, but coincided with increased ocean harvest, resulting in relatively high total harvest rates until the 1990s. The current harvest levels average about 45% for the three fall Chinook stocks present in the lower Columbia.

Because of their northerly migration patterns, fall Chinook are harvested in Canada and Alaska fisheries more than other salmonids. For example, the majority of fishery CWT recoveries of 1989-94 brood Cowlitz Hatchery fall Chinook were distributed between Washington ocean (30%), British Columbia (21%), Alaska (15%), Cowlitz River (11%), and Columbia River (8%) sampling areas. Also, CWT recoveries of Kalama fall Chinook 1992-1994 brood indicate the majority of the harvest occurred in British Columbia (36%), Alaska (38%), Washington ocean (6%), and Columbia River (14%) fisheries. Upriver bright fall Chinook stocks also have a northerly migration. CWT data analysis of the 1989-1994 brood years suggests that the majority of the URB fall Chinook harvest occurred in Alaska (24%), British Columbia (23%), and mainstem Columbia River (42%) fisheries. However, tule fall Chinook stocks originating from the Bonneville Pool are more southerly distributed. CWT data analysis of the 1971-1972 brood years from Spring Creek Hatchery indicates that the majority of Bonneville Pool Hatchery fall Chinook stock harvest occurred in British Columbia (28%) and Washington (38%) ocean commercial and recreational fisheries. Canadian interception of Columbia River fall Chinook was reduced beginning in the mid-1990s because of management concerns for depressed Canadian Chinook stocks. Current Canadian harvest is limited by the recent abundance-based management agreement negotiated through the PST process.

In Washington coastal and Columbia River fisheries, the harvest of fall Chinook was reduced in the 1990s because of the reduced abundance of fall Chinook and ESA limitations (Figure A-25). While LRH

stock fall Chinook have rebounded in abundance in recent years, fall Chinook harvest is limited by ESA constraints on LRH natural spawners (Coweeman index) and on Snake River Wild (SRW) fall Chinook. The ESA limits total harvest (combined ocean, Columbia River, and tributary) of Coweeman natural fall Chinook to 49% or less (Table A-8). The ESA restricts southern U.S. ocean harvest of SRW Chinook (a component of the URB stock) to a 30% reduction from the 1989–1993 average harvest rate. The ESA restricts Columbia River harvest of SRW Chinook to 31.29%, allocated at 23.04% for treaty Indian fisheries and 8.25% to non-Indian fisheries.² Ocean, Columbia River, and tributary fisheries are managed to attain a minimum of 5,700 LRW natural spawners to the North Lewis River. Although hatchery fall Chinook are not mass marked, and wild harvest rates are likely similar to hatchery harvest rates, differential harvest can be achieved between fall Chinook stocks depending on management strategies implemented in a given year.

Table A-8. Example of lower Columbia fall Chinook current harvest exploitation and distribution under current management.

Fishery	Tule*	LRW**	URB [§]	Comments
Alaska	3%	10%	10%	PSC abundance-based management
Canada	12%	9%	15%	PSC abundance-based management
Washington/Oregon/California ocean	15%	3%	2%	PSC, ESA, allocation constraints
Columbia River	10%	8%	20%	ESA, allocation, <i>U.S. v. Oregon</i> constraint
Tributary	5%	10%	1%	ESA, escapement goal driven
Total	45%	40%	48%	Wild and hatchery fish rates

*Lower river tule harvest driven by 49% limit for Coweeman fall Chinook

**Lower river wild harvest driven by 5,700 minimum natural escapement to North Lewis

[§]Upriver harvest driven by Snake River wild ESA constraint and *U.S. v. Oregon* Indian /non-Indian allocation agreement

Fall Chinook Harvest Management Details

PSC Fisheries

In southeast Alaska, Chinook salmon are harvested in ocean commercial troll, commercial net, and recreational fisheries. Total southeast Alaska Chinook catch (in numbers of fish) from 1987–2002 has ranged from 155,700 in 1996 to 373,900 in 2002 (Figure A-26).

The spring troll fisheries are designed to increase the harvest of Chinook salmon produced by Alaskan hatcheries by allowing trolling in the small nearshore areas close to the hatcheries where fish concentrate. Although there is no ceiling on the number of Chinook salmon harvested in the spring fisheries, the take of PST-governed Chinook salmon is limited according to the percentage of the Alaskan hatchery fish taken.

Summer and winter troll fisheries primarily harvest PST-governed Chinook salmon and these fish are counted toward the Alaska fisheries allocation. Southeast Alaska commercial net fisheries target fish other than Chinook salmon, but Chinook are harvested incidentally in these fisheries. In the recreational fisheries of southeast Alaska, the harvest of Chinook salmon can be substantial: the recreational fishery harvest in 2002 was 85,200 Chinook salmon, with 27,000 from Alaska hatcheries.

Directed Chinook harvest occurs in numerous fisheries through Canadian PSC-managed waters; Chinook also are incidentally harvested in sockeye-directed fisheries (Figure A-26). Canadian Chinook fisheries are managed through either abundance-based management agreements (AABM) or Individual Stock

² *U.S. v. Oregon* Management Agreement

Base Management (ISBM) limits (Table A-9). Management of each fishery is directed by the abundance of the stocks of concern. Selective fishery practices are used to protect stocks, and these include gear requirements such as single barbless hooks and on-board revival tanks for resuscitating salmon for release.




Figure A-26. Chinook salmon harvest in PSC fisheries in Alaska and Canada, by area, 1987–2002.

Table A-9. Management regime for Canadian Chinook salmon fisheries affected by the PST.

Fishery	Management Regime (AABM or ISBM)
North Coast BC commercial troll	AABM
Queen Charlotte Islands sport	AABM
North and Central BC (including commercial net, marine sport along mainland coast, freshwater sport, Native fisheries in both marine and freshwater)	ISBM
West Coast Vancouver Island (WCVI) including commercial troll and outside sport)	AABM
Southern BC (including commercial net fisheries in Johnstone Strait, Strait of Juan de Fuca, Strait of Georgia, and the Fraser River, commercial troll fishery in Strait of Georgia, sport fisheries in the “inside” of WCVI, marine and freshwater sport fisheries, and both marine and freshwater Native fisheries)	ISBM

PFMC Fisheries

Chinook salmon are one of two primary target species in Pacific Coast salmon fisheries in PFMC-managed waters (i.e. Canadian border to Mexico, 3-200 nautical miles offshore). PFMC management focuses on five major stocks of Columbia River Basin fall Chinook: lower river hatchery tulle stock (LRH) and lower river wild bright stock (LRW), Spring Creek Hatchery tulle stock (SCH), all of which are part of the ESA-listed lower Columbia River ESU; upriver bright stock (URB), which includes the ESA-listed Snake River fall Chinook ESU; and mid-Columbia bright stock (MCB).

The PFMC STT annually publishes stock-specific preseason run forecasts that shape fishery management planning and harvest targets for the coming year. Forecasts are prepared by WDFW, ODFW, and the Columbia River Technical Advisory Committee (TAC) and presented annually in the PFMC *Preseason*

Report 1. Since 1964, age-specific estimates of escapement and in-river fishery catches have been used to establish age-specific linear regression relationships of cohort returns in previous run years. Therefore, the relationship of cohort returns from past years can be used as a predictor of the coming year return; for example, abundance of age 3 Chinook in 2002 can be applied to a linear relationship of age 3 and age 4 Chinook to estimate the age 4 Chinook return in 2003. Total run- or stock-specific forecasts are calculated by adding the estimated age-specific returns of all age classes represented in the run.

Ocean fisheries planning for the area North of Cape Falcon is coordinated between the PFMC and PSC. These fisheries are subject to the Chinook ISBM obligations contained within the 1999 Letter of Agreement. Management objectives for the Chinook fisheries in the North of Falcon area are to satisfy standards for ESA-listed stocks and, to the extent possible, provide for viable ocean and in-river fisheries while protecting depressed Columbia River natural stocks and the needs of hatcheries for fall Chinook brood stock. Lower Columbia River and Bonneville Pool hatchery fall Chinook historically have been the major stocks contributing to ocean fishery catches in the North of Cape Falcon area, and typically drive annual fishery quota levels. Federal ESA standards and the need to limit impacts on Puget Sound and lower Columbia Chinook stocks guide fishery management decisions; harvest is generally constrained by Chinook harvest quotas (Table A-10). Fisheries in the North of Cape Falcon area are divided into outside (ocean) and inside (Puget Sound and in-river) fisheries; treaty troll, non-treaty troll, and numerous recreational fisheries occur in this area.

Ocean Chinook harvest in PFMC-managed waters occurs throughout the year. California ocean commercial troll fisheries occur from April to October, although most of the landings occur May–July (Figure A-27). Oregon ocean commercial troll fisheries generally occur from May to November, although in recent years, fisheries have occurred in April and, in 2002, the fishery opened in March for the first time since 1976 (Figure A-27). The largest harvests historically occurred in July and August; 2002 harvest was greatest in June, September, and October. Washington ocean non-Indian troll fisheries occur from May to September; most of the harvest typically occurs in May and June (Figure A-27). Treaty Indian commercial ocean troll fisheries occur throughout the year; the majority of harvest occurs from May to August (Figure A-27). The ex-vessel value and the price per pound of troll-caught Chinook in California, Oregon, and Washington ocean fisheries has declined since the 1980s (Figure A-28). Ex-vessel values have increased slightly in recent years compared to the 1990s, potentially because of increased harvest as a result of higher ocean productivity and salmon abundance.

Table A-10. PFMC pre-season adopted Chinook catch quotas (in thousands of fish) for ocean fisheries north of Cape Falcon and critical stocks driving management, 1983–2001.

Year	Critical Stocks	Treaty Troll	Non-Indian Troll	Sport
1983	Columbia River hatchery and depressed upriver stocks	—	114.0	88.0
1984	LRH and SCH	8.3	16.7	10.3
1985	SCH	10.5	47.5*	37.2
1986	SCH	12.5	51.0	37.1
1987	SCH	15.8	58.2**	44.6
1988	Columbia River upriver stocks	60.0	73.7	29.8
1989	Columbia River upriver stocks	32.0	47.5	47.5
1990	LRH	31.2	37.5	37.5
1991	LRH	33.0	40.0	40.0
1992	Columbia River tules and Snake River falls	33.0	47.0	33.0
1993	Columbia River tules and Snake River falls	33.0	35.0	25.0
1994	LRH and Snake River falls	16.4	0	0
1995	LRH and Snake River falls	12.0	0	0
1996	LRH and Snake River falls	11.0	0	0
1997	Snake River falls	15.0	11.5	5.2
1998	LRH	15.0	6.5	3.5
1999	LRW (Lewis River)	30.0	28.5	21.5
2000	Columbia River tules and LRW (Lewis River)	25.5	12.5	12.5
2001 [§]	Columbia River tules	37.0	30.0	30.0

*Plus 7,430 hooking mortality for pink fishery.

** Plus 3,250 hooking mortality for pink fishery.

[§] Sharing of impacts on ESA-listed Puget Sound Chinook also affected the shaping of ocean and inside fisheries.




Figure A-27. California, Oregon, Washington, and treaty Indian ocean commercial troll landings (in thousands of fish) by month, 1976–2002.




Figure A-28. Total value and price per pound (in 2002 dollars) for ocean troll Chinook landings in California, Oregon, and Washington 1979–2002.

The recreational ocean harvest of Chinook in California is generally greater in the charter boat sector than the private sector, although in recent years, private boat landings have exceeded charter boat landings (Figure A-29). In Oregon, the recreational ocean harvest of Chinook is dominated by private boats although, compared to the 1990s, the charter boat catch has increased in recent years (Figure A-29). In Oregon, the ocean recreational harvest occurs from April to November; most landings occur in July and August. In Washington, charter boat landings historically exceeded private boat landings; after years of no harvest in the mid 1990s, catch of the two boat types have increased similarly in recent years (Figure A-29).




Figure A-29. California, Oregon, and Washington ocean recreational salmon effort (in thousands of angler trips) by boat type, 1979–2002.

Ocean fisheries management in PFMC-managed waters for the 2003 seasons was constrained by:

1. endangered Sacramento River winter Chinook south of Point Arena,
2. threatened California Coastal Chinook south of Cape Falcon,
3. Klamath River fall Chinook south of Cape Falcon,
4. threatened lower Columbia River natural tule Chinook north of Cape Falcon, and
5. management goals for naturally produced coho stocks over the entire PFMC management area, including threatened Oregon and California coastal stocks.

Management criteria for each West Coast stock were established for the 2003 season to achieve desired escapement objectives and manage the allowable ocean harvest (Table A-11 and Table A-12).

Table A-11. Management criteria and projected key stock escapements (in thousands of fish) for Chinook salmon in PFMC-adopted ocean salmon fisheries, 2003*.

Key Stock/Criteria	Projected Ocean Escapement** or Other Criteria	Spawner Objective or Other Standard
Columbia Upriver Brights	253.2	57.3; minimum ocean escapement to obtain 43.5 adults over McNary Dam, with normal distribution and no mainstem harvest
Mid-Columbia Brights	93.6	16.6; minimum ocean escapement to attain 5.75 adults for Bonneville Hatchery and 2.0 for Little White Salmon Hatchery egg-take, assuming average conversion and no mainstem harvest
Lower Columbia River Hatchery Tules	116.9	23.4; minimum ocean escapement 14.3 adults for hatchery egg-take, with average conversion and no lower river mainstem or tributary harvest
Lower Columbia River Natural Tules	47%	≤49%; ESA guidance met by a total adult equivalent fishery exploitation rate on Coweeman tules (NMFS ESA consultation standard)
Lower Columbia River Wild (threatened) [§]	23.4	5.7; MSY spawner goal for North Lewis River fall Chinook (NMFS ESA consultation standard)
Spring Creek Hatchery Tules	101.9	11.1; minimum ocean escapement to attain 7.0 adults for Spring Creek Hatchery egg-take, assuming average conversion and no mainstem harvest
Snake River Fall (threatened)	67%	≤70% of 1988–93 average age 3 and 4 AEQ exploitation rate for all ocean fisheries (NMFS ESA consultation standard)
Klamath River Fall	35.0	≥35.0 adult spawners to natural spawning areas
Age 4 ocean harvest rate	16%	<16%; NMFS ESA consultation standard for threatened California Coastal Chinook
Federally recognized tribal fishery	50%	50% share of adult harvest; equates to 41.4 adult fish for the Yurok and Hoopa tribal fisheries
KMZ recreational fishery	14.8%	Share of adult ocean harvest (none specified for 2003)
CA/OR commercial fishery	51%/49%	Share of adult commercial ocean harvest for the States of California/Oregon (none specified for 2003)
Klamath River recreational fishery	26.1%	>15% share of nontribal adult harvest specified by California Fish and Game Commission; equates to 10.8 adult fish
Sacramento River Winter (endangered)	Yes	Duration and timing of commercial and recreational seasons south of Point Arena not to differ substantially relative to those of 2000 and 2001 (NMFS ESA consultation standard)
Sacramento River Fall	517.0	122.0-180.0; Sacramento River fall natural and hatchery adult spawners

* Projections assume a SE Alaska TAC of 366.7 Chinook per PST agreement. For Canadian Chinook fisheries, assumed TACs were 112.5 for WCVI and a 1.4 in the Strait of Georgia troll fishery. All other Canadian troll and sport fisheries were assumed to have the same impact rates as in 2002

** Ocean escapement is the number of salmon escaping ocean fisheries and entering fresh water

[§] Includes minor contributions from EF Lewis and Sandy Rivers

Table A-12. Tentatively adopted 2003 fishery management measures for PFMC fisheries to mitigate potential impacts on ESA-listed ESUs of lower Columbia River salmonids.

ESU	Stock Representation in Salmon FMP	ESA Consultation Standard	2003 Council Guidance
Lower Columbia River Chinook—threatened	<ul style="list-style-type: none"> • Cowlitz, Kalama, Lewis spring • Lower River Hatchery fall • NF Lewis fall 	<ul style="list-style-type: none"> • No specific requirements • Brood year adult equivalent exploitation rate on Coweeman tule fall Chinook < 49% • 5,700 MSY level adult spawning escapement 	<ul style="list-style-type: none"> • Meet hatchery escapement goals • 47% total ocean and freshwater AEQ exploitation rate • 23,400 adults to Columbia River mouth
Upper Willamette Chinook— threatened	Upper Willamette River spring	No specific requirements; rare occurrence in PFMC fisheries	Troll fisheries N of Cape Falcon do not begin before 5/1
Upper Columbia River spring Chinook—endangered	Upper Columbia River spring	No specific requirements; rare occurrence in Council fisheries	Troll fisheries N of Cape Falcon do not begin before 5/1
Snake River fall Chinook— threatened	Snake River fall	≥30% reduction from the 1988-93 average adult (age 3 & 4) exploitation rate for all ocean fisheries	33% reduction from 1988-93 average (age 3 & 4) AEQ ocean exploitation rate
Snake River spring/summer Chinook—threatened	Snake River spring/summer	No specific requirements; rare occurrence in PFMC fisheries	Troll fisheries N of Cape Falcon do not begin before 5/1

Columbia River Fisheries

Columbia River fall fishing seasons are set by the Columbia River Compact, which is charged by Congressional and statutory authority to establish Columbia River Indian and non-Indian fishing seasons in joint waters bordering Washington and Oregon. The Compact considers annual abundance forecasts (produced by state biologists and endorsed by federal and tribal biologists) for each fall Chinook management stock in order to assure seasons set by the Compact are consistent with Ocean and In-River Management Agreements, treaty Indian and non-Indian allocation mandates, conservation measures of the ESA, as well as *U.S. v. Oregon* and state established escapement goals. The Compact considers agency, tribal, and public testimony in public hearings prior to taking regulatory action.

2002 Columbia River Salmon Management Guidelines

The CRFMP expired on July 31, 1999. A Management Agreement for upper Columbia River fall Chinook, steelhead, and coho has been reached by all parties for fall fisheries occurring in 2002. The following guidelines will be in place for the 2002 fall fishery management period.

- ***Allowable SRW fall Chinook impacts in combined non-Indian and treaty Indian mainstem fisheries below the confluence of the Snake River for 2002 result in a 30% reduction from base period harvest rates. The corresponding impact rate is 31.29% of the aggregate URB run.***
- ***The freshwater URB impact rate of 31.29% will be allocated 23.04% for treaty Indian fisheries and 8.25% for non-Indian fisheries.***
- ***Treaty Indian fall fisheries will be managed to limit impacts on wild Group B index steelhead to no greater than 15%. All non-Indian fisheries outside the Snake River basin will be managed for an upriver wild steelhead impact rate to not exceed 2% on wild Group B index steelhead.***
- ***Upriver fall Chinook escapement goals include 7,000 adult fall Chinook (4,000 females) to Spring Creek Hatchery and 43,500 adult fall Chinook (natural and hatchery included) for spawning escapement above McNary Dam.***
- ***Ocean and lower river fisheries will be managed to provide for Bonneville Dam escapement of at least 50% of the upriver coho salmon return.***
- ***Non-Indian fisheries will be managed for an impact rate of less than 5% for Columbia River***

Columbia River fall Chinook runs are divided for stock-specific management of Columbia River fisheries; the six major fall Chinook stock components are LRH, LRW, URB, BPH, mid Columbia River Brights (MCB; includes hatchery production of URB stock downstream of McNary Dam), and Select Area Brights (SAB; includes bright stock of Rogue River origin released from net pens in Youngs Bay, OR). Each stock varies in annual abundance and therefore the stock mix in fisheries is different in any given year (Table A-13). The *U.S. v. Oregon* TAC accounts for specific stock abundances to make a pre-season projection of harvest of each stock by fishery, time, and area. The pre-season forecasts are used to establish harvest agreements between Indian and non-Indian fisheries, sport and commercial fisheries, and ocean and Columbia River fisheries. State biologists monitor actual fish runs and fishery harvest by stock (Table A-14) to assure fisheries are adjusted in-season to meet management requirements. Several emergency Compact hearings are held during the course of each fall season to close or add fisheries in response to in-season updates.

Table A-13. Stock accountability of fall Chinook returning to Columbia River, 1980–2002.

Return Year	Total Return	URB	BPH	MCB*	LRH	LRW	SAB
1980	320,000	76,800	97,800	0	105,600	38,800	
1981	278,900	66,600	86,300	4,400	94,900	25,000	
1982	363,100	79,000	120,700	8,800	139,500	13,000	
1983	237,600	86,100	28,900	14,400	88,100	16,800	
1984	309,400	131,400	47,500	11,800	102,400	13,300	
1985	362,800	196,400	33,200	5,700	111,000	13,300	1,600
1986	494,800	281,600	16,600	17,400	154,800	24,500	2,000
1987	871,000	420,700	9,100	57,000	344,100	37,900	2,300
1988	784,700	339,900	12,000	78,000	309,900	41,700	3,200
1989	552,000	261,300	26,800	93,100	130,900	38,600	1,200
1990	312,900	153,600	18,900	59,000	60,000	20,300	1,100
1991	275,500	103,300	52,400	35,400	62,700	19,800	2,000
1992	219,000	81,000	29,500	31,100	62,600	12,500	2,300
1993	214,900	102,900	16,800	27,400	52,300	13,300	2,100
1994	254,000	132,800	18,500	33,700	53,600	12,200	3,200
1995	242,800	106,500	33,800	34,100	46,400	16,000	6,000
1996	330,800	143,200	33,100	59,700	75,500	14,600	4,700
1997	321,500	161,700	27,400	58,900	57,400	12,300	3,800
1998	255,400	142,300	20,200	36,800	45,300	7,300	3,500
1999	309,500	166,100	50,500	50,600	40,000	3,300	2,900
2000	253,300	155,700	20,500	36,900	27,000	10,200	4,900
2001	548,800	232,600	125,000	76,400	94,300	15,700	5,000
2002	733,100	276,900	160,800	108,400	156,400	24,900	5,700

- URB stock below The Dalles Dam

Table A-14. Stock composition of adult fall Chinook landed in mainstem Columbia River fisheries, 2001.

	Stock						Total
	LRH	LRW	BPH	URB	MCB	Other*	
Non-Indian Fisheries							
Recreational	4,845	356	3,159	11,146	7,483	1,590	28,579
Early August commercial	528	112	673	394	25	161	1,893
Late Aug/Sept commercial	2,654	985	2,815	6,596	4,186	258	17,494
October commercial	53	285	88	338	2,812	9	3,585
Select area commercial	1,193	0	117	823	0	2,040	4,203
Subtotal	9,273	1,738	6,852	19,297	14,506	4,088	55,754
Treaty Indian Fisheries							
Sales to licensed buyers	0	0	33,808	18,520	7,800	110	60,238
C&S and other non-ticketed catch	0	0	18,528	16,295	8,770	0	43,593
Subtotal	0	0	52,336	34,815	16,570	110	103,831
Total	9,273	1,738	59,188	54,112	31,076	4,198	159,585

* includes select area brights, spring Chinook, and non-Columbia Chinook

The lower river run (i.e. below Bonneville Dam) is composed of LRH, LRW, and MCB stocks, as well as minor stock components of LRB and SAB fall Chinook. MCB stocks are also produced in basins above Bonneville Dam, such as the Wind and Little White Salmon River basins. Columbia River salmon management guidelines for the 2002 fall fisheries were driven by the following restrictions on fall Chinook:

- Allowable Snake River wild (SRW) (part of URB group) fall Chinook impacts in combined non-Indian and treaty Indian mainstem fisheries below the confluence of the Snake River result in a 30% reduction from base period harvest rates (31.29% impact rate of the aggregate URB run),
- Freshwater URB impact of 31.29% will be allocated 23.04% for treaty Indian harvest and 8.25% for non-Indian fisheries,
- Above Bonneville Dam fall Chinook escapement goals include 7,000 BPH adults to Spring Creek Hatchery and 43,500 URB adults past McNary Dam,
- Lower river hatchery (LRH) escapement goal of 14,700 adult Chinook and Coweeman wild combined ocean and Columbia River exploitation rate of less than 49%, and
- Lewis River wild (LRW) Chinook escapement goal of 5,700 adults.

Early fall seasons target fall Chinook, particularly the non-Indian commercial openings in Zones 4 and 5, as well as the treaty Indian commercial harvest in Zone 6 (Figure A-30).




Figure A-30. Commercial and subsistence harvest of fall Chinook in the Columbia River from 1970–2001.

Since 2000, an agreement between non-Indian commercial and sport fishing interests has been established to address allocation of fall Chinook between non-Indian fisheries and general season structure. These agreements have been negotiated annually during the NOF pre-season planning process. Sharing of non-Indian SRW ESA impacts is addressed as well as equitable sharing of total

Chinook harvest between commercial, mainstem Columbia sport, and tributary sport fisheries. The 2002 non-Indian fall Chinook management agreement included the following elements:

Box 1. 2002 Non-Indian Columbia River Fall Fishery Chinook Allocation Agreement

- *Expected total catch of fall Chinook in the mainstem Columbia River downstream of the Snake River and in lower Columbia River tributaries is 85,400 of which 45,300 (53%) are expected to be harvested by the sport fishery and 40,100 (47%) by the commercial fishery.*
- *This agreement is limited by the non-Indian allocation of URB fall Chinook impacts of 8.25% as per the 2002 U. S. v. Oregon Fall Management Agreement. Non-Indian catch estimates are based on pre-season abundance forecasts referenced in Model Run "2002 MR-6" .*
- *URB fall Chinook impacts in fisheries downstream of the Snake River are allocated preseason at 4.36% to the sport fishery and 3.89% to the commercial fishery. The Columbia River Compact will use this URB impact allocation as guidance for making in-season management decisions concerning the Columbia River sport and commercial fisheries. Actual URB impacts in the fisheries may differ from pre-season estimates based on actual fishery catches, stock composition, and run-size updates. The U. S. v. Oregon TAC will update the URB run-size beginning in mid-September.*
- *The Buoy 10 sport fishery is modeled at 90% of the Chinook catch estimated for a full fishery to the end of the year (with a two fish daily limit) which is expected to deliver enough Chinook to continue the fishery through Labor Day. URB impacts with this fishery are projected to be 1.70%; or 39% of the total sport impacts of 4.36%.*
- *The mainstem sport fishery below McNary Dam is modeled at 95% of the Chinook catch estimated for a full fishery to the end of the year (with a two fish daily limit), which is expected to provide enough Chinook to continue the fishery through September, unless the mid-September URB run size and fishery updates indicate this fishery cannot continue past mid-September. URB impacts associated with this fishery are 2.66%; or 61% of the total sport impacts of 4.36%. For 2003 fall fishery discussions, the mainstem sport fishery will begin at 100%.*

Expectations for the commercial fishery include:

- *An early August salmon fishery up to four nights during the first week of August with potential for fishing during the early part of the second week of August in Zones 2 and 3 only. During the first week of August, the open area will include Zone 1 upstream to Longview Bridge and an 8-in minimum mesh restriction. Projected catch is 16,800 salmon. Chinook[URB impacts not used in this fishery will transfer to August Zone 4-5 fishery.*
- *Late August Zone 4-5 fishery during the last two week of August. Fishing is expected to occur 2-3 nights per each week with breaks in between fishing days. This fishery will not occur past August 29. Mesh size is 9-in minimum. Chinook/URB impacts not used in this fishery will transfer to September fisheries. Expected catch is 8,300 Chinook plus any transfers from the early August commercial fishery.*
- *Late August Zone 4-5 fishery during the last two week of August. Fishing is expected to occur 2-3 nights per each week with breaks in between fishing days. This fishery will not occur past August 29. Mesh size is 9-in minimum. Chinook/URB impacts not used in this fishery will transfer to September fisheries. Expected catch is 8,300 Chinook plus any transfers from the early August commercial fishery.*

- *Late fall fishery to begin the week of September 15. Fishery to occur in as much of Zones 1-5 as possible and will target coho or Chinook as determined by remaining impacts and in-season run strength. The late September Chinook harvest will be determined by the mid-September URB run size update and the actual URB impacts remaining that can be used by the commercial fishery.*
- *No sturgeon retention will be allowed in the August fisheries. Directed sturgeon fishing may occur during September or October to meet commercial allocation.*

Columbia River fall commercial fisheries are set by time, area, and gear type to correspond to timing differences between different fall Chinook stocks and other species to focus harvest on particular stocks and species at rates consistent with management intent. The commercial fishing areas are divided into zones with landings recorded by individual landing zone. Zones 1-5 are located downstream of Bonneville Dam. Zone 6 is located between Bonneville and McNary dams and is an exclusive tribal commercial fishing area. Non-Indian sport fisheries can occur in Zones 1-6 (Box 1).

Recent year commercial fisheries have been set in August primarily in Zones 4-5 to access URB, BPH, and MCB fish. The peak of the fall Chinook abundance in the lower river areas (Zones 1-2) occurs in late August and early September. Commercial fisheries have not been set in the lower area during this peak time to avoid over-harvest (Figure A-31 and Figure A-32). Commercial fisheries in the lower zones are typically set after mid-September to access the lower river and upriver Chinook stocks and coho after the peak of the Chinook runs has cleared the mainstem Columbia. Treaty Indian commercial fisheries are focused in September to harvest fall Chinook and summer steelhead (Figure A-31 and Figure A-32).




Figure A-31. Number of adult Chinook landed during early fall, late fall, and treaty Indian mainstem Columbia River seasons.




Figure A-32. Number of days fished during early fall, late fall, and treaty Indian mainstem Columbia River commercial seasons, 1982–2001.

Columbia River sport fisheries typically open in August. The Buoy 10 (estuary) fishery is managed under a total catch guideline to assure Chinook limits are not exceeded. In some years, there are emergency closures in late August and early September. The mainstem sport fishery upstream of the estuary area (upstream of Grays Point) is intended to remain open for the entire fall season, but on occasion it has closed early to avoid exceeding agreed Chinook harvest levels. Chinook harvest in both of these sport fisheries can be significant (Figure A-33).




Figure A-33. Buoy 10 fishery recreational catch and combined Oregon and Washington angler catch of Chinook on the lower Columbia River, 1982–2000.

Tributary fall Chinook sport fisheries for LRH tules occur principally in the Washougal, Cowlitz, Kalama, Grays, and Elochoman Rivers with most harvest occurring from late August through September. Annual harvest rates within each basin vary depending on abundance (Figure A-34 and Figure A-35). In large run years, harvest rates in individual tributaries can exceed 20%. In low run years, tributaries may be closed if needed to meet hatchery escapement needs. Fall Chinook fishing is closed in the Coweeman and EF Lewis Rivers and in Abernathy Creek to protect natural spawning Chinook.




Figure A-34. Fall Chinook tributary harvest rate in the NF Lewis, Grays, and Elochoman Rivers, 1988–99. Harvest rate equals sport catch divided by run size at tributary mouth.

North Lewis River fall Chinook sport fishing occurs from late August into October as Lewis River fish spawn later than tule stocks. Fishing is open when LRW fall Chinook projections indicate there are sufficient returns to harvest Chinook and meet the natural spawning escapement goal of 5,700 natural spawners. The fishery was open annually except for 1996–2000, when run forecasts indicated low returns. The LRW stock rebounded in 2001 and the sport fishery was reopened. The North Lewis River fall Chinook sport fishery is the only lower Columbia tributary fishery which targets healthy natural produced fall Chinook, as hatchery fall Chinook are not produced from the Lewis River.




Figure A-35. Fall Chinook tributary harvest rate in the Kalama, Cowlitz, Toutle, and Washougal Rivers, 1988–99. Harvest rate equals sport catch divided by run size at tributary mouth.

A.1.10. Assessment of Current Status and Limiting Factors

Listing Status

The Lower Columbia River Chinook salmon ESU was originally listed as a threatened species under the Endangered Species Act (ESA) on March 24, 1999 (Fed. Reg., V64, N56, p.14308). The BRT established by NMFS to examine the status of Chinook determined in 1998 that the estimated overall abundance of Chinook salmon in the Lower Columbia ESU is not cause for immediate concern. However, they found that apart from the relatively large, and apparently healthy fall-run population in the Lewis River, production in the ESU appears to be predominantly hatchery-driven with few identifiable native, naturally reproducing populations. Long- and short-term trends in abundance of individual populations are mostly negative, some severely so. About half of the populations comprising this ESU are very small, increasing the likelihood that risks due to genetic and demographic processes in small populations will be important. Numbers of naturally spawning spring-run Chinook salmon are very low. The BRT cautioned that it is possible that some native spring Chinook runs are now extinct, but that this loss is masked by the presence of naturally spawning hatchery fish. The BRT was particularly concerned about the inability to identify any healthy native spring run populations. The large numbers of hatchery fish in the ESU make it difficult to determine the proportion of naturally produced fish. While studies show that genetic and life history characteristics of populations in the Lower Columbia ESU still differ from those in other ESUs, the BRT identified the loss of fitness and diversity within the ESU as an important concern (NMFS 1998). NMFS assessed the status of the Lower Columbia River Chinook salmon ESU and reaffirmed the threatened listing on June 28, 2005 (Fed. Reg., V70, N123, p.37160). Subsequently, the current critical habitat was designated for this ESU on September 2, 2005 (Fed. Reg., V70, N123, p.52630).

Current Status

Lower Columbia Chinook salmon populations began declining by the early 1900s because of habitat alterations and unsustainable high harvest rates given the changing habitat conditions. Long- and short-term trends in abundance of individual populations are mostly negative, some severely so. About half of the populations comprising this ESU are very small, increasing genetic and demographic risks. Today, the once abundant natural runs of spring and fall Chinook have been largely replaced by hatchery production. Apart from the relatively large, and apparently healthy fall-run population in the Lewis River, production in the ESU appears to be predominantly hatchery-driven with few identifiable native, naturally reproducing populations.

Spring Chinook

Extinction risks were estimated to be very high in all seven Washington populations of spring Chinook, and eight of nine Washington and Oregon populations that comprise the ESU (Figure A-36). It is possible that some native spring Chinook runs are now extinct, but that this loss is masked by the presence of naturally spawning hatchery fish. No healthy spring Chinook populations exist in this ESU. Only one population is currently not at very high risk of extinction – Oregon’s Sandy River spring Chinook is at moderate risk. Current numbers of highly viable populations (0 in both the Cascade and Gorge strata) and average viability scores (Cascade = 0.71, Gorge = 0.25) fall well below the standards established by the WLC-TRT for viability (2 populations per strata at or above 3.0, strata average viability of at least 2.25).

High extinction risks of spring Chinook correspond to very low values of all population viability attributes (Table A-15). Abundance and productivity of naturally-spawning spring Chinook salmon in Washington are very low. Recent wild abundance is uncertain due to large numbers of naturally-

spawning hatchery spring Chinook but habitat assessments suggest that current natural production capacity is under 3,000 fish. Historical abundance was estimated to be approximately 50,000 to 100,000 fish based on historical records and habitat conditions. Total abundance of the combined natural and hatchery run has fluctuated widely over the last 35 years, reaching dangerous low levels of just 3,000 to 5,000 fish during 1996-1999 (Figure A-5).

Spatial structure of spring Chinook populations has been severely reduced by loss of access into core headwater production areas in the Cowlitz, Cispus, Lewis, and White Salmon rivers. The upper Cowlitz and Lewis systems historically accounted for the majority of the Washington lower Columbia spring Chinook production. While most historical habitats in the Kalama and Toutle remain accessible, these populations were historically much smaller and current distribution has been further reduced by habitat degradation in these systems.

Diversity of spring Chinook is estimated to have been greatly reduced due to pervasive hatchery effects and small population bottlenecks within the natural populations during recent low run size years. Remaining natural production areas for spring Chinook in Washington are very limited and hatchery-origin fish typically comprise a large fraction of the spawners in natural production areas.




Figure A-36. Status of historical spring Chinook populations among lower Columbia River subbasins.

Fall Chinook

Extinction risks were estimated to be very high in 12 of 15 Washington populations of fall Chinook (80%), and 16 of 22 Washington and Oregon populations (73%) that comprise the ESU (Figure A-37, Figure A-38). Only one population is currently at low or very low risk of extinction – Washington’s Lewis

River late fall (bright) Chinook. No tule fall Chinook population is currently at low or very low risk of extinction. The strongest tule fall Chinook populations in Washington currently include the lower Cowlitz (moderate risk) and Coweeman (high risk). Current numbers of highly viable tule fall Chinook populations (0 in each of Coast, Cascade, and Gorge strata) and average viability scores (Coast = 0.5, Cascade = 0.6, Gorge = 0.4) fall well below the standards established by the WLC-TRT for viability (2 populations per strata at or above 3.0, strata average viability of at least 2.25). The Cascade late fall Chinook stratum meets the strata average objective (2.5) but includes only one of the two required high or highly viable populations

Very high extinction risks of fall Chinook correspond to very low values abundance and productivity (Table A-15). Recent wild abundance of tule fall Chinook is uncertain due to large numbers of naturally-spawning hatchery fall Chinook but spawning ground surveys and habitat assessments suggest that current natural production capacity is under 5,000 fish. The majority of this occurs in the lower Cowlitz and Coweeman rivers. Historical abundance was estimated to be approximately 100,000 fish based on historical records and habitat conditions. Natural spawner numbers have varied widely over the last four decades but are typically very low. For instance, spawner numbers in the Coweeman and East Fork Lewis Rivers, where hatchery influence is very low, have ranged from a few hundred to a few thousand fish per year, but have regularly fallen to dangerous low levels of just 200 to 300 fish spawners per year (Figure A-6). In contrast, spawner numbers of the healthy Lewis River bright fall Chinook population have never fallen below 3,000 and spawning escapement has averaged approximately 10,000 fish per year (Figure A-7).

Spatial structure of most fall Chinook populations remains relatively high as fish continue to have access to most areas of historical spawning habitat. Exceptions include the upper Cowlitz and White Salmon rivers where dams block access. However, current distribution of most populations has been significantly reduced by habitat degradation in lower basin mainstem spawning areas utilized by fall Chinook.

Diversity of tule fall Chinook is estimated to have been greatly reduced due to pervasive hatchery effects and small population bottlenecks within the natural populations during recent low run size years. Hatchery-origin fish typically comprise a large fraction of the spawners in natural production areas, making it difficult to assess the true natural production capacity of the existing habitat. Large numbers of fall Chinook have produced in lower Columbia River hatcheries for decades and included long history of inter-population transfers and also significant influence of from out-of-basin stocks. In contrast, the single late fall Chinook population has been largely uninfluenced by hatchery production and has not seen the small population bottlenecks observed in other fall Chinook populations.




Figure A-37 Status of historic bright late fall Chinook salmon populations among lower Columbia River subbasins. Extinction risks are based on viability scores.




Figure A-38. Status of historical tule fall Chinook salmon populations among lower Columbia River subbasins. Extinction risks are based on viability scores

Table A-15. Baseline status and recovery objectives for lower Columbia River Chinook populations (relative to listing period baseline).

Population	Contribution	Baseline viability				Obj.	Prod. target	Abundance		
		A&P	S	D	Net			Historical	Baseline	Target
Coast Fall										
Grays/Chinook	Contributing ²	VL	H	VL	VL ²	M+	+500%	800	<50	1,000
Eloch/Skam ^C	Primary	VL	H	L	VL ²	H	+150%	3,000	<50	1,500
Mill/Aber/Germ	Primary ¹	VL	H	L	VL ²	H	+155%	2,500	50	900
Youngs Bay (OR)	Stabilizing	-- ³	-- ³	-- ³	L	L	-- ³	-- ³	-- ³	-- ³
Big Creek (OR) ^C	Contributing ¹	-- ³	-- ³	-- ³	VL	L	-- ³	-- ³	-- ³	-- ³
Clatskanie (OR)	Primary	-- ³	-- ³	-- ³	VL	H	-- ³	-- ³	-- ³	-- ³
Scappoose (OR)	Primary ¹	-- ³	-- ³	-- ³	L	H	-- ³	-- ³	-- ³	-- ³
Cascade Fall										
Lower Cowlitz ^C	Contributing	VL	H	M	VL ²	M+	+50%	24,000	500	3,000
Upper Cowlitz	Stabilizing	VL	VL	M	VL	VL	--	28,000	0	--
Toutle ^C	Primary ¹	VL	H	M	VL ²	H+	+265%	11,000	<50	4,000
Coweeman ^G	Primary	VL	H	H	VL ²	H+	+80%	3,500	100	900
Kalama	Contributing ²	VL	H	M	VL ²	M	+110%	2,700	<50	500
Lewis ^G	Primary	VL	H	H	VL ²	H+	+280%	2,600	<50	1,500
Salmon	Stabilizing	VL	H	M	VL	VL	--	n/a	<50	--
Washougal	Primary	VL	H	M	VL ²	H+	+190%	2,600	<50	1,200
Clackamas (OR) ^C	Contributing	-- ³	-- ³	-- ³	VL	M	-- ³	-- ³	-- ³	-- ³
Sandy (OR)	Contributing ¹	-- ³	-- ³	-- ³	VL	M	-- ³	-- ³	-- ³	-- ³
Cascade L Fall										
Lewis NF ^{C,G}	Primary	VH	H	H	VH ¹	VH	0%	23,000	7,300	7,300
Sandy (OR) ^{C,G}	Primary	-- ³	-- ³	-- ³	L	VH	-- ³	-- ³	-- ³	-- ³
Cascade Spring										
Upper Cowlitz ^{C,G}	Primary	VL	L	M	VL ²	H+	>500%	22,000	300	1,800
Cispus ^{C,G}	Primary	VL	L	M	VL ²	H+	>500%	7,800	150	1,800
Tilton	Stabilizing	VL	VL	VL	VL	VL	0%	5,400	100	100
Toutle	Contributing	VL	H	L	VL	M	>500%	3,100	100	1,100
Kalama	Contributing ²	VL	H	L	VL	L	>500%	4,900	100	300
Lewis NF ^C	Primary	VL	L	M	VL	H	>500%	15,700	300	1,500
Sandy (OR) ^{C,G}	Primary	-- ³	-- ³	-- ³	M	H	-- ³	-- ³	-- ³	-- ³
Gorge Fall										
L. Gorge (WA/OR)	Contributing	VL	M	L	VL ²	M	>500%	n/a	<50	1,200
U. Gorge (WA/OR) ^C	Contributing ¹	VL	M	L	VL ²	M	>500%	n/a	<50	1,200
White Salmon ^C	Contributing	VL	L	L	VL	M	>500%	n/a	<50	500
Hood (OR)	Contributing ⁴	-- ³	-- ³	-- ³	VL	L ⁴	-- ³	-- ³	-- ³	-- ³
Gorge Spring										
White Salmon ^C	Contributing	VL	VL	VL	VL	L+	>500%	n/a	<50	500
Hood (OR)	Primary	-- ³	-- ³	-- ³	VL	VH	-- ³	-- ³	-- ³	-- ³

¹ Increase relative to the interim plan.

² Reduction relative to the interim plan.

³ Addressed in the Oregon Management Unit plan.

⁴ OR analysis indicates a low probability of meeting their delisting objective of high viability for this population.

^C Designated as a historical core population by the Technical Recovery Team.

^G Designated as a historical legacy population by the Technical Recovery Team.

Current Limiting Factors

Net Effect of Manageable Factors

Declines in status of lower Columbia River Chinook salmon result from the combined impacts of human activities involving freshwater habitat, estuary habitat, dam construction and operation, fishing, fish hatcheries, and ecological factors such as predation. Impacts of each factor are compounded across the salmon life cycle to drive most populations to current very low levels. Net effects of quantifiable and potentially manageable impacts translate into an estimated 75-100% (average 96%) reduction in abundance and productivity of lower Columbia Chinook populations in Washington (Figure A-39). Thus, current fish numbers represent only 0-25% of the historical production potential in the absence of potentially manageable impacts. Total reductions would be even greater if all human impacts could be effectively quantified. No single factor accounts for the majority of the reduction in fish numbers and the significance of specific factors varies from population to population (Table A-16).




Figure A-39. Net effect and relative contribution of potentially manageable impact factors on Chinook salmon in Washington lower Columbia River subbasins. Net effect is the approximate reduction from historical fish numbers as a result of manageable factors included in this analysis.

Table A-16 Productivity improvements and impact reduction benchmarks consistent with recovery of Washington lower Columbia River Chinook populations if reductions are distributed across factors in proportion to their baseline impact.

Population	Productivity		Baseline impacts (listing date reference)						Impacts at target					
	Target	Trib	Est	Dams	Harv	Hat	Pred	Δ	Trib	Est	Dams	Harv	Hat	Pred
<u>Coast Fall</u>														
Grays/Chinook	+500%	0.40	0.23	0.00	0.65	0.50	0.09	61%	0.16	0.09	0.00	0.26	0.20	0.03
Eloch/Skam	+150%	0.30	0.23	0.00	0.65	0.50	0.09	29%	0.21	0.17	0.00	0.46	0.35	0.06
Mill/Aber/Germ	+155%	0.40	0.23	0.00	0.65	0.49	0.10	28%	0.29	0.17	0.00	0.47	0.35	0.07
<u>Cascade Fall</u>														
Lower Cowlitz	+50%	0.70	0.23	0.00	0.65	0.50	0.10	8%	0.64	0.21	0.00	0.60	0.46	0.09
Upper Cowlitz	0%	0.80	0.23	1.00	0.65	0.50	0.10	0%	0.80	0.23	1.00	0.65	0.50	0.10
Toutle	+265%	0.60	0.23	0.05	0.65	0.50	0.10	32%	0.41	0.16	0.03	0.44	0.34	0.07
Coweeman	+80%	0.50	0.23	0.00	0.65	0.23	0.10	18%	0.41	0.19	0.00	0.53	0.19	0.08
Kalama	+110%	0.40	0.23	0.00	0.65	0.50	0.10	21%	0.31	0.18	0.00	0.51	0.39	0.08
Lewis EF	+280%	0.40	0.23	0.00	0.65	0.50	0.11	42%	0.23	0.13	0.00	0.38	0.29	0.06
Salmon	0%	0.90	0.23	0.00	0.65	0.50	0.11	0%	0.90	0.23	0.00	0.65	0.50	0.11
Washougal	+190%	0.30	0.23	0.00	0.65	0.50	0.11	34%	0.20	0.15	0.00	0.43	0.33	0.07
<u>Cascade L Fall</u>														
Lewis NF	0%	0.10	0.23	0.16	0.50	0.05	0.11	0%	0.10	0.23	0.16	0.50	0.05	0.11
<u>Cascade Spring</u>														
Upper Cowlitz	>500%	0.90	0.15	0.90	0.50	0.50	0.22	50%	0.45	0.08	0.45	0.25	0.25	0.11
Cispus	>500%	0.90	0.15	1.00	0.50	0.50	0.22	50%	0.45	0.08	0.50	0.25	0.25	0.11
Tilton	0%	0.80	0.15	1.00	0.50	0.50	0.22	0%	0.80	0.15	1.00	0.50	0.50	0.22
Toutle	>500%	0.90	0.15	0.00	0.50	0.50	0.22	50%	0.45	0.08	0.00	0.25	0.25	0.11
Kalama	>500%	0.90	0.15	0.00	0.50	0.50	0.22	50%	0.45	0.08	0.00	0.25	0.25	0.11
Lewis NF	>500%	0.40	0.15	0.95	0.50	0.50	0.22	50%	0.20	0.08	0.48	0.25	0.25	0.11
<u>Gorge Fall</u>														
L. Gorge (WA/OR)	>500%	0.70	0.23	0.30	0.65	0.50	0.11	50%	0.35	0.11	0.15	0.33	0.25	0.06
U. Gorge (WA/OR)	>500%	0.70	0.22	0.58	0.65	0.50	0.14	50%	0.35	0.11	0.29	0.33	0.25	0.07
White Salmon	>500%	0.70	0.22	0.58	0.65	0.50	0.14	50%	0.35	0.11	0.29	0.33	0.25	0.07
<u>Gorge Spring</u>														
White Salmon	>500%	0.70	0.14	0.96	0.50	0.50	0.27	50%	0.35	0.07	0.48	0.25	0.25	0.13

Fisheries

Fishing impacts on lower Columbia wild spring Chinook averaged 53% at listing and has since been reduced to 22%. Current mortality is incidental to target fisheries for fin-clipped Willamette, lower Columbia, and upper Columbia hatchery fish. Additional harvest of wild spring Chinook occurs in the ocean incidental to target fisheries for Alaskan, Canadian, Columbia River Hatchery, and California Hatchery Chinook stocks (Figure A-40). The exploitation rate of spring Chinook has fluctuated over time, ranging from 20 to 65%. The current exploitation of hatchery spring Chinook is similar to historical exploitation rates, while wild spring Chinook exploitation is considerably lower than historical rates.




Figure A-40. Approximate spring Chinook fishery exploitation rates over time and allocation of current exploitation rates among fisheries

Fishing impact limits on lower Columbia fall Chinook averaged 65% for tules and 50% for brights at listing. Recent rates have been less than 49% for tules and less than 40% on brights. Columbia basin fisheries targeting upriver bright and Columbia hatchery Chinook account for a third to half of total fishing rates, with the remainder primarily distributed between Oregon, Washington, Canada, and Alaska ocean fisheries targeted on Alaska, Canadian, Columbia River hatchery, and California hatchery Chinook (Figure A-41). Current fishing rates on fall Chinook are approximately half their historical average. For instance, Chinook fishing rates remained fairly constant at 70-80% through the 1980s and 1990s, but have declined to approximately 40%.




Figure A-41. Approximate fall Chinook fishery exploitation rates over time and allocation of current exploitation rates among fisheries. Time series is for tule fall Chinook.

Hatcheries

Hatchery influence is significant for most Washington lower Columbia Chinook populations (Table A-17). Spring Chinook populations are primarily naturally spawning hatchery fish. Hatchery releases of spring Chinook smolts range from 0 to 1.267 million. Hatchery fractions of adults generally average 90% with reintroduction attempts in the upper Cowlitz basin relying entirely on hatchery stock. Current spring Chinook hatchery broodstock are primarily derived from local populations and moderately affected by hatchery practices (category 2) or derived from other populations within the same ESU (category 3). The indexed potential for negative impacts of hatchery spawners on wild population fitness was estimated to range from 27 to 45%. However, the high incidence of hatchery spawners suggests that the fitness of natural and hatchery fish is now probably quite similar and natural populations could collapse without continued hatchery subsidy under current habitat conditions. Inter-specific hatchery predation impacts on juvenile fall Chinook range from 0% in basins without significant releases of coho, steelhead or spring Chinook to 15 and 27% in the Lewis and Cowlitz basins where large hatchery programs are underway.

For Tule fall Chinook, potential fitness impacts of hatchery fish range from 0 to 34% (Table A-17). Hatchery fish do not contribute to Coweeman and Lewis tule Chinook populations, hence, their genetic legacy designation by the Technical Recovery Team. Hatchery fractions generally exceed 50% for other Cascade and Coast tule populations but are less among Gorge populations. Current hatchery releases of tule Chinook smolts range from 0 to 5 million per year per subbasin although historical releases were greater. Hatchery broodstocks are primarily derived from local populations and moderately affected by hatchery practices (category 2) for Cascade populations. Many Chinook salmon hatchery programs were developed initially from out-of-subbasin or multiple stocks but this practice has been largely discontinued. In most cases, brood stock mixing was limited to a few stocks and performed only during the initial years of establishing the hatchery program. After the hatchery program had been established, brood stock collection came from adults returning to the hatchery facility within the basin, aside from minimal outside brood stock usage during years of hatchery shortfalls. Most hatchery programs now use adults returning to the hatchery facility for brood stock; future genetic risks for outside brood stock usage appear minimal. In contrast, Coast and Gorge tule Chinook hatchery programs primarily originated from other populations within the same ESU (category 3) because natural populations in those areas were typically small.

Bright fall Chinook in the Lewis basin have included a small fraction of hatchery fish until recently when the program was discontinued. The potential for fitness impacts was estimated to be less than 1% because of the low incidence of hatchery fish and the native, local brood source (category 1).

Supplementation has not been the goal of most Chinook hatchery programs; these are intended to mitigate for losses of Chinook by providing fish for harvest opportunities. The exceptions are a fall Chinook hatchery program at the Sea Resources Hatchery on the Chinook River, spring Chinook, coho, and winter steelhead programs at the Cowlitz hatcheries for the upper Cowlitz River, and chum programs at the Grays River hatchery for the Grays River and at the Washougal Hatchery for lower Gorge chum. A spring Chinook, coho, and winter steelhead supplementation program is expected to be developed at the Lewis River hatcheries for the upper Lewis.

The Columbia River Hatchery Scientific Review Group (HSRG) recently completed a review of hatchery programs affecting salmon and steelhead populations in the Lower Columbia River. The objective of the review was to change the focus of the Columbia River hatchery system from a harvest supplementation/hydropower mitigation focus to an ecosystem-based approach founded on the idea that harvest goals are sustainable only if they are compatible with conservation goals. Individual population reviews and recommendations can be found at http://www.hatcheryreform.us/mfs/reports/appendix/welcome_show.action.

Table A-17. Presumed reductions in wild population fitness as a result of natural hatchery spawners and survival as a result of interactions with other hatchery species for Washington lower Columbia River Chinook populations.

Population	Annual releases ^a	Hatchery fraction	Fitness category	Assumed fitness	Fitness impact	Interacting releases ^j	Interspecies impact
Coast Fall							
Chinook/ Grays	107,500 ^b	0.37	3	0.5	0.18	190,000	0.01
Eloch/Skam	2,000,000	0.69	3	0.5	0.34	1,050,000	0.05
Mill/Aber/Germ	0 ^c	0.47	3	0.5	0.24	0	0
Cascade Fall							
Lower Cowlitz	5,000,000	0.67	2	0.7	0.20	5,319,500	0.27
Upper Cowlitz	0	0.67	2	0.7	0.20	--	--
Toutle	2,500,000	0.90	2	0.7	0.27	850,000	0.04
Coweeman	0 ^d	0.00	0	--	0.00	20,000	0
Kalama	5,000,000	0.67	2	0.7	0.20	1,380,000	0.07
Lewis/Salmon	0	0.00	0	--	0.00	115,000	0.01
Washougal	4,000,000	0.57	2	0.7	0.17	620,000	0.03
Gorge Fall							
L Gorge	0 ^e	na	2	0.7	na	0	0
U. Gorge (Wind)	0 ^f	0.17	4	0.3	0.12	1,420,000	0.07
White Salmon	0	0.22	3	0.5	0.11	0	0
Cascade L Fall							
Lewis NF	0 ^g	0.13	1	0.9	0.01	3,070,000	0.15
Cascade Spring							
Cowlitz	1,267,000 ^h	0.90	2	0.7	0.27	--	--
Cispus	-- ^h	0.90	2	0.7	0.27	--	--
Tilton	0	0.90	2	0.7	0.27	--	--
Toutle	0	0.90	3	0.5	0.45	--	--
Kalama	500,000	0.90	3	0.5	0.45	--	--
Lewis NF	1,050,000 ⁱ	0.90	3	0.5	0.45	--	--
Gorge Spring							
White Salmon	0 ^j	1.00	4	0.3	0.70	--	--

^a Annual release goals.

^b Number refers to fall Chinook hatchery program underway to restore a naturally producing population in the Chinook River. The Grays River fall Chinook hatchery program stopped releasing smolts in 1998; hatchery returns were expected to significantly diminish starting with the 2002 return.

^c Abernathy hatchery stopped releasing fall Chinook in 1995; hatchery returns were expected to significantly diminish starting with the 1999 return.

^d Hatchery fall Chinook have not been released in the Coweeman River basin since the early 1980s and tagged hatchery strays have not been recovered during spawning surveys since that time.

^e There are no hatchery fall Chinook programs in the lower gorge tributaries; fall Chinook from the Spring Creek NFH were released in Hamilton in 1977.

^f There are no hatchery fall Chinook programs in the Wind River basin. Fall Chinook were historically produced at the Carson NFH and released in the basin, however, production shifted to spring Chinook in 1981.

^g The Lewis River fall Chinook hatchery program was discontinued in 1986. There is no hatchery fall Chinook program in Salmon Creek.

^h 300,000 fingerling spring Chinook from Cowlitz Trout Hatchery are released annually in an attempt to restore the upper Cowlitz population. An additional 967,000 yearlings are released in the lower Cowlitz from Cowlitz Salmon Hatchery.

ⁱ Current releases are in the lower Lewis. Reintroduction into the upper Lewis is also under consideration in the hydroelectric re-licensing process.

^j No hatchery spring Chinook are released into the White Salmon. However, 1,000,000 and 1,420,000 hatchery spring Chinook are released in the Little White Salmon and Wind rivers, respectively.

^j Includes steelhead, coho and spring Chinook for fall Chinook. Not applicable for spring Chinook.

Tributary Habitat

EDT analyses suggest that stream degradation has substantially reduced the habitat potential for Chinook salmon in all Washington lower Columbia River subbasins where analyses have been completed (Figure A-42). Declines in habitat quality and quantity for Chinook salmon have reduced current productivity potential to 0-76% and current equilibrium fish numbers to 0-50% of the historical template. Substantial stream habitat improvements would be necessary to reach optimum conditions (i.e. PFC) for Chinook salmon in most subbasins. Restoration of optimum habitat quality would be expected to increase habitat capacity by 100 to 24,000 adult Chinook per subbasin.

Fall Chinook salmon rely on the lower and middle mainstem reaches of large streams and rivers. Channel instability, low habitat diversity, and sedimentation consistently limit habitat suitability for fall Chinook in these areas. Spring Chinook use the upper basins of large river systems. Many of these upstream areas continue to provide suitable habitat for spring Chinook but dams limit access in the Lewis and Cowlitz basins. More detailed descriptions of stream habitat conditions and effects on fish in each subbasin may be found in Volume II of the Technical Foundation.




Figure A-42. Current, optimal, and historical subbasin productivity and equilibrium numbers inferred for Chinook salmon from stream reach habitat conditions using EDT.

Dams

Dam impacts on Washington lower Columbia Chinook were estimated to range from 0 to 100% (Figure A-43). Dams on the Cowlitz have inundated or blocked access to over 90% of the spring Chinook and 45% of the fall Chinook habitat based on EDT assessments. On the North Fork Lewis River, 90% of the spring Chinook and 7% of the fall Chinook habitat has been inundated or blocked. Passage mortality at Bonneville Dam was assumed to average 10% for juveniles and an additional 10% for adults based on a synthesis of the available literature. Bonneville Dam inundated approximately half of the fall Chinook habitat on the Wind and White Salmon rivers. Assessments also included an assumed 20% reduction in fall Chinook productivity in the Columbia River mainstem as a result of Bonneville Dam operations. Dam operations in the Cowlitz and Lewis River have similar potential to affect downstream habitat conditions for Chinook but the significance of this impact is unknown.




Figure A-43. Assumed dam impacts on Washington lower Columbia Chinook populations.

Mainstem and Estuary Habitat

Mainstem and estuary habitat impacts were estimated to account for approximately a 30-40% reduction in productivity of fall Chinook which migrate as subyearlings and a 20% reduction for the primarily spring-migrating spring Chinook.

Predation

Predation mortality rates of juvenile fall and spring Chinook by pikeminnow and terns at the time of ESA listing was assumed to average 20% to 25% depending on travel distance from the subbasin to the ocean. Pikeminnow and tern management is projected to reduce salmonid predation by approximately 50%. Tern predation is almost entirely an artifact of recently established colonies on dredge spoil islands in the estuary but the current rate (9%) is less than half that observed prior to downstream translocation of the Rice Island colony (20%). Pikeminnow predation was greatest for populations that originate in Bonneville Reservoir tributaries, and must pass the pikeminnow gauntlet in Bonneville Dam forebay and tailrace, and then travel the entire 145 miles from Bonneville to the Estuary. Predation rates by seals and sea lions on adult Chinook were assumed to be four times greater on spring migrants (12%) than on fall migrants (3%).

Summary Assessment

1. Human activities, including fishing, hatchery operation, alteration of stream, river, and estuary habitats, hydropower development and operation, and potentially manageable predation have

collectively reduced productivity of spring and fall Chinook populations to 0-15% of historical levels. Recovery efforts will require significant improvements in multiple areas because no single factor accounts for the majority of the reduction in fish numbers.

2. Current fishing impacts on spring Chinook are modest and provide limited opportunities for increasing their numbers through additional regulation of fisheries. Higher fishing impacts on fall Chinook salmon provide some opportunity for increasing numbers through additional fishery constraints. Impacts have been reduced since listing. Additional reductions would likely require changes in ocean and freshwater fisheries. Fall Chinook impacts are widely distributed among U.S. ocean, Canada ocean, Alaska ocean, and Columbia River fisheries. Since Lower Columbia fall Chinook comprise only a small portion of the catch in many fisheries, additional constraints for their protection will forgo harvest of larger numbers from healthy wild and especially hatchery populations. Intensive fishery management processes provide significant opportunities for limiting fishing risks by tailoring annual harvests to fish availability.
3. Reduced productivity of wild populations as a result of interbreeding with potentially less-fit hatchery fish is among the most significant of hatchery concerns for wild stock recovery although these negative effects are at least partially offset by the demographic benefits of additional spawners. Potential negative impacts increase with the proportion of hatchery spawners and the genetic and phenotypic disparity between wild and hatchery fish. Potential fitness impacts among Washington lower Columbia fall Chinook populations range from 0 to 34%. Potential impacts are substantially greater among spring Chinook populations (27-70%) where hatchery fish comprise the majority of many remaining runs. Inter-specific hatchery predation impacts on juvenile fall Chinook range from 0% in basins without significant releases of coho, steelhead or spring Chinook to 15 and 27% in the Lewis and Cowlitz basins where large hatchery programs are underway.
4. Stream habitat conditions significantly limit Chinook salmon in all Washington lower Columbia River subbasins where EDT analyses have been completed. Substantial stream habitat improvements would be necessary to reach optimum conditions (i.e. PFC) in all subbasins. The significance of stream habitat suggests that recovery may not be feasible without substantial improvements in habitat quantity and quality.
5. Estuary and mainstem habitats are critical to Chinook salmon life history with assumed habitat impacts greater for fall Chinook (30-40% reduction) than spring Chinook (20% reduction).
6. Hydropower development in tributaries is currently the most significant factor limiting spring Chinook populations and effective recovery may not be feasible without effective passage measures at Cowlitz and Lewis dams. Hydro impacts on Gorge fall Chinook populations are also significant but can be only partially addressed by passage improvements. For instance, inundation of limiting habitat for fall Chinook in Lower Gorge Tributaries may constrain restoration of large and productive natural spawning in those areas.



A.2. Coho Salmon

Coho salmon (*Oncorhynchus kisutch*) is a widespread species of Pacific salmon, with production in most major river basins around the Pacific Rim from central California to Korea and northern Hokkaido, Japan (Laufle et al. 1986), as well as in many smaller independent tributaries through the region. In the lower Columbia River basin, coho salmon historically returned to spawn in all accessible tributary reaches. Until the early 1800s, these watersheds remained essentially untouched by human development. Heavy growth of coniferous trees and understory vegetation armored many river stretches. The Columbia River systems' cool streamflows, clean gravel beds, and deep pools supported healthy populations of coho, other salmon, and steelhead. These pristine environments began to change in the mid-1800s, often causing declines in salmonid production. Coho runs were further affected by hydro development and harvest pressure on the lower Columbia River. Harvest emphasis moved to coho as Chinook abundance dropped; peak commercial catches of coho in the Columbia River occurred in 1925 (Lichatowich and Mobrand 1995).

Present coho populations in tributaries to the Washington side of the lower Columbia River have been heavily influenced by extensive hatchery releases. Investigations report that a number of local populations of coho salmon in the area have become extinct, and that the abundance of many others is depressed (Brown and Moyle 1991, Nehlsen et al. 1991, Frissell 1993, NMFS 1995).

A.2.1. Life History and Requirements

The freshwater life history cycle of lower Columbia coho salmon populations follows the timing of seasonal changes in river flow and water temperatures in lower Columbia River tributaries. The region generally has a mild climate with warm, relatively dry summers and cool, wet winters. The river environments coho enter are characterized by relatively low elevations in the headwaters (1,640-3,280 ft [500-1,000 m]), with moderate amounts of precipitation (80-95 in/year [200-240 cm/year]). These rivers display relatively low flows during late summer and early fall, increased river flows and decreased water temperatures beginning in early October, and a single flow peak in December or January.




Figure A-44. Life cycle for lower Columbia River coho salmon.

Upstream Migration Timing

Coho runs to the Columbia River show considerable temporal variability in river entry and spawn timing. Coho salmon begin to return to the Columbia River in August, continuing through December/ January and peaking in September/October (Figure A-44). This variability resembles the pattern of river entry in other river systems, such as the Chehalis in southwest Washington, the Skagit in northern Washington, and the Klamath in northern California (Leidy and Leidy 1984, WDF et al. 1993).

Coho generally return in two runs:

- Early-returning (Type S) coho enter the Columbia River in mid-August and begin entering tributaries in early September, with spawning peaks from mid-October to early November.³
- Late-returning (Type N) coho pass through the lower Columbia from late September through December and enter tributaries from October through January.⁴ Most spawning occurs from November to January, but some spawning ranges to February and as late as March.

In some regions, individual coho stocks show exceptionally early or late run timings; these stocks are often referred to as summer or winter runs, respectively (Godfrey 1965), and are thought to have evolved in response to particular flow conditions (Sandercock 1991). The relationship between populations with unusually timed runs and normally timed runs within the same basin is not well understood. For example, in some cases, such as the Soleduck (Washington coast) and Clackamas (Willamette River) rivers, differently-timed, sympatric runs are thought to be largely reproductively isolated from each other (Houston 1983, Cramer and Cramer 1994), while in the Grays Harbor basin, there is believed to be reproductive overlap (WDF et al. 1993). Unusually timed runs are found in many

³ referred to as Type S because their ocean migration is generally south of the Columbia River.

⁴ referred to as Type N because of a more northern ocean distribution.

geographic areas. However, because there is no evidence to suggest that all runs of a certain type are closely related, differently timed runs are considered to be a component of overall life history diversity within each area (NMFS 1995).

Total residence time of coho in freshwater streams is highly variable and dependent upon both environmental and population specific factors, and can range from a few days to months. Residence time in the spawning areas of the stream (i.e. survey life, or the number of days the average spawner is alive in a survey area) has ranged from 3-15 days across the Pacific coast region for 30 reported population observations, with an average of 11.4 days (Perrin and Irvine 1990).

Spawning

In general, earlier migrating fish spawn farther upstream within a basin than later migrating fish, which enter rivers in a more advanced state of sexual maturity (Sandercock 1991). Spawning usually occurs between November and early February, but generally peaks October–December and can extend into March (WDF et al. 1993). Tributary spawning extends from October through at least February, and into March in some river systems.

The timing of coho spawning can also reflect water temperature changes in a particular river system. Lister et al. (1981) found that spawn timing of coho salmon in tributaries of the Cowichan River (British Columbia) was strongly correlated to tributary water temperature. Coho salmon spawning in warmer tributaries spawned later than those spawning in colder tributaries. Such factors make determining and comparing when coho will enter a river or spawn difficult because of the temperature variability within basins (NMFS 1995).

Other environmental factors influence coho spawning as well. Adult coho returning to spawn need adequate flows and water quality, and unimpeded passage to their natal grounds. Thus, the onset of coho salmon spawning in lower Columbia tributaries is tied to the first significant fall freshet. They often mill near the river mouths or in lower river pools until freshets occur. They also need deep pools with vegetative cover and instream structures such as root wads for resting and shelter from predators. In addition, successful spawning and incubation depend on the presence of appropriate sized gravel. In research on the upper Toutle and Green rivers within the Cowlitz basin, Burner (1951) described coho redd characteristics. Coho prefer substrate of 6 in (15 cm) or smaller; only 10% of the coho redds were constructed in gravels greater than 6 in (15 cm) in diameter (Burner 1951).

While spawner size in naturally spawning populations normally shows considerable spatial and temporal variability, scientists have found that coho salmon, throughout their range, are, over time, becoming smaller, and the rates of decrease are specific to populations of fish or to certain areas (Ricker 1981, Bigler and Helle 1994). While the size of coho salmon adults is declining fastest in the Puget Sound/Strait of Georgia area, smaller fish are also returning to Columbia River tributaries. Results of a regression analysis of coho salmon size (FL [cm] or total weight oz [kg]) over time, found that from 1952–92, the change in estimated weight for fish in the Columbia River fishery was statistically significant ($P=0.00$; NMFS 1995). Fecundity in coho increases with length (Salo and Bayliff 1958, Shaplov and Taft 1954).

It is not clear whether such size reductions are because of harvest practices, effects of fish culture, declining ocean productivity, density-dependence effects in the marine and freshwater environments attributable to large numbers of hatchery releases, or a combination of these factors. It is also not known whether there have been permanent genetic changes related to changes in fish size. Regardless of its cause or potential genetic effects, reduced adult size in itself poses a number of serious risks to natural populations of coho, and could be a sign of other factors placing the population at risk (NMFS 1995).

Incubation and Emergence

Coho eggs normally incubate for 1–6 months before hatching. During incubation, the eggs need stable gravel that is not choked with sediment, thus river channel stability is vital at this life history stage. Floods have their greatest impact to salmon populations during incubation, and flood impacts are worsened by human activities—particularly those related to timber harvest and urban development. In a natural river system, the upland areas are forested, and the trees and their roots store precipitation and slowly release stormwater into the stream. Trees within the riparian area also contribute large pieces of wood to the river that create habitat diversity and slow streamflow. Natural systems also have floodplains that connect directly to the river, and provide habitat and temporarily store floodwater. In a healthy river, erosion or sediment input is great enough to provide new gravel for spawning and incubation, but does not overwhelm the system, raising the riverbed and increasing channel instability. A stable incubation environment is essential for coho and other salmon, but is highly dependent on the functioning of nearly all habitat components in an ecosystem.

As with other salmonids, the length of time required for coho eggs to incubate is largely dependent on water temperature, and to a lesser extent, on dissolved oxygen content. The colder the temperature, the slower the development rate; however, for a given temperature, hatching time often differs between eggs from different fish or even different eggs from the same fish (Shapovalov and Taft 1954). Consequently, a wide range of values have been reported for the mean number of days that elapsed from fertilization to hatching for coho salmon. Data indicates that coho require approximately 300-450 temperature units (TU) for hatching. In general, the hatching time is shorter for North American stocks than Asian stocks, even at northern latitudes.

After hatching, coho alevins migrate downward in the gravel; migration distance is related to gravel size (Dill 1969). If the gravel was ≤ 1.25 in (3.2 cm) in diameter, alevins migrated down about 4 in (10 cm). If the gravel diameter ranged from 1.25 to 2.5 in (3.2 to 6.3 cm), the alevins migrated down more than 8 in (20 cm). This migration appears to be an adaptation to prevent premature emergence of alevins located close to the gravel surface.

For coho salmon on the Big Qualicum River, Fraser et al. (1983) documented the total heat requirement for fry emergence was $1,036 \pm 138$ degree ($^{\circ}$ C) days. The total time from egg deposition to fry emergence averaged 167 days (range 149-188 d). In three Oregon coastal streams, Koski (1966) observed that the average time from egg deposition to fry emergence was 110 days (range 104-115 d).

Most mortality from the egg to fry stage is a result of winter flooding and the associated disruptive gravel movement. Coho have high egg to fry survival compared to other salmonids; Neave (1949) attributes this high survival to the selection of better spawning sites in areas of good flow stability and less crowding. Under average conditions, approximately 15-27% of eggs will survive to emergence (Neave 1949, Crone and Bond 1976). In three Oregon coastal streams, Koski (1966) reported that egg to fry survival ranged from 0-78% and averaged 27.1%. Also, survival to emergence was positively correlated with gravel sizes >0.1 in (3.35 mm) and <1 in (26.9 mm). However, survival decreases if the gravel bed has a high concentration (up to 50%) of fine sediment and sand (Tagart 1984). Furthermore, if the gravel is heavily compacted with fine sediment, fry may not be able to get out of the gravel (Koski 1966); when the gravel/sand mixture was 70% sand, survival to emergence was only 8% (Phillips et al. 1975).

Spawn timing affects time of fry emergence. In the lower Columbia River, peak spawning time for early run coho (Type S) is in late October, and for late run coho (Type N), peak spawning is generally from December to early January. In the Washougal River system, coho spawn from mid-October and continue through November. Incubation extends from late October through January, with emergence occurring in late January and early February (WDF 1990). On the Cowlitz River, fry emergence occurs from January to April (WDW 1990). In the Cowlitz and Lewis rivers at 50° F (10° C), fertilization to eyed-

egg stage takes about 3.5 weeks, eyed-egg to hatching about 2.5 weeks, and hatching to emergence about 8 weeks (Howell et al. 1985). On the Wind and Little White Salmon rivers, coho fry emerge in late winter/early spring, generally from mid-January to February (WDW 1990). These patterns likely typify the incubation timing occurring in other Washington tributaries to the lower Columbia River.

Freshwater Rearing

After emergence, coho fry move to shallow, low velocity rearing areas, primarily along the stream edges and in side channels. They congregate in quiet backwaters, side channels, and small creeks, especially in shady areas with overhanging branches (Gribanov 1948). The vast majority of coho juveniles remain in the river for a full year after leaving the gravel. They often rear in the same habitat areas as Chinook, either commingled with or entering an area as the Chinook fry are leaving. Stein et al. (1972) observed that coho juveniles at the head of riffles were able to defend the area against Chinook fingerling. Although juvenile coho are found in both riffle and pool habitat, they are best adapted to holding in pools (Hartman 1965). They do not compete well with trout for rearing space in riffles. Godfrey (1965) noted that coho fry from 1.5-1.75 in (38-45 mm) may migrate upstream considerable distances to reach lakes or other rearing areas; in lakes, coho fry generally occupy the nearshore littoral zone (Mason 1974). However, the majority rear in streams. As they grow, the juveniles move into faster water and disperse into tributaries and areas that adults cannot access (Neave 1949). Coho fry are active during daylight hours and seem to tolerate a wide range of light intensities; this adapts coho well to the small, shallow streams they normally occupy where light conditions are highly variable (Hoar 1958).

The two most important factors for coho freshwater survival are water discharge rate and temperature. There is a correlation between summer flows and the catch of adult coho salmon 2 years later (Neave 1948, 1949, Smoker 1953). During summer months, the amount and quality of juvenile rearing habitat can decline due to low flows and high water temperatures. This may lead to a physical reduction of available habitat, increased stranding, decreased dissolved oxygen, and increased predation (Cederholm and Scarlett 1981). Coho fry production also has been shown to be a function of the stability of winter flows (Lister and Walker 1966, Seiler 2003).

The most productive coho streams are those with alternating pools and riffles of about equal area (i.e. 1:1 pool to riffle ratio; Ruggles 1966). Invertebrate production is maximized in the riffle area and the pool habitat is the optimum environment for coho fry holding and feeding (Mundie 1969). Coho tend to be more aggressive in defense of their territories where the current is fast and most of the available food is coming from upstream. In areas where the current is slow or slack, the food can appear from any direction and coho tend to move in loose aggregates while scrambling for food (Mundie 1969). As coho juveniles grow into yearlings, they become more predatory on fry of their own or other species (Gribanov 1948).

While juvenile coho are highly territorial and can occupy the same area for a long period of time (Hoar 1958), coho abundance can be limited by the number of suitable territories available (Larkin 1977). McMahon (1983) determined that pools of 10-80 m³ (353-2825 ft³) in size were optimum for coho production, provided there was adequate shading from streamside vegetation; however, if the canopy is very dense, then coho biomass will be reduced (Chapman and Knudsen 1980). Streams with more structure (logs, undercut banks, etc.) support more coho (Scrivener and Andersen 1982), not only because they provide more territories (useable habitat), but they also provide more food and cover. There is a positive correlation between the amount of terrestrial insect material in coho stomachs and the extent the stream was overgrown with vegetation (Chapman 1965). In addition, the leaf litter in the fall contributes to aquatic insect production (Meehan et al. 1977).

Coho fry are continually displaced downstream by freshets throughout the active juvenile growth period (Fraser et al. 1983). If the downstream area is unoccupied, displaced fry may take up residence;

however, if fry already occupy the space, displaced fry will be displaced further downstream (Ruggles 1966). Evolutionarily, the displacement may distribute fry far from the spawning grounds, allowing them to make more effective use of the available habitat (Allen 1969). However, in many cases, fry are displaced to less favorable sites, where they become more vulnerable to predators or are prematurely driven to the estuary. Some coho fry displaced downstream may migrate back upstream, or they may move along the shore in low salinity water and enter other streams to continue rearing (Otto and McInerney 1970). Of those coho fry that are displaced to the ocean or voluntarily choose to migrate to sea in their first year, survival to the adult stage is uncommon (Crone and Bond 1976). However, Otto (1971) points out that the type of estuary has a substantial bearing on the ability of coho fry to survive. Crone and Bond (1976) demonstrated that coho fry could survive salinities as high as 29 ppt if they had been acclimated at lower salinities first. Regardless, adult production from coho fry that enter the sea during their first year is expected to be very low.

Pool habitat is also important during all stages of juvenile development. Preferred coho pool habitat includes deep pools with riparian cover and woody debris. In the autumn as the temperatures decline and juvenile coho feeding activity decreases, juvenile coho move into deeper pools and hide under logs, tree roots, and undercut banks (Hartman 1965). The fall freshets redistribute them (Scarlett and Cederholm 1984), and over-wintering generally occurs in available side channels, spring-fed ponds, and other off-channel sites to avoid winter floods (Peterson 1980). The lack of side channels and small tributaries may limit coho survival in some areas (Cederholm and Scarlett 1981). Fall freshets may cause considerable downstream migration before suitable overwintering habitat can be found. For example, in Washington's Clearwater River, coho move as much as 24 miles (38 km) downstream before entering a tributary (Scarlett and Cederholm 1984). Coho also have been observed overwintering in lakes, such as Tenmile Lake in Oregon.

High summer water temperatures can exceed the 77°F (25°C) upper lethal temperature for juvenile coho. Brett (1952) observed that juvenile coho preferred a temperature range of 54-57°F (12-14°C), which is close to optimum for growth efficiency. Rapid increases or decreases in temperature can also cause significant mortality in juvenile coho (Brett 1952). Most mortality in the fry stage occurs in the first summer. Godfrey (1965) reported an average fry to smolt survival of 1.27-1.71% for two British Columbia streams, one Washington stream, and one Oregon stream. British Columbia survival from egg to smolt has been estimated at 1-2% (Neave and Wickett 1953). In the Big Qualicum River, Fraser et al. (1983) estimated fry to smolt survival was 7.3%. The long freshwater residence time likely results in higher freshwater mortality, than juveniles with shorter freshwater residence time, but may contribute to a lower marine mortality because smolts are larger when they go to sea (Drucker 1972).

Juvenile Migration

Most juvenile coho, in the region south of central British Columbia, migrate seaward as smolts in late spring, typically during their second year. Factors that tend to affect the time of migration include; the size of the fish, flow conditions, water temperature, dissolved oxygen levels, day length, and the availability of food (Shapovalov and Taft 1954). The size of coho smolts is fairly consistent over the species' geographic range; a FL of 4 in (100 mm) seems to be the threshold for smoltification (Gribanov 1948). Generally, the timing of outmigration is earlier in the southern coho populations compared to northern populations. For example, coho smolt emigration in California starts as early as mid-March and peaks in mid-May (Shapovalov and Taft 1954), but in the Resurrection Bay area of Alaska, smolt migration begins in late May and has been observed into early September (McHenry 1981). In the lower Columbia River, coho smolt emigration likely occurs from March to June, with peak movement in April and May. In addition, the timing of smolt outmigration may respond to small-scale habitat variability, with smolts residing in ponds and lakes often having different outmigration timing and being a different

size than smolts residing in streams within the same basin (Swales et al. 1988, Irvine and Ward 1989, Rodgers et al. 1993, Nielsen 1994).

Changes in the environment can also cue coho smolts to migrate. For example, Tripp and McCart (1983) observed the main peak of coho emigration coincided with a time of maximum stream discharge. In addition, a second peak of migration was observed during a time of increasing water temperature. For a given river system, there are annual variations in emigration timing that are related to these environmental factors. Changes in a tributary created by habitat degradation, habitat restoration and/or flow control often influence outmigration timing and though the relationships are not yet clear. Nearshore ocean conditions can also affect the timing of smolt outmigration in some tributaries to the lower Columbia River (NMFS 1995).

Estuary Rearing and Growth

Coho use estuaries primarily for interim feeding while they adjust physiologically to salt water. However, connectivity of available feeding and refuge areas may be important for species such as coho that move quickly through the estuary. For example, radio tagged coho in Grays Harbor estuary moved alternatively from low velocity holding habitats to strong current, passive downstream, movement areas (Moser et al. 1991). Additionally, Dittman et al. (1996) suggest that habitat sequences at the landscape level may be important even for species and life history types that move quickly through the estuary during the important smoltification process, as salmon gather the olfactory cues needed for successful homing and these cues may depend on the environmental gradients experienced during migrations.

Juvenile coho salmon were present in the Columbia River estuary from March to August of each year of sampling by Bottom et al. (1984); coho abundance was greatest in May and June and relatively low for other months (Bottom et al. 1984). Juvenile coho salmon comprised 18% of the total juvenile salmonid catch (Bottom et al. 1984). Coho juveniles were distributed throughout the freshwater, estuarine, and marine regions of the estuary; they were most frequently associated with water column habitats, however, tagged hatchery coho released in the lower Columbia (i.e. Grays River (rm 34) and Big Creek (rm 29)) were more likely to be found in shallow bays and intertidal areas than upriver coho (Bottom et al. 1984). Juvenile coho salmon moved through the estuary relatively quickly and appeared to increase their migration rate through the estuary (Bottom et al. 1984). As with other salmonids, juvenile hatchery coho released further upstream in the basin migrated at a faster rate than juveniles released lower in the system (Bottom et al. 1984).

Recent sampling of juvenile salmonids in the Columbia River plume has started to illustrate patterns of habitat use by salmonids in the plume and nearshore ocean habitats (Fresh et al. 2003), although limited years of data are currently available. First, juvenile salmon distance offshore appears to be positively related to river flow as measured at Bonneville Dam; generally, Chinook and coho salmon yearling were captured further offshore in the plume environment as river flow increased (Fresh et al. 2003). Second, preliminary evidence suggests that some juvenile salmonids (chum, steelhead, and yearling coho) may preferentially utilize the plume front compared to other areas in the plume or adjacent ocean habitats (Fresh et al. 2003). Although reasons for the apparent preference to the plume front are not clear, this area may be a more productive habitat than elsewhere in the plume and adjacent ocean.

Ocean Migrations

Most research indicates that, upon entering the ocean, coho remain in nearshore environments over the continental shelf for a couple of months before they disperse on more seaward migrations; this holds true from California to Alaska (Shapovalov and Taft 1954, Milne 1964, Godfrey 1965). This pattern

may help coho avoid pelagic predators and reduce feeding competition with immature salmon that are older by a year or more.

Some Washington and British Columbia stocks migrate only short distances to good feeding areas and remain there until they approach maturity (Godfrey et al. 1975). In the Strait of Georgia, coho smolts quickly disperse throughout the strait; the number of coho that remain in the strait depends on coho density and feeding conditions (Healey 1980). If fish find themselves in poor feeding areas within the strait, they move to outside waters; however, those fish that locate good feeding areas remain within the Strait of Georgia.

Coho salmon typically spend 18 months in the ocean before returning to fresh water. Thus, many returning coho are 3 years old and have spent 18 months in fresh water and 18 months in salt water. Jacks, however, return earlier at age 2. These sexually mature males return to fresh water to spawn after only 5 to 7 months in the ocean.

Data collected from CWT recovery studies shows that coho salmon released from Columbia River hatcheries are recovered primarily in Oregon (36-67%) and Washington (22-54%), with lower but consistent recoveries from British Columbia (2-16%) and California (1-15%). These ocean distribution patterns were determined from CWT recovery data for 66 North American hatcheries between 1973–92 from the PSMFC's 1994 Regional Mark Information System. Compared to Oregon coast coho salmon, Columbia River fish are recovered less frequently in California and more frequently in Washington. Although they share the same general recovery pattern, coho salmon from Washington-side Columbia River hatcheries are caught more frequently in Washington and British Columbia, and less frequently in Oregon than are those from Oregon-side hatcheries. This is presumably the result of Washington hatcheries producing both Type S and Type N coho, while Oregon hatcheries produce only Type S coho. Washington has maintained both stocks in Columbia River hatcheries because early and late returning coho are indigenous to Washington streams and the mix of stocks provides fishing access off the Washington coast as well as in the Columbia River and Washington tributaries.

A.2.2. Distribution

Historically, coho were present in all lower Columbia River tributaries. Before the early 1800s, the watersheds remained essentially untouched by human development. Heavy growth of coniferous trees and understory vegetation armored many river stretches. The river systems' cool streamflows, clean gravel beds, and deep pools supported healthy populations of coho and other salmon and steelhead. These pristine environments began to change in the mid-1800s, often causing declines in salmonid production. Currently, very few wild coho salmon spawn annually throughout the lower Columbia River subbasins. Until recently, Columbia River coho salmon were managed as a hatchery stock. In some cases, coho salmon returning to Columbia River hatcheries above the brood stock needs for the hatchery are allowed to bypass the hatchery rack or collection facility and allowed to spawn naturally. Spawning is expected to occur in most areas accessible to coho, although production from naturally spawning hatchery fish is likely low.

Two general coho stocks are present in the lower Columbia River today: early (Type S) refers to an ocean distribution generally south of the Columbia River with an early adult run timing in the Columbia River; late (Type N) refers to an ocean distribution generally north of the Columbia River with a late run timing in the Columbia River. CWT data provides valuable information on fish distribution and harvest rate of stocks originating in Columbia River hatcheries; however, the fishery distribution does not necessarily reflect ocean migratory patterns as fishing effort levels vary in specific areas each year. CWT recoveries data are useful when describing catch distribution for a given time period. The recoveries illustrated by the following table represent fishery distribution during the late 1990s (Table A-18). These fishery recoveries reflect years when Oregon ocean fisheries were minor compared to Washington

ocean and Columbia River fisheries. Oregon fisheries were curtailed in the early 1990s in response to management of the presently ESA-listed Oregon Coastal natural coho.




Figure A-45. Distribution of historical coho salmon populations among Washington lower Columbia River subbasins.

Table A-18. Coho salmon CWT recoveries in various fisheries.

Hatchery	Sampling Areas			
	Columbia River	Oregon Ocean	Washington Ocean	California Ocean
<i>Early Coho</i>				
Grays River ¹	58%	21%	19%	1%
Elochoman ²	53%	7%	40%	—
North Toutle ²	47%	1%	37%	—
Fallert Creek (Kalama) ²	49%	9%	42%	—
Lewis River ²	21%	21%	58%	—
<i>Late Coho</i>				
Elochoman ²	59%	11%	29%	—
Cowlitz Salmon ³	55%	1%	30%	—
Kalama Falls ²	58%	10%	32%	—
Lewis River ²	56%	21%	31%	—
Washougal ²	57%	13%	30%	—

¹ 1994, 1996, and 1997 brood years; ² 1995, 1996, and 1997 brood years; ³ 1994 and 1997 brood years

A.2.3. Genetic Diversity

The physical environments that support West Coast coho salmon—and the life history traits and genetic characteristics exhibited by these fish—indicate a substantial degree of ecological and genetic diversity. These environments range from the relatively dry climate in central California with strong and consistent upwelling offshore, to the extremely wet Olympic Peninsula with its snow and rain-fed rivers.

While information on historical runs is scarce, individual coho runs to the different tributaries probably showed a great deal of flexibility within their range. These runs became more conformed through selection by hatchery and harvest practices. In the 1940s, two separate runs of coho were reported to enter the Cowlitz River. The early run entered the Cowlitz from late August–September, with a spawning peak in late October. The late run entered from October–March, with a spawning peak in late November. Further, the Toutle River, a tributary to the Cowlitz River, historically produced an early-returning stock, with most fish returning from August–October. These early Toutle River coho are generally more southerly distributed in the ocean than the early component of the Cowlitz stock (WDW 1990).

Genetic diversity has largely been lost in the lower Columbia River because of widespread hatchery production with many out-of-basin (but mostly within-ESU) stock transfers. In the 1950s, salmon hatchery construction expanded on the lower Columbia River tributaries and hatcheries began to trap brood stock. Over time, transferring brood stock, eggs, and juvenile coho between hatcheries and planting hatchery fish off-station became commonplace throughout the watershed, resulting in a widely-mixed coho stock (WDF et al. 1993).

A.2.4. ESU Definition

In 1995, the BRT formed by NMFS concluded during its coho status review that, historically, there probably was an ESU that included coho salmon from all tributaries of the Columbia River below the Klickitat River on the Washington side and below the Deschutes River on the Oregon side. This ESU also would have included coho salmon from coastal drainages in southwest Washington between the Columbia River and Point Grenville (between the Copalis and Quinault rivers). The team based its determination in part on the similarities between the different physical environments. The Columbia River estuary, Willapa Bay, and Grays Harbor all have extensive intertidal mud and sandflats and similar estuarine fish faunas, and they differ substantially from estuaries to the north and south. Their similarity results from the shared geology of the area and the transportation of Columbia River sediments northward along the Washington coast. Other commonalities include:

- moving west to east, rivers that drain into the Columbia River have their headwaters in areas that are increasingly dry.
- Columbia River tributaries that drain the Cascade Mountains have proportionally higher flows in late summer and early fall than rivers on the Oregon coast.

Genetic analysis conducted since the 1995 status review, however, have led the BRT to conclude that SW Washington coho salmon form their own ESU, separate from LCR coho salmon. Geneticists from NMFS collected and compared allozyme data over 10 years from more than 100 coho salmon samples from locations ranging from California to Alaska, with a primary focus on Oregon, Washington, and southern British Columbia. Results from the study showed regional patterns of allele frequency.

The samples arranged into several different clusters. This cluster analysis placed SW Washington coast coho in a different cluster from LCR coho salmon, which form a supercluster with Oregon coast fishes (NMFS 2001). Within the SW Washington and LCR clusters, several subclusters and three branches have only one or two members.

The NMFS BRT concluded it could not identify any remaining natural populations of coho salmon in the lower Columbia River (excluding the Clackamas and Sandy rivers) or along the Washington coast south of Point Grenville that warrant protection under the ESA.

A.2.5. Life History Diversity

Little is known about the specific life history traits of wild coho salmon within the Washington tributaries of the lower Columbia River. There does not appear to be many significant life history differences among ‘same type’ coho stocks in the Washington subbasins of the lower Columbia River. Two general coho stocks are present in the lower Columbia River today; Type S (early) refers to an ocean distribution generally south of the Columbia River with an early adult run timing in the Columbia River and Type N (late) refers to an ocean distribution generally north of the Columbia River with a late run timing in the Columbia River. For early coho in the Washington subbasins of the lower Columbia, migration timing is generally from mid-August to September and peak spawning occurs in late October. For late coho stocks, migration timing is generally from late September to October and peak spawning occurs in December to early January. Dominant adult age class for lower Columbia coho is 1.2, indicating 1 year in fresh water and 2 years in salt water. Any natural spawning that does occur is thought to happen in all areas accessible to coho; specific spawning areas for each subbasin are noted in the Subbasin Chapters in Volume II.

For both stocks, fry emerge in the spring, spend 1 year in fresh water, then migrate to sea during the spring of their second year. In the Cowlitz River, there appears to be a late-late run component that may be an artifact of hatchery practices. Late stock hatchery programs throughout the lower Columbia have always taken brood stock for the late run from the early portions of the run. Once the annual brood stock needs were met, hatchery collection efforts ceased, leaving the later component of the run to spawn naturally. Therefore, a remnant late-late run from natural production exists in the Cowlitz basin and it is possible that other similar runs exist elsewhere in the lower Columbia.

There appears to be a significant difference in average length between the males and females of Columbia River hatchery coho stocks (Jim Haymes, WDFW, personal communication). Based on CWT-marked coho recovered at hatcheries on the Washington side of the Columbia River, the females are larger in Type N coho, and to a lesser extent, in Type S coho. This length difference is not seen in other Washington hatchery coho stocks – if anything, the females tend to be smaller at most hatchery racks (Jim Haymes, WDFW, personal communication).

A.2.6. Abundance

Since 1970, the Columbia River produced adult coho ocean population has ranged from a low of 96,700 in 1996 to over 3 million in 1971. The returns to the Columbia River mouth (after ocean harvest) have ranged from 74,800 in 1995 to over 1.5 million in 1986 (Figure A-46).

While records of coho escapement from the 1940s and 1950s report runs as high as 77,000 coho to the Cowlitz River (WDF/WGC 1948) and 15,000 coho to the Lewis River, such large natural returning runs to lower Columbia River tributaries are now gone. Estimated coho escapement to other lower Columbia River systems in the early 1950s was generally in the range of a few thousand coho: Grays River (2,500), Kalama River (3,000), and the Washougal River (3,000). Today, coho stocks in these and other lower Columbia River tributaries in Washington are considered depressed, primarily because of chronically low escapement and production. Natural spawning is presumed to be quite low in most areas, and subsequent juvenile production is below stream potential. Much of this natural production has been replaced by hatchery production.




Figure A-46. Columbia River coho ocean and freshwater run size.

Natural coho production is now being reintroduced in some once productive habitat areas. For example, in the Cowlitz system, coho are now reseeding the highly productive habitat in the upper basin above the hydroelectric projects through a salmon reintroduction program. In 1994, a trap and haul program began with adults returning to the Cowlitz Falls Fish Collection Facility near Mayfield Dam; adult coho have been released in the upper Cowlitz, Cispus, and Tilton rivers and allowed to spawn naturally. Nearly 1 million coho smolts were estimated to have migrated out of the upper basin in the spring/summer of 2001. This shows that the upper river is still capable of producing significant numbers of fish, however, only a portion of the migrating smolts (375,000) were subsequently trapped and hauled below the dams. Downstream passage of juvenile coho continues to be a challenge for this reintroduction effort. Estimates of coho smolt production potential in the Cowlitz River basin above the dams range from 6,319 (Stockley 1961) to 261,254 fish (Easterbrooks 1980). In the Lewis, reintroduction of coho to the upper basin above the hydro-system is also being considered.

A.2.7. Productivity

The productive capacity of the freshwater environment for coho has been estimated by a number of researchers. In the Big Qualicum River, Lister and Walker (1966) determined that 19.1 smolts were produced per 100 m² of wetted stream area measured at low flow. In three Oregon coastal streams, Chapman (1965) reported 18-67 smolts produced per 100 m² over a 4-year period. In low gradient side channels of the Cowichan River, Armstrong and Argue (1977) determined that 125-141 smolts were produced per 100 m². In an evaluation of coho smolt production in ten Western Washington streams, average smolt production per square mile ranged from 417 to 1,798; additionally, average smolt production per mile² in select lower Columbia tributaries from 1997 to 2002 ranged from 17 to 765 (Jim Haymes, WDFW, personal communication). In a study of hatchery coho in isolated headwater streams in Canada, Tripp and McCart (1983) found that the average production was 8.4-8.5 smolts per 100 m²; this low production may be a result of hatchery smolt fitness or because high-gradient headwater streams are not usually productive areas. The average coho production likely falls between these extremes reported in the literature. In addition, smolt production in streams is 7-10 times greater than lakes (Foerster and Ricker 1953).

The NPCC’s smolt density model was run on many subbasins within the lower Columbia River to estimate potential coho smolt production. Estimates of coho smolt production for Washington subbasins include: 125,874 for the Grays River, 43,393 for the Elochoman, 123,123 for the lower Cowlitz, 131,318 for the Tilton River and Winston Creek, 155,018 above Cowlitz Falls, 142,234 for the Toutle, and 37,797 for the Coweeman.

A.2.8. Hatchery Production

In past years, hatchery production of coho salmon in the lower Columbia River/southwest Washington coast ESU has far exceeded that of any other area with respect to the number of hatcheries and quantities of fish produced. Many hatcheries within this ESU released 1-3 million smolts annually, with the two largest hatcheries, Cowlitz and Lewis, releasing an average of 6-7 million smolts (Table A-19). However, in recent years, Washington lower Columbia hatchery programs have reduced coho production, either as a result of reduced funding, reprogramming federal-funded production to facilities above Bonneville Dam, or mitigation adjustments. The current annual release goal for 2003 brood totals 9.7 million yearling smolts (Table A-20).

Table A-19. Average annual releases of coho salmon juveniles (fry and smolts) from Washington lower Columbia hatchery facilities during release years 1987–91 (NMFS 1995).

Hatchery	5-Year Average
Washougal	3,885,612
Lewis	6,180,000
Kalama Falls	990,000
Fallert Creek (Lower Kalama)	831,605
Toutle	478,090
Cowlitz	7,956,089
Elochoman	2,013,032
Grays River	744,655
Sea Resources	125,500
Total	23,204,583

Table A-20. Current (2003 brood) annual release goals of coho salmon smolts from Washington lower Columbia hatchery facilities.

Basin	Brood Source (Hatchery)	Early (Type S)	Late (Type N)	Total
Little White Salmon	Little White Salmon	1,000,000		1,000,000
Washougal	Washougal		500,000	500,000
NF Lewis	Lewis River	880,000	815,000	1,695,000
Kalama	Fallert Creek (S)	350,000	350,000	700,000
	Kalama Falls (N)			
NF Toutle (Green R.)	NF Toutle	800,000		800,000
L. Cowlitz	Cowlitz Salmon		3,200,000	3,200,000
Elochoman	Elochoman	418,000	512,000	930,000
Columbia (Steamboat slough)	Grays River	200,000		200,000
Grays	Grays River	150,000		150,000
Deep River	Grays River	400,000		400,000
Chinook	Sea Resources	52,500		52,500
TOTAL		4,250,500	5,377,000	9,627,500

Extensive stock transfers have occurred within the Lower Columbia River/ Southwest Washington Coast Coho ESU. Most transfers of coho salmon have used stocks from within the ESU, although transfers from outside the ESU have also occurred, including those from the Oregon coast, Olympic Peninsula, and Puget Sound/Strait of Georgia ESUs. Outplanting records show a similar pattern to transfers between hatcheries, with extensive use of within-ESU stocks, in addition to less frequent use of stocks from the other three ESUs. Most movement of coho salmon, either as hatchery transfers or off-station releases, has occurred within each of the three areas of this ESU (Oregon-side Columbia River, Washington-side Columbia River, and southwest Washington coast), with little movement of fish among the three areas (NMFS 1995). There has been liberal exchange of early and late stock coho among hatcheries on the Washington side, with the exception of the Cowlitz Hatchery which has maintained the original late stock without transfers into the program. On the other hand, Cowlitz Type N stock coho have been used widely in several Washington lower Columbia tributary hatchery programs.

Because of past hatchery practices, many coho stocks in Washington-side tributaries of the lower Columbia are now considered mixed and of composite production. In the once productive Cowlitz River basin, for example, DeVore (1987) accounted for the 1982-brood hatchery release and concluded wild/natural production was minor. Of the 4,635 naturally spawning coho in the Cowlitz in 1985, an estimated 91% were from hatchery smolt releases. Hatchery coho have been planted in the Cowlitz since at least 1915, when the Tilton River Hatchery operated downstream of Morton until 1921. Stock mixing probably began in 1915 (DeVore 1987). Since 1968, the Cowlitz Salmon Hatchery has been producing coho salmon. The mitigation goal is to maintain annual returns of 25,500 coho adults to the hatchery.

Many streams, especially those downstream of the Cowlitz River, have had years of hatchery production with an earlier-timed stock than historically spawned in each river. For example, the Grays and Elochoman hatcheries have maintained an early Type S hatchery program with fish primarily originating from Toutle River stock, although the natural returns to these rivers were principally fish that spawned from late November to March. The effect of introducing an early coho stock to a basin that historically supported a late coho stock (or vice versa) is not completely understood; there may be

little or no interaction between early hatchery stock and any remnant late stock because of the temporal segregation of the runs.

There may be some benefit to the way late stock production has historically been managed. Hatchery programs have been operated to take eggs prior to mid-December to assure adults produced would be accessible to most of the freshwater fisheries. Because of this practice to maximize exposure to the fisheries, there could be a late spawning remnant of natural production which has not been significantly mixed with hatchery fish nor subjected to the same harvest pressure as the earlier-timed fish. This appears to be the case in the Cowlitz basin as there is documentation of a “late” late coho run. These fish could be extremely important to future recovery efforts. Currently, as part of the reintroduction program above Cowlitz Falls Dam, adult coho are being released into the upper Cowlitz River basin.

Historical juvenile hatchery coho salmon releases by basin have generally ranged from 1-5 million annually (Figure A-47). In recent years, releases into the Cowlitz River have exceeded releases in other basins and have averaged about 5 million juveniles annually, including coho reintroduced to the Upper Cowlitz above the dams. Grays River Hatchery coho production has declined substantially from 1970s and 1980s production levels.

Since 1982, adult coho returns to Washington hatcheries below Bonneville Dam have ranged from 4,759 (1987) to 91,407 (2001) for early stock, and from 11,776 (1995) to 177,941 (2001) for late stock (Figure A-49). During 1978–2002, hatchery releases at the Little White Salmon NFH peaked at approximately 3.7 million in 1984. The lowest number of releases from this hatchery was approximately 750,000 in 1993 (Figure A-48).

Coho Salmon Hatchery Releases in the Lower Columbia River by Basin, 1967-2002




Figure A-47. Hatchery releases of coho salmon in the lower Columbia River by basin, 1967–2002.

Coho Salmon Hatchery Releases by Brood Year from the Little White Salmon NFH 1980-2002




Figure A-48. Hatchery releases of coho salmon by brood year from the Little White Salmon NFH, 1980–2002.




Figure A-49. Columbia River adult coho escapement to hatcheries.

From 1983 to 1992, the average annual hatchery escapement to the Cowlitz River was 28,572, and to the Cowlitz Salmon Hatchery was over 75,000 coho in 2001 and 2002 (Figure A-50). The Cowlitz Salmon Hatchery produces Type N (late) coho while the North Toutle Hatchery produces Type S (early) coho. In 1980, some North Toutle Type S coho strayed to Cowlitz Hatchery as a result of the eruption of Mt. St. Helens.




Figure A-50. Hatchery returns of adult coho salmon in the Cowlitz River basin by hatchery and stock type, 1960-2002.

Significant coho production also has occurred in the Lewis, Kalama, and Washougal River basins.

- The Lewis River Hatchery produces both Type S and Type N coho while the Speelyai Hatchery produces Type S coho (Figure A-51). The largest annual hatchery coho return to the Lewis River basin was over 95,000 adult fish in 1999.
- In the Kalama basin, the Kalama Falls Hatchery was the primary producer of coho salmon with the largest annual return of over 40,000 adult coho in 1966 (Figure A-52). In recent years, similar-sized hatchery returns of Type S coho have been documented at the Kalama Falls and Fallert Creek hatcheries.
- In the Washougal basin, historical production at the Washougal Hatchery was Type S coho; the largest annual hatchery return was approximately 45,000 coho in 1968 (Figure A-53). In recent years, Washougal Hatchery production has shifted to Type N coho and annual adult returns have averaged about 5,000 fish.




Figure A-51. Hatchery returns of adult coho salmon in the Lewis River basin by stock type, 1950–2002.




Figure A-52. Hatchery returns of adult coho salmon in the Kalama River basin by stock type, 1951–2002.




Figure A-53. Hatchery returns of adult coho salmon in the Washougal River basin by stock type, 1956–2002.

No consistent pattern is apparent in historical hatchery coho returns to the Grays or Elochoman rivers (Figure A-54 and Figure A-55); generally, hatchery returns in the 1960s and 1970s were higher than those in the 1980s and 1990s.




Figure A-54. Hatchery returns of adult coho salmon in the Grays and Chinook River basins by stock type, 1961–2002.




Figure A-55. Hatchery returns of adult coho salmon in the Elochoman River basin by stock type, 1954–2002.

A.2.9. Fishery

Impacts to lower Columbia River coho salmon are harvested in ocean commercial, sport, and tribal fisheries; in Columbia River sport, commercial, and treaty Indian fisheries; and in tributary sport fisheries. These fisheries, and their management structure, are briefly discussed in the harvest overview section of this report. Like other salmon stocks in the Columbia River, integrating the management of coho ocean and Columbia River fisheries is essential to meeting conservation requirements for ESA-listed or critical stocks and to promote fishery opportunity on healthy hatchery and wild populations. Inside the Columbia River, early and late stock coho are managed separately; differences in the timing of fish runs enable managers to structure seasons to meet separate harvest objectives for the stocks.

Coho Harvest Over Time

Coho salmon received significant harvest pressure beginning in the late 1800s, particularly on the lower Columbia River. Peak commercial catches of wild coho in the Columbia River occurred in 1925 (Lichatowich and Mobernd 1995); since the 1960s, Columbia River commercial catch has consisted primarily of hatchery-produced coho. Commercial landings of coho salmon in Washington, Oregon, and California from 1882 to 1982 have been estimated by Shepard et al. (1985). These estimates show relatively constant landings since 1895, ranging mainly between 1.0 and 2.5 million fish, with a low of 390,000 fish (1920) and a high of 4.1 million fish (1971). Columbia River coho became an important marine, as well as freshwater, harvest species in the 1960s.

Ocean harvest of coho in the Oregon Production Index (OPI) peaked in the 1970s and early 1980s (Figure A-56) and resulted in high coho exploitation when combined with freshwater fisheries aimed at harvesting large hatchery production (Figure A-57). For example, ocean and Columbia River combined harvest rates of Columbia River-produced coho ranged from 70 to 90% during 1970-1983. During this

time, naturally produced coho were managed like hatchery stocks and were subject to similar harvest rates. In the mid-1980s, ocean fisheries harvest was reduced to protect several Puget Sound and Washington coastal wild coho stocks. Beginning in the early 1990s, Columbia River coho commercial seasons were closed before November to reduce harvest of late Clackamas River wild coho. Coho in the Oregon Coast ESU were listed as threatened under the ESA in 1998; subsequent harvest restrictions to protect Oregon Coast coho likely also benefited naturally produced lower Columbia River coho.




Figure A-56. Coho ocean harvest rate based on Oregon Production Index.




Figure A-57. Approximate coho fishery exploitation rates over time.

Coho were subject primarily to Columbia River harvest until 1950s. Ocean harvest peaked 1970s–80s. Coho remain an ocean sport fishery focus. Sport harvest in Lower Columbia estuary began to be significant in 1980s. Columbia commercial harvest focused on late September–October. Differential harvest of wild fish commenced in 1960s when late fall fisheries were reduced. Selective harvest in ocean and Columbia began in 1998 and provided greater differences in wild and hatchery harvest rates.

Beginning with the 1995 brood, most Columbia River hatcheries mass marked hatchery-released fish with an adipose fin clip. Since marked fish began returning as adults in 1998, fisheries managers have been able to prosecute selective sport fisheries for marked hatchery coho where all unmarked fish were required to be released. In addition, because there are run timing differences between some hatchery and wild stocks, Columbia River commercial fisheries have employed select area and time strategies to target hatchery fish to reduce impacts on wild coho. As a result of these selective management strategies employed during 1998–2002, combined fisheries harvest of ESA-listed coho was less than 15% annually, while harvest of Columbia River hatchery coho was maintained near 50 percent.

Recent harvest management practices have resulted in greater commercial harvest of late hatchery coho compared to early coho (Figure A-58). Peak migration time for early coho in the Columbia River is September; harvest of early coho is currently restricted because of harvest constraints on fall Chinook and Sandy River wild coho which also migrate during September. Columbia River commercial coho harvest effort is concentrated in October during the peak migration of late hatchery coho; there are no concurrent harvest restrictions for other salmonids during this period.




Figure A-58. Columbia River Zone 1-5 commercial harvest of early and late stock coho.

Current Coho Harvest Distribution

Lower Columbia wild coho returning to Oregon tributaries were placed on the Oregon State Endangered Species List in 1999. Impacts to Oregon state listed coho have been managed under an abundance-based management plan similar to OCN coho.

CWT data analysis of hatchery coho from the mid-to late 1990s brood years consistently show greater harvest percent age of late coho with less percent age accounted for in the escapement compared to early coho. For example, CWT analysis of Fallert Creek (lower Kalama) early coho from the 1995–1997 brood years indicated that 30% were captured in a fishery and 70% were accounted for in escapement. However, 76% of Kalama Falls late coho from the 1995–1997 brood years were captured in a fishery and 24% were accounted for in escapement. In the Cowlitz basin, 34% of Toutle Hatchery early coho from the 1995–1997 brood years were captured in a fishery while 66% were accounted for in escapement. Meanwhile, 64% of Cowlitz Hatchery late coho from the 1994 and 1997 brood years were captured in a fishery while 36% were accounted for in escapement.

CWT data also provide some insight into the general distribution of fish. CWT hatchery coho from the 1995–1997 brood years, regardless of whether they were early or late coho, were primarily (50-60%) recovered in the Columbia River sampling area, 20-40% were recovered in the Washington ocean sampling area, and 10-20% of coho CWT recoveries were reported in the Oregon ocean sampling area. The one notable exception to this pattern is early coho from the Lewis River Hatchery; 58% CWT were reported in the Washington ocean, 21% in the Columbia River, and 21% in the Oregon ocean. In general, lower Columbia River hatchery and wild coho are harvested in West Coast ocean or Columbia River sport or commercial fisheries (Table A-21).

Columbia River coho do not migrate as far north as Columbia River Chinook; consequently, few Columbia River coho are harvested in Alaska or Canadian fisheries. Commercial ocean troll fisheries typically focus on Chinook, but Indian and non-Indian ocean troll coho harvest can be significant in years of large hatchery abundance. Selective fisheries for adipose fin-marked hatchery coho have been implemented in most PFMC area ocean fisheries since 1998.

Table A-21. Example of lower Columbia coho harvest exploitation and distribution under current management (combined early and late stock).

Fishery	H*	W*	Comment
Alaska	0%	0%	Do not typically migrate far north
Canada	<1%	<1%	Constrained by PSC and Thompson coho management
WA/OR/CA/Ocean	30%	9%	Selective sport and troll fisheries
Columbia River	15%	8%	Sport selective, commercial time and area restricted
WA Tributaries	6%	1%	Sport selective
Total Exploitation	51%	18%	Late stock hatchery harvest rate higher than early stock hatchery harvest rate. Wild stock that enter the Columbia November and later have a lower harvest rate.

* H=Hatchery, W=Wild

The Oregon production index (OPI) area coho stocks include all Washington, Oregon, and Northern California natural and hatchery stocks from streams south of Leadbetter Point, Washington. Historically, OPI stocks contributed primarily to Oregon and northern California ocean fisheries and, to a lesser extent, ocean fisheries off Washington and British Columbia. In recent years, more of the coho harvest has shifted to southern Washington coastal fisheries as a result of management actions aimed at reducing impacts to Oregon coastal natural coho stocks. The largest naturally produced component of the OPI area coho stock is Oregon coast natural (OCN) coho, which is managed as an aggregate stock with four identified components from Oregon systems south of the Columbia River. There are three threatened ESUs within the naturally produced OPI coho stocks; central California coast (CCC) coho (1996), southern Oregon/northern California (SONC) coho (1997), and Oregon coast (OC) coho (1998). OPI area coho harvest is driven by harvest restrictions on these listed stocks. No directed coho fisheries are allowed in any commercial and recreational fisheries off the California coast to protect threatened

CCC coho. Marine fishery impacts on threatened CCC and SONC coho must be no more than 13% based on projected impacts on Rogue/Klamath hatchery coho. Combined marine and freshwater impacts on OCN coho should not exceed levels in the abundance-based fishery management plan (15% in recent years).

Ocean commercial troll and recreational fisheries south of Cape Falcon (near Cannon Beach) have been closed to unmarked coho retention since 1993. Selective commercial troll fisheries for marked coho have occurred since 2000 in areas from Cape Falcon to the Queets River in Washington; all directed coho recreational fisheries in the OPI area have been selective for hatchery coho since 1998. Terminal recreational harvest in Oregon coastal systems is limited to areas where surplus hatchery returns are expected. Improved hatchery coho populations in the OPI area have expanded hatchery coho ocean commercial and recreational harvest opportunities in recent years (Figure A-59).




Figure A-59. Harvest of Oregon Production Index Coho in the ocean and Columbia River, 1970–2000.

In the Columbia River, numerous commercial coho fisheries still exist, including non-Indian commercial harvest in the lower river as well as treaty Indian commercial harvest in Zone 6 above Bonneville Dam (Figure A-60). Columbia River commercial coho fisheries are limited by Chinook constraints in the early fall season, which results in limited early stock coho harvest. Most commercial coho harvest is focused in late September and October when late stock hatchery coho abundance is highest (Figure A-58). Late fall seasons, primarily in Zone 3, target coho in the lower river below the mouth of the Lewis River. Columbia River commercial fisheries are closed before November to avoid harvest of late wild Clackamas coho, chum, and winter steelhead. Columbia River commercial fisheries retain all coho, but are managed by time and area to reduce impacts to wild fish. Columbia River commercial harvest of coho was low during the 1990s but has increased in recent years because of improved hatchery coho populations; coho harvest in treaty Indian fisheries in the Columbia River has generally been low (Figure A-60).

Coho Salmon Harvest in Columbia River Commercial and Subsistence Fisheries, 1970-2003




Figure A-60. Commercial and subsistence harvest of coho salmon in the Columbia River from 1970–2001.

Columbia River hatchery coho are very important to the Lower Columbia estuary (Buoy 10), mainstem, and tributary sport fisheries (Figure A-61). Selective fisheries for adipose-marked hatchery coho have been implemented in Columbia River and tributary sport fisheries since 1998, except in fisheries above Bonneville Dam. A substantial Columbia River estuary sport fishery exists between Buoy 10 and the Astoria-Megler Bridge; harvest is primarily early run coho, however, harvest of late coho can also be substantial (Figure A-62). Angler trips to the Buoy 10 fishery in years of high hatchery coho abundance can exceed 150,000 during August and September.

Tributary sport fisheries for coho also occur in many basins throughout the lower Columbia (Figure A-63). Data from the late 1980s indicate that average annual harvest was over 1,000 coho in some tributaries (e.g. 1,183 in the Elochoman [1981–88], 1,494 in the Cowlitz [1986–90], 1,272 in the Kalama [1979–86], and 3,500 in the NF Lewis [1980–98]).




Figure A-61. Recreational harvest of coho salmon in the Lower Columbia River for 1970–2001.




Figure A-62. Recreational harvest of Columbia River early/late stock adult coho in Columbia River estuary.




Figure A-63. Recreational harvest of Columbia River early/late stock coho in Washington tributaries of Columbia River.

Coho Harvest Management Details

PSC Fisheries

Coho salmon management and harvest in Alaska and Canada are governed by the 1999 Letter of Agreement (LOA) negotiated as part of the PST. The LOA specifies provisions for in-season conservation and information sharing for northern boundary coho salmon. The LOA specifies catch-per-unit-of-effort (CPUE) levels in Southeast Alaska commercial troll fisheries that trigger conservation measures. The LOA also contains total commercial harvest levels in July that trigger early region-wide troll closures. Targeted coho fisheries in Canada are currently limited to southern British Columbia; the 2002 management objective for all Canadian fisheries was to limit the total exploitation rate on Thompson River coho (Canadian population) to 3% (Table A-21).

PFMC Ocean Fisheries

Coho and Chinook are the two primary salmonid species harvested in Pacific Coast ocean fisheries occurring in PFMC managed waters, extending from the Canadian border to Mexico, and 3-200 nautical miles offshore. The PFMC STT annually publishes stock-specific preseason run forecasts provided by state and tribal biologists. These forecasts shape fishery management planning and harvest targets for the coming year; forecasts are presented annually in the PFMC Preseason Report 1 in February. The majority of coho harvested in U.S. ocean fisheries originate from rivers within the Oregon Production Index (OPI) area; stocks include hatchery and natural production from the Columbia River, Oregon coast, and northern California. The individual stock components originating in the OPI area in which abundance is estimated annually include: 1) public hatchery (OPIH), 2) Oregon coastal natural river (OCNR), 3) Oregon coastal natural lake (OCNL), 4) private hatchery (PRIH), and hatchery smolt production from the Oregon coastal Salmon Trout Enhancement Program (STEP).

The OPI area public hatchery stock is composed primarily of production from Columbia River facilities and net pens, with lesser contribution from facilities in Oregon coastal rivers and the Klamath River basin. OPIH forecasts are generated using multiple linear regression methods and a relationship

established between coho jacks and the subsequent year’s returning adults for the major stock components (i.e. Columbia River, Oregon coastal, Klamath River). The OPIH stock predictor is partitioned into Columbia River early and late stocks, and coastal stocks north and south of Cape Blanco, Oregon.

Integrated management of ocean and Columbia River coho fisheries is essential to the conservation and recovery of federal and state ESA-listed coho stocks. Therefore, fishery planning and management actions by PSC, PFMC, and the Columbia River Compact must be consistent and complementary. Federal ESA-listed coho stocks driving management and harvest constraints include the Oregon coast (OCN) ESU, southern Oregon/ northern California coasts (SONC) ESU, and central California coast (CCC) ESU; Oregon state-listed lower Columbia River wild coho (LCN, a federal candidate species) limits coho fisheries harvest. OCN and LCN coho are assumed to have similar temporal and spatial distributions in ocean fisheries. Harvest limits on OCN coho therefore benefit LCN coho.

In 1997, PFMC approved an amendment to the Fishery Management Plan that changed the basis for coho fisheries management from spawner escapement objectives to exploitation rate limits. The maximum allowable exploitation rates for OCN vary in response to changes in observed brood year-specific parental spawner abundance and marine survival conditions (Table A-22). Similar exploitation rate matrices were developed for ocean fisheries mortality of LCN coho (Table A-23). Because the exploitation rate matrices incorporate the same marine survival index, and OCN and LCN coho likely experience the same ocean conditions, managers must be mindful of situations where improved parental spawner abundance of OCN coho allows for ocean exploitation levels that make it impossible to achieve the fishery exploitation rates for LCN.

Table A-22. Harvest management matrix identifying allowable fishery impacts and ranges of resulting recruitment based on parental spawner abundance and marine survival (OCN work group revisions to original PFMC matrix).

Parental Spawner Status *	Marine Survival Index (based on return of jacks per hatchery smolt)			
	Extremely Low (<0.0008)	Low (0.0008-0.0014)	Medium (>0.0014-0.0040)	High (>0.0040)
High: >75% of full seeding	≤8%	≤15%	≤30%	≤45%
Medium: >50% to ≤75% of full seeding	≤8%	≤15%	≤20%	≤38%
Low: >19% to ≤50% of full seeding	≤8%	≤15%	≤15%	≤25%
Very Low: >4 fish/mile to ≤19% of full seeding	≤8%	≤11%	≤11%	≤11%
Critical **: ≤4 fish/mile	0-8%	0-8%	0-8%	0-8%

Sub-aggregate and Basin-specific Spawner Criteria Data							
Sub-aggregate	Miles of Available Spawning Habitat	100% of Full Seeding	Critical		Spawner Status Intervals		
			4 fish/mile	12% of full seeding	19% of full seeding	50% of full seeding	75% of full seeding
Northern	899	21,700	3,596	n/a	4,123	10,850	16,275
North-Central	1,163	55,000	4,652	n/a	10,450	27,500	41,250
South-Central	1,685	50,000	6,740	n/a	9,500	25,000	37,500
Southern	450	5,400	n/a	648	1,026	2,700	4,050
Total	4,197	132,000		15,636	25,099	66,050	99,075

* Parental spawner abundance status for the OCN aggregate assumes the status of the weakest sub-aggregate.

** Critical parental status is defined as ≤4 fish per mi for the Northern, North-Central, and South-Central sub-aggregates; because of high quality spawning habitat in the Rogue River basin, critical status for the Rogue River (Southern sub-aggregate) is defined as 12% of full seeding of high quality habitat.

Table A-23. Harvest management matrix for Lower Columbia Natural (LCN) coho with maximum allowable ocean fishery mortality rates.

Parental Escapement*	Marine Survival Index (based on return of jacks per hatchery smolt)			
	Critical (<0.0008)	Low (<0.0015)	Medium (<0.0040)	High (>0.0040)
High >75% of full seeding	<8%	<15%	<30%	<45%
Medium >50% to ≤75% of full seeding	<8%	<15%	<20%	<38%
Low >20% to ≤50% of full seeding	<8%	<15%	<15%	<25%
Very Low >10% to ≤20% of full seeding	<8%	<11%	<11%	<11%
Critical ≤10% of full seeding	0-8%	0-8%	0-8%	0-8%

* Full Seeding: Clackamas River = 3,800; Sandy River = 1,340.

Fisheries off the Oregon and Washington coasts are developed through the PFMC and NOF processes and are subject to agreements of the PST. The NOF process integrates ocean and river management objectives, constraints, and agreements to formulate a coordinated management plan. For example, coho fisheries in 2002 were structured to address standards for ESA-listed stocks (especially OCN coho) and PST obligations regarding Thompson River coho (BC stock). U.S. fisheries, including those in Puget Sound, were limited to a total exploitation rate under 10% on Thompson River coho. Low expected abundance levels of lower Columbia River hatchery coho reduced coho quotas off the Washington and Oregon coasts compared to 2001.

In establishing ocean salmon fisheries that impact OPI area coho stocks, PFMC is guided by the NMFS 1999 Supplemental Biological Opinion and Incidental Take Statement for the three ESA-listed coho stocks in the OPI area. To protect threatened CCC coho, no directed coho fisheries or retention of coho is allowed in commercial and recreational fisheries off the California coast. Marine fishery impacts on threatened CCC and SONC coho must be no more than 13% based on projected impacts on Rogue/Klamath hatchery coho. Marine and freshwater impacts on OCN coho should not exceed levels in the abundance-based fishery management plan (15% in recent years).

The PFMC management process includes evaluating proposed fishing seasons and quotas by assessing their ability to meet management criteria for key coho stocks present in West Coast fisheries. Table A-24 displays the management criteria and projected results of ocean salmon seasons adopted by the PFMC for 2003.

A recent important management tool in the PFMC fishery management process is the use of selective fisheries for hatchery-marked adipose fin-clipped fish. For planning purposes, the STT estimates the rate of marked fish expected to be caught in particular fisheries to anticipate potential effects on wild fish. The 2003 expected mark rates for selective coho ocean fisheries are presented in Table A-25.

Table A-24. Management criteria and projected key stock escapements (in thousands of fish) for coho salmon in PFMC-adopted ocean salmon fisheries, 2003.

Key Stock/Criteria	Projected Ocean Escapement* or Other Criteria (Council Area Fisheries)	Spawner Objective or Other Standard
<i>Columbia River</i>		
Upper Columbia	52%	50%; minimum % of the run to Bonneville Dam.
Columbia River Hatchery Early	246.4	38.7; minimum ocean escapement to attain hatchery egg-take goal of 19.6 early adult coho, assuming average conversion and no mainstem or tributary harvest
Columbia River Hatchery Late	145.9	19.4; minimum ocean escapement to attain hatchery egg-take goal of 15.2 late adult coho, with average conversion and no lower river mainstem or tributary harvest
<i>Coastal Natural</i>		
Quillayute Fall	21.2	6.3-15.8; MSY adult spawner range (not annual target); annual management objectives may be different and are subject to agreement between WDFW and the treaty tribes
Hoh	10.4	2.0-5.0; MSY adult spawner range (not annual target); annual management objectives may be different and are subject to agreement between WDFW and the treaty tribes
Queets Wild	19.6	5.8-14.5; MSY adult spawner range (not annual target); annual management objectives may be different and are subject to agreement between WDFW and the treaty tribes
Queets Supplemental	1.1	
Grays Harbor	52.3	35.4; MSP level of adult spawners; annual management objectives may be different and are subject to agreement between WDFW and the treaty tribes
Oregon Coastal Natural (threatened)	14.4%	≤15%; marine and freshwater fishery exploitation rate
Northern California (threatened)	9.6%	≤13%; marine fishery exploitation rate for R/K hatchery coho (NMFS ESA consultation standard)
<i>Puget Sound</i>		
Skagit	37% (5.4%) 97.9	≤60%; 2003 total exploitation rate ceiling based on comanager comprehensive coho management plan ^a ; 30.0 MSP level of adult spawners identified in FMP
Stillaguamish	37% (7.8%) 27.7	≤50%; 2003 total exploitation rate ceiling based on comanager comprehensive coho management plan ^b ; 17.0 MSP level of adult spawners identified in FMP
Snohomish	33% (7.8%) 147.6	≤60%; 2003 total exploitation rate ceiling based on comanager comprehensive coho management plan ^b ; 70.0 MSP level of adult spawners identified in FMP
Hood Canal	41% (5.9%) 25.8	≤45%; 2003 total exploitation rate ceiling based on comanager comprehensive coho management plan ^b ; 21.5 MSP level of adult spawners identified in FMP
Strait of Juan de Fuca	14% (5.8%) 18.0	≤40%; 2003 total exploitation rate ceiling based on comanager comprehensive coho management plan ^b ; 12.8 MSP level of adult spawners identified in FMP
<i>Canada</i>		
Interior Fraser (Thompson River)	8.3%	≤10%; total exploitation rate for all U.S. fisheries south of the U.S./Canada border

* Ocean escapement is the number of salmon escaping ocean fisheries and entering fresh water.

** Annual management objectives may be different than FMP goals and are subject to agreement between WDFW and the treaty tribes under U.S. District Court orders. Total exploitation rate includes all fisheries and is calculated as total fishing mortality divided by total fishing mortality plus spawning escapement.

Table A-25. Expected mark rates for Council-adopted ocean salmon fisheries with selective coho retention, 2003.

Area	Fishery	June	July	Aug.	Sep.	2002 Observed
<i>North of Cape Falcon</i>						
Neah Bay (Area 4)	Recreational	39%	57%	45%	52%	39%
	Non-Indian troll	—	47%	47%	52%	NA
La Push (Area 3)	Recreational	64%	54%	64%	18%	28%
	Non-Indian troll	—	55%	50%	71%	NA
Westport (Area 2)	Recreational	75%	74%	72%	74%	56%
	Non-Indian troll	—	60%	70%	50%	NA
Columbia River (Area 1)	Recreational	89%	87%	83%	83%	58%
	Non-Indian troll	—	77%	78%	77%	NA
Buoy 10	Recreational	—	—	81%	81%	74%
<i>South of Cape Falcon</i>						
Cape Falcon to Humbug Mt.	Recreational	—	—	—	—	56%
Tillamook	Recreational	80%	75%	67%	—	—
Newport	Recreational	77%	75%	68%	—	—
Coos Bay	Recreational	74%	71%	58%	—	—

Coho production in the OPI area has exceeded 4.3 million fish (1976) and been as low as 216,400 fish in 1995, (Figure A-64). Production was consistently low throughout the 1990s but since 2000 has increased and is similar to the average production since 1970 (1.5 million coho). The highest ocean escapement to the Columbia River was over 1.5 million fish in 1986; the lowest Columbia River escapement was 75,200 coho in 1995. Columbia River escapements since 2000 have exceeded the 1970-2002 escapement average (407,200 coho). Historically, most of the production was harvested in ocean fisheries; ocean fisheries accounted for almost 90% of the OPI production in some years, while ocean escapement to the Columbia River was less than 10% of OPI production during these years (Figure A-65). In recent years, ocean fisheries account for about 10-15% of the total OPI coho production, while Columbia River escapement has been approximately 70% of the total OPI coho production. The remaining coho include escapement to the Oregon Coast and California OPI areas.

Commercial troll fisheries have been closed to coho retention south of Cape Falcon since 1993. In 2000 and 2001, commercial troll selective fisheries for marked hatchery coho occurred from Cape Falcon, OR to the Queets River, WA. In 2002, commercial troll selective fisheries for marked hatchery coho occurred from Cape Falcon to Leadbetter Point, WA. Limitations on Chinook harvest (such as fishery closures in July and a 4-spread requirement on gear) have also been used to reduce impacts to OCN coho.




Figure A-64. Coho salmon ocean fisheries harvest and ocean escapement of the primary coho stock components within the OPI area, 1970–2002.




Figure A-65. Exploitation rate of ocean fisheries on OPI area coho stocks and percent of total OPI area production accounted for by coho ocean escapement to the Columbia River.

Ocean coho harvest in PFMC-managed waters generally occurred from May–October. California ocean commercial troll fisheries occur from May–October, although most landings are in June and July (Figure A-66). Oregon ocean commercial troll fisheries generally are from June–October, with the largest harvests in July and significant harvest in August (Figure A-66). Washington ocean non-Indian troll fisheries are from May–September and most of the harvest occurs in July and August (Figure A-66). Washington treaty Indian commercial ocean troll fisheries occur throughout the year, with the majority of harvest in July–August, although in recent years, the September harvest has been substantial (Figure A-66).

The ex-vessel value and the price per pound of troll-caught coho in California, Oregon, and Washington ocean fisheries has declined since the 1980s (Figure A-67). Minimal fishing occurred in all areas throughout the 1990s; recent year hatchery selective fisheries have occurred in Oregon and Washington. The total ex-vessel value of these fisheries in recent years has been a fraction of their historical value. Price per pound also has generally been low, except for the Washington 2002 fisheries when the price was comparable to some historical years.

Retention of coho in ocean recreational fisheries has been restricted since 1993. Since 1998, coho-directed recreational fisheries in the OPI area have been selective for adipose fin-clipped hatchery-marked fish. Improving hatchery coho populations in the OPI area in recent years have allowed increasing opportunities for a hatchery-marked coho fishery. Recreational ocean harvest of coho in California is generally greater in the private sector than by charter boats; harvest has been minimal since the 1994 season for either boat type (Figure A-68). In Oregon, recreational ocean harvest of coho is dominated by private boats (Figure A-68). In Washington, coho landings by charter boats historically exceeded private boat landings, but the private boat harvest has been greater in recent years (Figure A-68).

Angler effort in California has remained relatively steady over the past 20 years due to stable hatchery Chinook runs, primarily from the Sacramento River. Beginning in the early 1980s, angler effort was reduced significantly in Washington ocean fisheries in response to constraints on Chinook and coho, and angler effort in Oregon lessened due to constraints on Oregon coastal wild coho. Angler effort in both Washington and Oregon has rebounded recently because of improved Chinook and coho abundance and implementation of selective fisheries (Figure A-69).




Figure A-66. California, Oregon, Washington, and treaty Indian ocean commercial troll landings (in thousands of fish) by month, 1976–2002.




Figure A-67. Total value and price per pound (in 2002 dollars) for ocean troll coho landings in California, Oregon, and Washington, 1971–2002.




Figure A-68. Ocean recreational coho catch 1979–2002.




Figure A-69. California, Oregon, and Washington ocean recreational salmon effort (in thousands of angler trips) by boat type, 1979–2002.

Columbia River Fisheries

Coho are harvested in Columbia River mainstem and Select Area commercial fisheries, as well as in Buoy 10, mainstem Columbia, and tributary sport fisheries. Coho also are harvested in treaty Indian fisheries in the Columbia River and tributaries upstream of Bonneville Dam. The Columbia River Compact manages coho fisheries under the requirements of a *U.S. v. Oregon* Fall Management Agreement for upper Columbia coho and Oregon ESA limitations for Lower Columbia Natural coho. The resulting management requirements are:

- pass a minimum of 50% of upper Columbia coho through ocean and lower Columbia River fisheries to escape over Bonneville Dam, and
- fishery impacts to lower Columbia natural coho not to exceed management matrix levels as adopted by the OFWC (14% in 2002).

Maximum allowable freshwater impacts were developed for OCN coho (Table A-26) to guide Columbia River coho fisheries. These rates were adopted by OFWC and are implemented by ODFW and WDFW through the Columbia River Compact.

Combined total harvest rates, including ocean and Columbia River, also were adopted to ensure that total exploitation rates are consistent with state coho management requirements (Table A-27). Established for Oregon state-listed coho, these fishing rates also provide harvest protection for wild Washington coho.

The Buoy 10 area at the mouth of the Columbia River provides the most popular and productive sport coho fishing for Columbia River stocks. Buoy 10 angler trips exceed 100,000 in years of high coho abundance, and the combined Oregon and Washington economic impact has been as high as 9 million (Table A-28). The coho harvest in the Buoy 10 sport fishery has exceeded 100,000 fish four times since 1986, and exceeded 200,000 fish in 1992 (Figure A-61). Coho salmon are actively feeding when entering the Columbia estuary and fishing can be quite successful during mid-August to mid-September. Sport harvest of coho is less productive in the mainstem Columbia upstream of the estuary area.

Table A-26. Harvest management matrix for Lower Columbia Natural (LCN) coho with maximum allowable freshwater fishery mortality rates.

Parental Escapement*	Marine Survival Index (based on return of jacks per hatchery smolt)			
	Critical (<0.0008)	Low (<0.0015)	Medium (<0.0040)	High (>0.0040)
High: >75% of full seeding	<4%	<7.5%	<15%	<22.5%
Medium: >50% to ≤75% of full seeding	<4%	<7.5%	<11.5%	<19%
Low: >20% to ≤50% of full seeding	<4%	<7.5%	<9%	<12.5%
Very Low: >10% to ≤20% of full seeding	<4%	<6%	<8%	<10%
Critical: ≤10% of full seeding	0-4%	0-4%	0-4%	0-4%

* Full Seeding: Clackamas River = 3,800; Sandy River = 1,340.

Table A-27. Cumulative exploitation rates for LCN coho under the combined management protocols proposed for setting ocean and in-river fishery harvest rates.

Parental Escapement*	Marine Survival Index (based on return of jacks per hatchery smolt)			
	Critical (<0.0008)	Low (<0.0015)	Medium (<0.0040)	High (>0.0040)
High >75% of full seeding	<11.7%	<21.4%	<40.5%	<57.4%
Medium >50% to ≤75% of full seeding	<11.7%	<21.4%	<29.2%	<49.8%
Low >20% to ≤50% of full seeding	<11.7%	<21.4%	<22.7%	<34.4%
Very Low >10% to ≤20% of full seeding	<11.7%	<16.3%	<18.1%	<19.9%
Critical ≤10% of full seeding	0-11.7%	0-11.7%	0-11.7%	0-11.7%

* Full Seeding: Clackamas River = 3,800; Sandy River = 1,340.

Table A-28. Angler trips and economic impact (in 2002 dollars) of the Buoy 10 recreational fishery, 1982–2002.

Year	Angler Trips (000s)	Economic Impact (000s)	
		Oregon	Washington
1982	17.3	n/a	n/a
1983	7.1	n/a	n/a
1984	67.4	n/a	n/a
1985	32.2	n/a	n/a
1986	102.2	n/a	n/a
1987	125	\$2,169	\$3,928
1988	183	\$3,075	\$6,212
1989	156	\$2,346	\$5,148
1990	80	\$1,264	\$2,386
1991	172	\$2,672	\$5,544
1992	115	\$1,762	\$3,638
1993	76	\$1,179	\$2,182
1994	9	\$189	\$230
1995	25	\$491	\$615
1996	18	\$373	\$420
1997	56	\$910	\$1,728
1998	30	\$507	\$860
1999	50	\$907	\$1,276
2000	73	\$1,335	\$2,000
2001	126	\$2,636	\$2,940
2002	84	\$1,803	\$1,849

WDFW statewide rules declare that salmon fisheries are closed unless otherwise specified in Special Rules. Depending on the strength of adult salmon returns, WDFW promulgates regulations allowing spring Chinook, fall Chinook, and coho salmon fisheries in lower Columbia River tributaries. Coho fisheries typically overlap fall-run Chinook fisheries in the Washington tributaries. Salmon-directed fisheries will vary from year to year and from stream to stream depending on the health of salmonid populations and sizes of runs forecast for each particular stream. Fisheries for adipose fin-clipped hatchery coho salmon destined for the Grays, Elochoman, Cowlitz, Toutle, Kalama, Lewis, Washougal, and Little White Salmon rivers occur from August through January in most years. Anglers experience good success rates for coho in the tributary fisheries (Figure A-61). Selective fishery regulations have been in place for all lower Columbia River sport fisheries since 1998.

Fall commercial fisheries before late-September primarily harvest early coho and fall Chinook. Commercial fisheries after early October primarily harvest late hatchery coho stocks and sturgeon; fisheries between these two time periods harvest both early and late coho stocks. Late fall seasons in October primarily target hatchery coho in the lower river below the mouth of the Lewis River.

Commercial fishing in Columbia River off-channel areas (i.e. Select Area fisheries) commenced in 1962 when salmon seasons were adopted for Youngs Bay, OR. Initially, openings were concurrent with the late fall mainstem gill net seasons but seasons have been separate since 1977. Recent declines in mainstem fishing opportunities prompted BPA to fund a research project to expand net-pen programs to select off-channel fishing areas. The result of this effort was the Select Area Fishery Enhancement (SAFE) Project, which has expanded to Tongue Point/South Channel and Blind/Knappa Slough in Oregon and Deep River and Steamboat Slough in Washington. Coho fisheries occur in all five Select Areas; these fisheries primarily target hatchery coho returning to specific release sites. Coho-targeted Select Area fisheries occur from August through October; most harvest occurs in September and October. The 2001 fall Select Area fisheries harvest totaled 33,687 coho salmon.

Coho salmon are the target species for late fall lower Columbia River commercial fisheries. Late fall coho seasons end before November to avoid impacts to late returning wild Clackamas coho, chum, and winter steelhead. Late returning wild Washington coho also benefit from the November season closure. Coho are also incidentally harvested in early fall commercial fisheries targeting fall Chinook. Coho salmon are also harvested in treaty Indian commercial and subsistence fisheries in Zone 6 above Bonneville Dam (Figure A-60). No prohibitions are in place on wild coho retention for the treaty Indian fisheries, but coho harvest in the treaty Indian commercial fishery is usually minor because of constraints to protect wild steelhead.

The PFMC uses a model to estimate catch, mortality, and escapement of early and late Columbia River coho; the model also partitions the fish into lower and upriver coho. Results of the 2002 model run are summarized in Table A-29; coho salmon exploitation rates can be inferred from the model. Note the change in the ratios of marked and unmarked coho in fisheries as marked coho are removed from the population prior to the fish entering the next fishery.

Table A-29. Estimated catch, mortality, and escapement of marked and unmarked Columbia River basin coho salmon, 2002.

Harvest & Interim Abundance	Marked (Hatchery)		Unmarked (Wild)	
	No. of Fish	Exploitation Rate	No. of Fish	Exploitation Rate
Ocean abundance	326,649		49,234	
Alaska & Canada harvest	120	0.04%	24	0.05%
<i>U.S. v. Oregon</i> area ocean abundance	326,529		49,210	
<i>U.S. v. Oregon</i> area catch and mortality (w/o treaty troll)	123,761	38%	5,173	11%
Ocean natural mortality	65,344		9,910	
Columbia River mouth abundance	137,424		34,127	
Buoy 10 catch and mortality	19,074	6%	1,115	2%
Mainstem recreational catch and mortality	977	0.3%	23	0.05%
August commercial catch	74	0.02%	26	0.05%
September commercial catch	7,975	2%	2,025	4%
October commercial catch	8,429	3%	1,571	3%
Tributary escapement	100,895		29,367	
<i>Total exploitation*</i>		49.36%		20.15%

* Does not include treaty Indian ocean troll fisheries or tributary recreational harvest.

A.2.10. Assessments of Current Status and Limiting Factors

Listing Status

Lower Columbia coho ESU was listed as a threatened species under ESA on June 28, 2005 (Fed. Reg., V70, N123, p.37160). In a 1995 status review of coho salmon, NMFS found that if an evolutionarily significant unit of coho salmon (such as Clackamas River late-run coho) still exists in the lower Columbia River, it is not presently in danger of extinction, but is likely to become so (NMFS 1995). However, the Oregon Fish and Wildlife Commission conducted its own status review and concluded that lower Columbia coho produced in Oregon basins, including the Clackamas and Sandy Rivers, are at risk of extinction, and listed them as a state endangered species in 1998. NMFS was subsequently petitioned to list lower Columbia coho salmon and to designate critical habitat. The ESU was listed in 2005 but critical habitat designation is still being developed.

Current Status

Coho historically spawned in all accessible lower Columbia River tributaries but the run now consists of very few wild fish. Until recently, Columbia River coho salmon were managed primarily as a hatchery stock. Present coho populations have been heavily influenced by extensive hatchery releases.

Every one of the 17 Washington lower Columbia River coho populations is estimated to be at a very high risk of extinction. Twenty-one of the 24 populations in the ESU, including Oregon, are at a very high risk (Figure A-70). It is possible that some native coho populations are now extinct, but that this loss is masked by the presence of naturally spawning hatchery fish. No healthy coho populations exist in this ESU. The strongest remaining populations occur in Oregon and include the Clackamas River and Scappoose Creek (both of moderate viability). Current numbers of highly viable populations (0) and

average viability scores (Coast = 0.6, Cascade = 0.6, Gorge = 0.3) fall well below the standards established by the WLC-TRT for viability (2 populations per strata at or above 3.0, strata average viability of at least 2.25).

Abundance and productivity of naturally-spawning coho is very low in all Washington populations (Table A-30). Current wild abundance is uncertain due to large numbers of naturally-spawning hatchery coho but habitat assessments suggest that current natural production capacity is under 2,000 fish. Historical abundance was estimated on the order of several hundred thousand coho. Annual escapement data is lacking for naturally-produced Washington coho but hatchery return data shows high annual variability and a declining trend in productivity (Figure A-46). Since 1970, the population of Columbia hatchery-origin adult coho ocean has ranged from over 3 million in 1971 to a low of 97,000 in 1996. Returns to the Columbia River mouth (after ocean harvest) have declined from over 1.5 million in 1986 to 75,000 in 1995. Recent numbers have rebounded somewhat from very low levels during the early 1990s but the Columbia River coho run continues to be dominated by fish released from hatcheries and naturally-produced fish from hatchery-origin parents.

Reductions in spatial structure have also contributed to the very low viability of many coho populations. While many populations continue to have access to most of their historical range, the area of productive habitat has been significantly reduced by habitat degradation in many systems, particularly in the smaller, rain-driven, lower elevation tributaries. Access has been lost to upstream areas of the Cowlitz and Lewis rivers which historically supported very large, diverse, and productive runs of early and late coho.

Diversity of spring Chinook is estimated to have been greatly reduced due to pervasive hatchery effects and small population bottlenecks within the natural populations. Hatchery-origin fish typically comprise a large fraction of the spawners in natural production areas. Widespread inter-basin (but within ESU) stock transfers have homogenized many populations. While historical population structure likely included significant genetic differences among populations in each watershed, unique natural populations of coho salmon can no longer be genetically distinguished in the lower Columbia River (excluding the Clackamas and Sandy rivers in Oregon).




Figure A-70. Current status of historical demographically-independent lower Columbia River coho salmon populations.

Table A-30. Status and recovery goals for lower Columbia River coho populations (relative to listing baseline).

Population	Contribution	Baseline viability				Obj.	Prod. target	Abundance		
		A&P	S	D	Net			Historical	Baseline	Target
Coast										
Grays/Chinook ^L	Primary	VL	H	VL	VL ²	H	+370%	3,800	<50	2,400
Eloch/Skam ^L	Primary	VL	H	VL	VL ²	H	+170%	6,500	<50	2,400
Mill/Ab/Germ ^L	Contributing	VL	H	L	VL ²	M	>500%	2,800	<50	1,800
Youngs (OR) ^L	Stabilizing	-- ³	-- ³	-- ³	VL	VL	-- ³	-- ³	-- ³	-- ³
Big Creek (OR) ^L	Stabilizing ²	-- ³	-- ³	-- ³	VL	VL	-- ³	-- ³	-- ³	-- ³
Clatskanie (OR) ^L	Primary ¹	-- ³	-- ³	-- ³	L	VH	-- ³	-- ³	-- ³	-- ³
Scappoose (OR) ^L	Primary	-- ³	-- ³	-- ³	M	VH	-- ³	-- ³	-- ³	-- ³
Cascade										
Lower Cowlitz ^L	Primary	VL	M	M	VL ²	H	+100%	18,000	500	3,700
Upper Cowlitz ^{E,L}	Primary ¹	VL	M	L	VL	H ¹	>500%	18,000	<50	2,000
Cispus ^{E,L}	Primary ¹	VL	M	L	VL	H ¹	>500%	8,000	<50	2,000
Tilton ^{E,L}	Stabilizing	VL	M	L	VL	VL ²	0%	5,600	<50	--
Toutle SF ^{E,L}	Primary	VL	H	M	VL ²	H	+180%	27,000	<50	1,900
Toutle NF ^{E,L}	Primary	VL	M	L	VL ²	H	+180%	<50	<50	1,900
Coweeman ^L	Primary	VL	H	M	VL ²	H	+170%	5,000	<50	1,200
Kalama ^L	Contributing	VL	H	L	VL ²	L	>500%	800	<50	500
NF Lewis ^{E,L}	Contributing	VL	L	L	VL ²	L	+50%	40,000	200	500
EF Lewis ^{E,L}	Primary	VL	H	M	VL ²	H	>500%	3,000	<50	2,000
Salmon ^L	Stabilizing	VL	M	VL	VL	VL	0%	na	<50	50
Washougal ^L	Contributing	VL	H	L	VL ²	M+	>500%	3,000	<50	1,500
Clackamas (OR) ^{E,L}	Primary	-- ³	-- ³	-- ³	M	VH	-- ³	-- ³	-- ³	-- ³
Sandy (OR) ^{E,L}	Primary	-- ³	-- ³	-- ³	VL	H	-- ³	-- ³	-- ³	-- ³
Gorge										
L Gorge (WA/OR) ^L	Primary	VL	M	VL	VL ²	H	+400%	na	<50	1,900
U Gorge (WA) ^L	Primary	VL	M	VL	VL ²	H	+400%	na	<50	1,900
U Gorge/Hood (OR) ^E	Contributing ⁴	-- ³	-- ³	-- ³	VL	H ⁴	-- ³	-- ³	-- ³	-- ³

¹ Increase relative to the interim plan.

² Reduction relative to the interim plan.

³ Addressed in the Oregon Management Unit plan.

⁴ Improvement increments are based on abundance and productivity, however, this population will require improvements in spatial structure or diversity to meet recovery objectives.

⁴ OR analysis indicates a low probability of meeting their delisting objective of high viability for this population.

^E Early run (Type S) coho stock.

^L Late run (Type N) coho stock.

(Core and Legacy populations not designated by the Technical Recovery Team for coho.)

Current Limiting Factors

Net Effects of Manageable Factors

Declines in status of lower Columbia River coho salmon result from the combined impacts of human activities involving freshwater habitat, estuary habitat, dam construction and operation, fishing, fish hatcheries, and ecological factors such as predation. Impacts of each factor are compounded across the salmon life cycle to drive most populations to current very low levels. Net effects of quantifiable and potentially manageable impacts translate into an estimated 91-100% (average 96%) reduction in abundance and productivity of lower Columbia River coho populations in Washington (Figure A-71). Thus, current fish numbers represent only 0-9% of the historical production potential in the absence of potentially manageable impacts. Total reductions would be even greater if all human impacts could be effectively quantified. No single factor accounts for the majority of the reduction in fish numbers and the significance of specific factors varies from population to population (Figure A-71, Table A-31).




Figure A-71. Net effect and relative contribution of potentially manageable impact factors on coho salmon in Washington lower Columbia River subbasins. Net effect is the approximate reduction from historical fish numbers as a result of manageable factors included in this analysis. Fisheries

Table A-31. Productivity improvements and impact reduction benchmarks consistent with recovery of Washington lower Columbia River coho populations if reductions are distributed across factors in proportion to their baseline impact.

Population	Productivity Target	Baseline impacts (listing date reference)							Impacts at target					
		Trib	Est	Dams	Harv	Hat	Pred	Δ	Trib	Est	Dams	Harv	Hat	Pred
<u>Coast</u>														
Grays/Chinook	+370%	0.70	0.16	0.00	0.50	0.50	0.14	43%	0.40	0.09	0.00	0.29	0.29	0.08
Eloch/Skam	+170%	0.60	0.16	0.00	0.50	0.50	0.14	30%	0.42	0.11	0.00	0.35	0.35	0.10
Mill/Ab/Germ	>500%	0.50	0.16	0.00	0.50	0.50	0.15	50%	0.25	0.08	0.00	0.25	0.25	0.08
<u>Cascade</u>														
Lower Cowlitz	+100%	0.70	0.16	0.00	0.50	0.50	0.15	17%	0.58	0.13	0.00	0.42	0.45	0.13
Upper Cowlitz	>500%	0.40	0.16	1.00	0.50	0.50	0.15	50%	0.20	0.08	0.50	0.25	0.25	0.08
Cispus	>500%	0.50	0.16	1.00	0.50	0.50	0.15	50%	0.25	0.08	0.50	0.25	0.25	0.08
Tilton	0%	0.95	0.16	1.00	0.50	0.50	0.15	0%	0.95	0.16	1.00	0.50	0.50	0.15
Toutle NF	+180%	0.90	0.16	0.00	0.50	0.50	0.15	12%	0.79	0.14	0.00	0.44	0.44	0.13
Toutle SF	+180%	0.90	0.16	0.00	0.50	0.50	0.15	12%	0.79	0.14	0.00	0.44	0.44	0.13
Coweeman	+170%	0.80	0.16	0.00	0.50	0.20	0.15	23%	0.62	0.12	0.00	0.39	0.15	0.12
Kalama	>500%	0.70	0.16	0.00	0.50	0.50	0.15	20%	0.56	0.12	0.00	0.40	0.40	0.12
NF Lewis	+50%	0.40	0.15	0.85	0.50	0.24	0.16	6%	0.38	0.14	0.80	0.47	0.22	0.15
EF Lewis	>500%	0.80	0.15	0.00	0.50	0.21	0.16	50%	0.40	0.08	0.00	0.25	0.11	0.08
Salmon	0%	0.90	0.15	0.00	0.50	0.50	0.16	0%	0.90	0.15	0.00	0.50	0.50	0.16
Washougal	>500%	0.80	0.15	0.00	0.50	0.50	0.16	50%	0.40	0.08	0.00	0.25	0.25	0.08
<u>Gorge</u>														
L Gorge	+400%	0.50	0.15	0.00	0.50	0.50	0.16	59%	0.20	0.06	0.00	0.20	0.20	0.07
U Gorge	+400%	0.50	0.14	0.10	0.50	0.75	0.19	39%	0.31	0.09	0.06	0.31	0.46	0.12

Fisheries

Fishery impact rates on wild lower Columbia River coho averaged 53% at listing and have been reduced to 22% at present. The primary fisheries targeting Columbia River hatchery coho salmon occur in West Coast ocean and Columbia River mainstem fisheries (Figure A-72). Hatchery-selective harvest regulations or time and area strategies have been widely implemented to limit impacts to wild coho. The exploitation rate of coho prior to the 1990s fluctuated from approximately 60 to 90%. Exploitation of wild and hatchery coho decreased significantly during the 1990s. The exploitation rate of wild coho has continued to decrease to current levels, while the exploitation of hatchery coho has remained similar to the 1990s rate.




Figure A-72. Approximate coho salmon fishery exploitation rates over time and allocation of current exploitation rates among fisheries.

Hatcheries

Hatchery influence continues to be significant for most Washington lower Columbia coho populations (Table A-32). Most coho hatchery programs are intended to mitigate the loss of natural coho production by providing fish for harvest opportunity. Hatchery releases of coho salmon smolts range from 0 to 3.2 million in subbasins where wild coho populations occurred historically. The average adult hatchery fraction for all Washington lower Columbia coho populations was 84%. Average hatchery fraction varied slightly by strata: 90%, 82%, and 85% for the Coast, Cascade, and Gorge strata, respectively. Reintroduction attempts in the upper Cowlitz, Cispus, and Tilton basins have relied almost entirely on hatchery stock. Current coho salmon hatchery broodstock are primarily derived from local populations and moderately affected by hatchery practices (category 2) or derived from other populations within the same ESU (category 3).

The indexed potential for negative impacts of hatchery spawners on wild population fitness was estimated to range from 11-50%. However, the high incidence of hatchery spawners suggests that the fitness of natural and hatchery fish is now probably quite similar and natural populations could collapse without continued hatchery subsidy under current habitat conditions. In general, the highest potential impacts occur in basins where hatcheries have used broodstock from outside of the local basin; this practice has recently occurred in hatchery coho programs in the Coast and Gorge strata. The lowest potential impacts appear to occur in basins that do not maintain active coho salmon hatchery programs and the hatchery programs in adjacent basins utilized local populations for broodstock (e.g. Coweeman, EF Lewis, Salmon).

The Columbia River Hatchery Scientific Review Group (HSRG) recently completed a review of hatchery programs affecting salmon and steelhead populations in the Lower Columbia River. The objective of the review was to change the focus of the Columbia River hatchery system from a harvest supplementation/hydropower mitigation focus to an ecosystem-based approach founded on the idea that harvest goals are sustainable only if they are compatible with conservation goals. Individual population reviews and recommendations can be found at http://www.hatcheryreform.us/mfs/reports/appendix/welcome_show.action.

Table A-32. Presumed reductions in wild population fitness as a result of natural hatchery spawners and survival as a result of interactions with other hatchery species for Washington lower Columbia River coho populations.

Population	Annual releases ^b	Hatchery fraction	Fitness category	Assumed fitness	Fitness impact	Interacting releases ^m	Interspecies impact
Coast Fall							
Chinook/ Grays	602,500 ^c	0.95	3	0.5	0.48	190,000	0.00
Eloch/Skam	930,000 ^d	0.99	3	0.5	0.50	1,050,000	0.01
Mill/Aber/Germ	0 ^e	0.88	3	0.5	0.44	0	0
Cascade							
Lower Cowlitz	3,200,000 ^f	0.85	2	0.7	0.26	5,319,500	0.07
Upper Cowlitz	0 ^h	0.96	2	0.7	0.29	0	0
Coweeman	0 ^e	0.38	2	0.7	0.11	20,000	0
SF Toutle	0 ^e	0.87	2	0.7	0.26	25,000	0
NF Toutle	800,000 ^g	0.86	2	0.7	0.26	825,000	0.01
Cispus	0 ^h	0.96	2	0.7	0.29	0	0
Tilton	0 ^h	0.96	2	0.7	0.29	0	0
Kalama	700,000 ⁱ	0.98	2	0.7	0.29	1,380,000	0.02
NF Lewis	1,695,000 ^j	0.69	2	0.7	0.21	3,070,000	0.04
EF Lewis	0 ^e	0.78	2	0.7	0.23	115,000	0.00
Salmon	0 ^e	0.67	2	0.7	0.20	20,000	0
Washougal	500,000 ^k	0.91	3	0.5	0.46	620,000	0.01
Gorge							
L Gorge	0 ^e	0.91	3	0.5	0.46	0	0
U. Gorge (LWS)	1,000,000 ^l	0.86	3	0.5	0.43	1,420,000	0.02
White Salmon	0 ^e	0.79	3	0.5	0.40	0	0

^a The TRT has not assigned genetic legacy designations to lower Columbia River coho populations.

^b Annual release goals.

^c Comprised of early coho (type S) released in the Grays, Deep, and Chinook Rivers from the Grays River and Sea Resources Hatcheries.

^d Elochoman Hatchery goals include 418,000 early coho (type S) and 512,000 late coho (type N).

^e Hatchery coho salmon are no longer released in the basin; hatchery fish in these basins appear to be strays from other programs.

^f The Lower Cowlitz coho hatchery program is composed of late coho (type N). One goal of the late stock coho salmon hatchery program is to provide restocking of the upper Cowlitz basin. Reintroduction efforts have been challenged in passing juvenile production through the system.

^g Comprised of early coho (type S) released in the NF Toutle and Green Rivers from the NF Toutle Hatchery.

^h Hatchery coho (predominately late coho type N) fry and adults have been released since 1997 and 1998, respectively, into the upper Cowlitz and Cispus Rivers. Outmigrating juvenile coho are collected and transported around the Cowlitz Falls Dam; collection efficiencies have ranged from 17-45%. Recent efforts have also released adults into the Tilton River basin; any juveniles produced in the Tilton need to be collected at Mayfield Dam.

ⁱ The Fallert Creek Hatchery goal is 350,000 early coho (type S); the Kalama Falls Hatchery goal is 350,000 late coho (type N).

^j Lewis River Hatchery goals include 880,000 early coho (type S) and 815,000 late coho (type N); fish are released in the lower Lewis River mainstem. Various possible salmonid reintroduction scenarios are currently being evaluated during the re-licensing process for the hydroelectric facilities on the Lewis River; the existing hatchery programs could become an integral part of any successful reintroduction program.

^k The Washougal River Hatchery releases late coho salmon (type N); broodstock is normally derived from Washougal or Lewis River hatchery returns.

^l The Little White Salmon hatchery goal is composed of early coho (type S).

^m Includes steelhead, coho, and spring Chinook.

Stream Habitat

EDT analyses suggest that stream degradation has substantially reduced the habitat potential for coho in all Washington lower Columbia River subbasins where analyses have been completed (Figure A-73).

Declines in habitat quality and quantity for coho salmon have reduced current productivity and equilibrium population sizes to 10-60% of the historical template. Substantial stream habitat improvements would be necessary to reach optimum conditions (i.e. PFC) for coho salmon in any subbasin. Restoration of optimum habitat quality would be expected to increase habitat capacity by 1,000 to 23,000 adult coho per subbasin, based on preliminary planning ranges.

Coho salmon rely on the middle mainstem to upper stream reaches where a lack of habitat diversity, sedimentation, and flow consistently limit habitat suitability. More detailed descriptions of stream habitat conditions and effects on fish in each subbasin may be found in Volume II of the Technical Foundation.




Figure A-73. Current, optimal, and historical subbasin productivity and capacity inferred for coho salmon from stream reach habitat conditions using EDT.

Dams

At present, there are no EDT assessments that quantify the amount of historical coho habitat that has been blocked or inundated by dam construction within specific basins. Steelhead results give some idea of the scale of effect although coho utilize many more downstream tributary areas than do steelhead. Similarly, coho dam passage rate data are sparse. If similar to steelhead, passage mortality at Bonneville Dam would be assumed to average 10% for juveniles and an additional 5% for adults based on a synthesis of the available literature. Coho salmon generally spawn and rear in headwater and upper mainstem reaches of subbasins and are less subject to hydropower effects on downstream habitats than are chum and fall Chinook.

Mainstem and Estuary Habitat

Mainstem and estuary habitat impacts were estimated to account for approximately a 10-20% reduction in productivity of coho salmon, if similar to steelhead. Coho salmon migrate through mainstem and estuary areas soon after emigration from tributary streams. Residence time in estuary and mainstem habitats is usually relatively brief, but smoltification and transition from fresh to salt water is a critical life stage.

Predation

Potentially manageable predation mortality was assumed to average 20% to 25% depending on travel distance from the subbasin to the ocean. Pikeminnow and tern management is projected to reduce salmonid predation by approximately 50%. Tern predation was almost entirely an artifact of recently established colonies on dredge spoil islands in the estuary but the current rate (9%) is less than half that observed prior to downstream translocation of part of the Rice Island tern colony (20%). Pikeminnow predation was greatest for populations that originate in Bonneville Reservoir tributaries (5%), pass the pikeminnow gauntlet in Bonneville Dam forebay and tailrace, and travel the entire 145-mile length from Bonneville to the Estuary. Predation rates by seals and sea lions on adult coho salmon added an assumed 3% mortality.

Summary Assessment

1. Human activities including fishing, hatchery operation, alteration of stream, river, and estuary habitats, hydropower development and operation, and potentially manageable predation have collectively reduced productivity of coho salmon populations to 0-8% of historical levels. Recovery efforts will require significant improvements in multiple areas because no single factor accounts for the majority of the reduction in fish numbers.
2. Implementation of selective fishery regulations in U.S. ocean and Columbia River fisheries has reduced impacts on wild coho salmon by over half. Additional reductions would require widespread changes in U.S. ocean and Columbia River fisheries. Because Lower Columbia wild coho salmon comprise only a small portion of the catch in many fisheries, additional constraints for their protection will forgo harvest of larger numbers from healthy wild and especially hatchery populations. Intensive fishery management processes provide significant opportunities for limiting fishing risks by tailoring annual harvests to fish availability.
3. Reduced productivity of wild populations as a result of interbreeding with potentially less-fit hatchery fish is among the most significant of hatchery concerns for wild stock recovery although these negative effects are at least partially offset by the demographic benefits of additional spawners. Potential negative impacts increase with the proportion of hatchery spawners and the genetic and phenotypic disparity between wild and hatchery fish. Potential fitness impacts among

Washington lower Columbia coho salmon populations range from 11 to 50%. Potential impacts are greatest in the Coast and Gorge strata populations where out-of-basin stocks continue to be used for broodstock. Inter-specific hatchery predation impacts on juvenile fall Chinook range from 0% in basins without significant releases of coho, steelhead or spring Chinook to a high of 7% in the Cowlitz basins where large hatchery programs are underway.

4. The current conditions of stream habitats significantly limit coho salmon in all Washington lower Columbia River subbasins where EDT analyses have been completed. Substantial stream habitat improvements would be necessary to reach optimum conditions (i.e. PFC) in any subbasin. The significance of stream habitat suggests that recovery may not be feasible without substantial improvements in tributary habitat quantity and quality.
5. Estuary and mainstem habitats are important to coho salmon life history with assumed habitat impacts of 10-20%.
6. Hydropower development in the Cowlitz and Lewis have blocked 50-95% of the historical coho salmon habitat, based on data for steelhead. Mainstem dam passage affects upper Gorge populations although passage success for coho salmon may be as high as steelhead, which tends to be greater than other salmon species.



A.3. Chum Salmon

Chum salmon (*Oncorhynchus keta*) have the widest natural geographic and spawning distribution of any Pacific salmonid, primarily because their range extends farther along the shores of the Arctic Ocean than other salmonids (Groot and Margolis 1991). They have been documented to spawn from Korea and the Japanese island of Honshu, east around the rim of the North Pacific Ocean, to Monterey Bay in southern California. The species' range in the Arctic Ocean extends from the Laptev Sea in the Russian Federation to the Mackenzie River in Canada (Bakkala 1970, Fredin et al. 1977). Chum salmon historically may have been the most abundant of all salmonids—Neave (1961) estimated that prior to the 1940s, chum salmon contributed almost 50% of the total biomass of all salmonids in the Pacific Ocean. Chum salmon also grow to be among the largest of Pacific salmon, second only to Chinook salmon in adult size, with individuals reported up to 42.9 in (108.9 cm) in length and 45.9 lbs (20.8 kg) in weight (*Pacific Fisherman* 1928). Average size for the species is around 7.9 to 15 lbs (3.6 to 6.8 kg) (Salo 1991).

The species is best known for its canine-like teeth and the striking body color of spawning males; a calico pattern, with the anterior two-thirds of the flank marked by bold, jagged, reddish lines and the posterior third by a jagged black line. Females are less flamboyantly colored and lack the extreme dentition of the males. The two most widely used common names, 'chum' and 'dog' salmon, reflect these traits. Chum salmon is the common name accepted by the American Fisheries Society, most likely derived from a word in the language of the Chinook peoples of the Columbia River area, *cam* (also translated as *sum* or *tzum*), which means calico.

In the Columbia River basin, chum salmon once migrated more than 310 miles (500 km) to spawn in the Walla Walla River (Nehlsen et al. 1991) and were productive in many lower Columbia River tributaries. Runs of nearly 1.4 million fish are believed to have returned annually to the Columbia River. The total minimum 2002 chum return to the Columbia River was estimated to be 19,914 fish, based on Washington tributary and lower Columbia mainstem spawning surveys (19,403), commercial fishery incidental catch (14), hatchery escapement (309), and the Bonneville Dam count (188). Production is generally limited to areas downstream of Bonneville Dam. All naturally produced chum populations in *the Columbia River and its tributaries in Oregon and Washington were federally listed as threatened in August 1999.

Intensive monitoring of chum spawning escapement is conducted in three Washington tributaries in the lower Columbia basin—Grays River, Hardy Creek, and Hamilton Creek—and in the mainstem Columbia River near Ives Island. The latter three populations are located immediately downstream of Bonneville Dam. Chum salmon populations exist in other river systems of the lower Columbia, but have not been consistently monitored and abundances are assumed to be extremely low.




Figure A-74. Life cycle for Columbia River Chum salmon.

A.3.1. Life History and Requirements

The freshwater adult life history cycle of lower Columbia chum salmon populations follows the timing of seasonal changes in river flow and water temperatures in lower Columbia River tributaries (Figure A-74). In general, the region has a mild climate with warm, relatively dry summers and cool, wet winters. The river environments these fish enter are characterized by relatively low elevations (1,640-3,280 ft [500-1,000 m]), with moderate amounts of precipitation (80-95 in/year [200-240 cm/year]). These rivers display relatively low flows during late summer and early fall, increased river flows and decreased water temperatures beginning in early October, and a single flow peak in December or January. Upstream migration of chum to tributary spawning areas often coincides with changes in streamflow and water temperature. In the lower Columbia River, streamflows typically begin to rise in October with the onset of fall rains. Water temperatures also drop at this time, creating conditions that favor salmon migration and spawning activity.

Upstream Migration Timing

Chum salmon returning to the Columbia River are considered a fall run. Adult fall run chum salmon return to the Columbia River from mid-October through November, but apparently do not reach the Grays River until late October-early December. Spawning occurs in the Grays River from early November to late December. Fish returning to Hamilton and Hardy creeks begin to appear in the tributaries in early November and their spawn timing is more protracted (mid-November to mid-January). Chum salmon have been reported in October in the Washougal, Lewis, Kalama, and Cowlitz rivers in Washington and in the Sandy River in Oregon (Salo 1991).

Chum seldom show persistence in surmounting river blockages and falls, which may be why they usually spawn in lower river reaches. However, in some river systems, such as Washington's Skagit River, chum salmon routinely migrate distances of at least 105 miles (170 km). They swim even greater distances in at least two other rivers. In Alaska's Yukon River and the Amur River in the Russian Federation, chum salmon migrate more than 1,550 miles (2,500 km) inland. Both of these rivers have low gradients and are without extensive falls or other blockages to migration. Chum salmon that historically traveled up the Columbia River to spawn in the Umatilla and Walla Walla rivers, however, would have had to pass Celilo Falls and a web of rapids and cascades. The falls would have presented a considerable obstacle and probably were passable by chum salmon only at high water flows.

Spawning

Chum salmon spawn primarily in the lower reaches of rivers, digging their redds in the mainstem, tributaries or in side channels of rivers from just above tidal influence to nearly 60 miles (100 km) from the sea. They spawn in shallower, slower-running streams and side channels more frequently than do other salmonids. However, literature on selection of spawning sites and redd characteristics for chum salmon (reviewed in Bakkala 1970, Smirnov 1975, Salo 1991), indicates that under specific circumstances chum salmon spawn in a variety of locations.

Water velocity in spawning areas varies widely for chum salmon. In Washington, Johnson et al. (1971) measured water velocities near 1,000 chum salmon redds and found that velocities where fish spawned varied from 0.0 to 5.5 ft/sec (0.0 to 167.6 cm/sec), and that over 80% of the fish spawned in velocities between 0.7 and 2.7 ft/sec (21.3 and 83.8 cm/sec). This range is similar to that found in other species of salmon. For example, velocities of streams where Chinook salmon spawn are reported to range from 0.3 to 4.9 ft/sec (10 to 150 cm/sec). Johnson et al. (1971) also attempted to correlate abundance indices of chum salmon in Washington with environmental variables such as stream discharge, velocity, and surface water temperatures, but found no relationship between run size and these variables. He concluded that he was unable to measure or to isolate the critical areas in which environmental factors influence run size.

Chum salmon in other parts of the world also choose spawning grounds with a variety of water velocities; for example, fall chum salmon spawned in pools where the velocity was reported to be quite insignificant (Soin 1954, Smirnov 1975). Working on Japan's Hokkaido Island, Sano and Nagasawa (1958) also found that fall chum salmon selected spawning areas with lower water velocities (0.3-0.7 ft/sec [10-20 cm/sec]) than did summer chum salmon in the Amur River area. These differences in the physical characteristics of spawning areas may act to isolate populations or runs in the same river (Salo 1991).

In some locations, subgravel flow (upwelled groundwater) may be important in the choice of redd sites by chum salmon. A summary of available information on Far Eastern chum salmon reported that throughout the Russian Federation and on Hokkaido Island, fall chum salmon *"utilize mostly spring areas of upper tributaries, [as] damage by freezing and other severe winter conditions is relatively minor in most years."* (Sano 1966). However, Sano also notes, based on studies by Smirnov in the 1940s, *"summer chum salmon spawn earlier in the season, and they do not particularly choose spring areas."*

Many Columbia River chum have been found to select spawning sites in areas of upwelling groundwater. New spawning grounds for chum were recently discovered along the Washington shoreline near the I-205 Glen Jackson Bridge where groundwater upwelling occurs. A significant proportion of chum returning to Hamilton Creek spawn in a spring-fed channel, and portions of the Grays River and Hardy Creek populations spawn in the area of springs. Hundreds of chum salmon once returned to spawn within spring-fed areas along Duncan Creek; efforts have been completed to restore passage to these productive areas and protect the springs that feed them. Adult and juvenile chum

salmon from the Ives Island population are being released into newly rehabilitated habitat in Duncan Creek.

Incubation and Emergence

One of the earliest detectable differences between chum salmon populations in different areas is the time it takes for eggs to incubate, hatch, and emerge as alevins from the gravel. Differences between populations are caused by physical factors such as stream flow, water temperature, dissolved oxygen, and gravel composition, and by such biotic factors as genetics, spawning time, and spawning density, all of which can affect survival (reviewed in Bakkala 1970, Salo 1991).

Water temperature is believed to have the most influence on the rate of embryonic development in chum salmon (reviewed in Bakkala 1970, Koski 1975, Salo 1991). The amount of heat, measured in TUs, required by fertilized chum salmon eggs to develop and hatch is about 400-600 TUs, and the heat required to complete yolk absorption is about 700-1,000 TUs. Lower water temperatures can prolong the time required from fertilization to hatching by 1.5-4.5 months. For example, fertilized eggs hatch in about 100-150 days (400-600 TUs) at 39°F (4°C), but hatch in only 26-40 days at 59°F (15°C). Each salmonid has an optimal temperature range that maximizes egg to fry survival. Schroder et al. (1974) reported significantly higher mortality of chum salmon eggs, alevins, and fry when early incubation temperatures were below 34.7°F (1.5°C). Upper thermal limits for chum salmon incubation have not been reported.

The time to hatching also varies among populations and among individuals within a population (Salo 1991). Koski (1975) found differences in the time to hatching between early and late-returning chum salmon at Big Beef Creek, a tributary to Hood Canal. For 2 years (1968-70), early-returning (peak September) and late-returning (peak late November or December) fish spawned and their offspring were reared in spawning channels in the creek. Fry emerged from February to June, but the timing of fry emergence differed between early- and late-returning fish by an average of 35 days each year. Early-run fish took longer to hatch, and this difference between the two runs was consistent from year to year. However, the longer hatching time of early-returning spawners led to fry with lower average weight and less lipid content than fry of late-returning spawners. Lower weight and fewer food reserves in early-return fry may decrease their chances of survival during early life history. The difference in incubation times for eggs from these early- and late-returning fish suggested a genetic difference between the two runs, and Koski (1975) concluded that natural selection apparently acted on hatching times: fry tended to emerge when they had their best chances of surviving in streams and estuaries.

Changes in hatching (incubation) times due to adaptation to cold water also have been found for chum salmon in the Susitna River, Alaska (Wangaard and Burger 1983) and in the Amur River (Disler 1954 cited in Bakkala 1970). At low incubation temperatures, these populations demonstrated faster embryonic development than embryos in other populations at the same temperature. In Canada, however, Beacham and Murray (1986) failed to find differences in hatching times among eggs from adults with early, middle, and late spawning times that had been incubated at constant temperatures of 39, 46, and 54°F (4, 8, and 12°C). Nevertheless, the time of emergence in that study depended on the timing of spawning: earlier-spawning fish laid larger eggs that took longer to develop than did smaller eggs from later-spawning fish.

Factors such as dissolved oxygen, gravel size, salinity, nutritional condition, and even the behavior of alevins in the gravel can influence the time to hatching and emergence from the gravel. For example, Fast and Stober (1984) found that developing chum salmon embryos in small coastal streams required less oxygen than had been reported for either coho salmon or steelhead, but it is unknown to what extent chum salmon in different areas vary in their oxygen requirements. The relative importance of various factors influencing early development in different populations has not been evaluated.

Despite a large amount of variability in incubation environments, even over short distances, chum salmon display a variety of developmental responses that result in similar emergence and outmigration times among fry within a specific area. Variability in some of these responses appears to reflect differences among individual fish, but it also reflects differences among populations in adult run and spawning times, egg size, and temperature-development requirements.

Chum do not have a clearly defined smolt stage, but are nonetheless capable of adapting to seawater soon after emerging from gravel. Chum salmon usually retain parr marks when they first enter seawater. In Japan, chum salmon fry weighing less than 0.06 oz (2 g) maintained normal levels of plasma sodium (Na⁺) when they moved from fresh water into sea water (Iwata 1982). This ability, however, declines slightly with continued residence in fresh water. The capability of chum salmon fry for early osmoregulation in seawater may be important for adults homing back to natal streams. For example, hatchery chum salmon were 10 times less likely to stray within a river system if they were released into the river as fingerlings rather than as smolts (McHenry 1981 cited in Lister et al. 1981).

Freshwater Rearing

Chum salmon do not typically have substantial freshwater rearing time. Most chum juveniles begin seaward migration with minimal time spent in natal streams.

Juvenile Migration

Less is known about chum salmon downstream migrations than juveniles of other salmonids (Salo and Bayliff 1958, Beall 1972, Koski 1975, Seiler et al. 1981, and reviewed in Salo 1991) because chum salmon outmigrants:

1. are smaller than outmigrants of other salmonids,
2. migrate at night,
3. usually have shorter distances to migrate to reach salt water than do other species, and
4. do not school as tightly as some other salmonids (e.g., pink and sockeye fry).

Nonetheless, several key facets of fry outmigration are known. Downstream migration may take only a few hours or days in rivers where spawning sites are close to the mouth of the river, or it may take several months, as in the Yukon and Amur rivers, where spawning sites are located hundreds of kilometers upriver. The timing of outmigration is usually associated with increasing day length, warming of estuarine waters, and high densities of plankton (Walters et al. 1978). Juvenile chum salmon at southern localities, such as those in Washington and southern British Columbia, migrate downstream earlier (late January through May) than do fry in northern British Columbia and southeastern Alaska (April to June).

Several factors influence the timing of downstream migration, resulting in considerable variability in migration timing throughout the species range. These factors include time of adult spawning, stream temperatures during egg incubation and after hatching, fry size and nutritional condition, population density, food availability, stream discharge volume and turbidity, physiological changes in the fry, tidal cycles, and day length (Simenstad et al. 1982, Salo 1991). In the Russian Federation, Soldatov (1912 cited in Smirnov 1975) found that chum salmon outmigrations did not always immediately follow emergence; juveniles in many rivers remained up to 4 months in the river and grew to a considerable size before outmigration (Kostarev 1970 as cited in Salo 1991). In Washington, chum may reside in fresh water for as long as a month (Salo and Noble 1953, Bostick 1955, Beall 1972). Juveniles have been found to reside in fresh water for more than a month in the mainstems of the Skagit (Dames and Moore 1976) and Nooksack (Tyler 1964) rivers.

Because chum fry generally emigrate shortly after emergence, predation mortality during downstream emigration can be significant. Coho juveniles, resident trout, and cottids have been implicated as the primary predators, however, the species composition in each system plays a significant role. The estimated mean freshwater mortality as a result of predation ranges from 22% to 58%. In general, predation on smaller chum fry is thought to be high and predation decreases as chum fry size increases (Beall 1972, Hiyama et al. 1972). To compensate for this predation mortality, chum fry form schools (Pitcher 1986) and synchronize their movements (Miller and Brannon 1982). Historical information concerning the timing of chum salmon emigration in the lower Columbia River is limited. One existing report, however, describes emerging fry outmigrating past the Mayfield Dam site on the Cowlitz River in March and May 1955 and 1956. Thompson and Rothfus (1969) reported the passage at the Mayfield Dam site of approximately 137,250 chum outmigrants past the site between March–May 1955, and about 8,200 fry during the same period in 1956. A wild Chinook capture and tag project conducted by Washington Department of Fish and Wildlife (WDFW) in the North Lewis River during 1977–79 showed incidental capture of chum fry peaking in April and not present in the catch after mid-May. In recent years, seining projects conducted by WDFW in the Grays River and at Ives Island by WDFW and the Oregon Department of Fish and Wildlife (ODFW) indicate outmigration occurs from March through May and peaks from mid-April to early May. Similar activities are being conducted in Deep River to assure release of hatchery fish from net pens is timed to minimize predation.

Estuary Rearing and Growth

The period of estuarine residence appears to be the most critical phase in the life history of chum salmon and may play a major role in determining the size of the subsequent adult run back to fresh water (Mazer and Shepard 1962, Bakkala 1970, Mathews and Senn 1975, Fraser et al. 1978, Peterman 1978, Sakuramoto and Yamada 1980, Martin et al. 1986, Healey 1982, Bax 1983a, Salo 1991). Chum salmon juveniles, like other anadromous salmonids, use estuaries to feed before beginning long-distance oceanic migrations. However, chum and ocean-type Chinook salmon usually have longer residence times in estuaries than do other anadromous salmonids (Dorcey et al. 1978, Healey 1982). Bax (1983b) determined that the extent of juvenile mortality within 4 days of a hatchery release into the Hood Canal estuary was 31-46%. The most important determinant of estuarine survival may be the timing of entry into salt water because plankton abundance in estuaries is highly seasonal (Gunsolus 1978, Helle 1979, Gallagher 1979, Simenstad and Salo 1982).

Because chum salmon spend more time in the estuary, they are more susceptible to changes in the productivity of that environment than stream-type salmonids. Estuaries may be 'overgrazed' when large numbers of ocean-type juveniles enter the estuary *en masse* (Reimers 1973, Healey 1991). The loss of coastal wetlands to urban or agricultural development may more directly affect ocean-type populations than stream-type populations. For example, Thomas (1983) and Johnson et al. (2003b) have documented substantial loss of marsh and swamp habitat throughout the estuary and the lower Columbia River mainstem; further, many researchers (Healey 1982, Levy and Northcote 1982, Myers and Horton 1982, Simenstad et al. 1982, Levings et al. 1986, Bottom et al. 1984) have documented that small juvenile salmonids usually occupy shallow, protected habitats such as salt marshes, tidal creeks, and intertidal flats.

Chum salmon juveniles of early-returning adults tend to enter estuaries before juveniles of late-returning fish (Koski 1975). Because the juvenile emigration timing of lower Columbia River chum salmon from the natal streams is generally from March to May with peak migration in April, chum salmon likely begin arriving in the Columbia River estuary in April. Juvenile chum salmon were a minor portion of the catch during Columbia River estuary sampling efforts of Bottom et al. (1984); chum, sockeye, and cutthroat collectively represented 1% of the total juvenile salmonid catch. Chum salmon juveniles were captured in the estuary during April and May during both years of the study; chum

salmon were present in the estuary from February through June (Bottom et al. 1984). Juvenile chum salmon were primarily distributed within the freshwater or estuarine regions of the estuary, although there was one occurrence in the marine region (Bottom et al. 1984).

Residence times are known for only a few estuaries, even though residence timing has been studied since the 1940s (reviewed in Congleton 1979, Healey 1982, Simenstad et al. 1982, Bax 1983a). Observed residence times range from 4 to 32 days, with a period of about 24 days being the most common.

Migration patterns of juvenile chum salmon have been studied intensively in areas such as Hood Canal by following marked juveniles from hatchery populations of fall-run chum salmon and by monitoring outmigration (Bax 1982, 1983a, b; Bax et al. 1979, 1980; Bax and Whitmus 1981; Schreiner 1977; Whitmus and Olsen 1979; Whitmus 1985; Salo et al. 1980). Some fry remain near the mouth of their natal river when they enter an estuary, but most disperse within a few hours into tidal creeks and sloughs up to several kilometers from the mouth of their natal river. Movements of chum salmon fry in Hood Canal generally appear to follow a pattern that depends on the time of release from hatcheries, however, release time is not the only factor influencing migratory patterns (Bax 1982, 1983a). Chum salmon fry released into Hood Canal in early February and March spread out over a large area, but fish released in April and early May tended to remain inshore initially, moving offshore in summer. These movements were apparently associated with prey availability. Fish initially fed inshore on epibenthic organisms, then offshore on plankton later in the season.

In the Nanaimo and Fraser River estuaries, juveniles spend up to 3 weeks feeding in the inner estuary, with little local movement (Healey 1979, Levy et al. 1979). Chum salmon juveniles in the Nanaimo, Yaquina, Cowichan, and Courtenay estuaries are most abundant in nearshore areas during April and May, but are most abundant in the outer estuary during May and June (Myers 1980, Healey 1982). Chum salmon fry show daily tidal migrations in the Fraser and Nanaimo rivers, which have large deltas and marshlands (Healey 1982). However, fry in Hood Canal have not been observed to display daily tidal migrations (Bax 1983a), most likely because rivers entering Hood Canal do not have extensive delta or tidal marsh systems (with the exceptions of the Quilcene and Skokomish rivers).

Chum salmon spend more of their life history in marine waters than other Pacific salmonids. Chum salmon, like pink salmon, usually spawn in coastal areas, and juveniles outmigrate to seawater shortly after emerging from the gravel (Salo 1991). This ocean-type migratory behavior contrasts with the stream-type behavior of some other species in the genus *Oncorhynchus* (e.g., coastal cutthroat trout, steelhead, coho salmon, and most types of Chinook and sockeye salmon), which usually migrate to sea at a larger size, after months or years of freshwater rearing. Again unlike stream-type salmonids, survival and growth in juvenile chum salmon depend less on freshwater conditions than on favorable estuarine conditions, except for those chum salmon that undergo lengthy migrations such as in the Yukon and Amur rivers.

Recent sampling of juvenile salmonids in the Columbia River plume has started to illustrate patterns of habitat use by salmonids in the plume and nearshore ocean habitats (Fresh et al. 2003), although limited years of data are currently available. For example, preliminary evidence suggests that some juvenile salmonids (chum, steelhead, and yearling coho) may preferentially utilize the plume front compared to other areas in the plume or adjacent ocean habitats (Fresh et al. 2003). Although reasons for the apparent preference to the plume front are not clear, this area may be a more productive habitat than elsewhere in the plume and adjacent ocean.

Ocean Migrations

Little is known about the seaward migration of juvenile chum salmon from the Columbia River. Generally, however, migration of chum salmon juveniles out of estuaries appears to be closely correlated with prey availability. Chum salmon move offshore as they reach a size that allows them to feed on the larger neritic plankton, and this movement normally occurs as inshore prey resources decline (Salo 1991). This transition has taken place at 1.75 in (45 mm) fork length (FL) in Puget Sound and Hood Canal, Washington, but at 2.33 in (60 mm) FL in Prince William Sound, Alaska (Cooney et al. 1978).

Studies have shown that chum salmon in Puget Sound, Washington, and southern British Columbia generally entered the ocean earlier than did more northern and western populations (Hartt 1980, Hartt and Dell 1986). Hartt (1980) and Hartt and Dell (1986) summarized available data on the distribution, migration, and growth of chum salmon in their first year at sea and found that chum, pink, and sockeye salmon juveniles tended to group together and remained nearer shore (within 22 miles [36 km]) than juvenile coho and Chinook salmon and steelhead. As groups of chum salmon reached Alaska, they moved offshore in a generally southwestern direction, although movement was variable and appeared to be strongly influenced by currents (Hartt 1980, Hartt and Dell 1986). A difficulty in these studies is that few numbers of tagged fish were recovered. In the tag recovery information summarized by Hartt and Dell, over 110,000 juvenile salmon and steelhead were caught and 35,259 tagged, of which 4,412 were chum salmon, although only 6 tagged chum salmon (0.1%) were recovered.

A second factor that obscures patterns of oceanic distribution and migration is the extent of delayed ocean migrations and residualism by chum salmon. In the tagging studies by Jensen (1956), juvenile chum salmon remained in nearshore waters beyond the usual time of ocean migration, although the extent of this residualism was unclear (Jensen 1956, Hartt 1980, Fresh et al. 1980, Hartt and Dell 1986). Not all of the chum salmon juveniles tagged in Hood Canal and Puget Sound moved northward toward British Columbia; some remained in Puget Sound throughout the summer, perhaps not leaving until the next spring (Jensen 1956). In November, Hartt and Dell (1986) found juvenile chum salmon in central Puget Sound and in Hecate Strait that averaged 9 in (230 mm) in length, an indication of good growth. It has been hypothesized that these fish may not make an extended northwest migration along the British Columbia/Alaska coast, but may instead proceed directly offshore into the North Pacific Ocean (Hartt and Dell 1986).

The International North Pacific Fisheries Commission (INPFC) has collected a large amount of information on the distribution and origins of high-seas chum salmon. These tagging and scale studies by the INPFC show that although chum salmon from both Asia and North America are distributed throughout the North Pacific Ocean and Bering Sea, Asian chum salmon apparently migrate farther across the Pacific Ocean than do North American fish. Neave et al. (1976) reported that North American chum salmon were rarely found west of the mid-Pacific Ocean beyond long. 175°E, while Asian chum salmon were often found far east of this line. Asian chum salmon have extended their distribution in recent years into the central and eastern North Pacific Ocean, perhaps because of the large increase in releases of hatchery fish in Japan (Kaeriyama 1989, Salo 1991), and because of the change from high-sea to inshore fisheries by Japan's fishing industry (Kaeriyama 1989, Ogura and Ito 1994). Bigler and Helle (1994) and Helle and Hoffman (1995) suggested that the overlap of continental groups may be detrimental to North American chum salmon because maturing chum salmon in the North Pacific Ocean may be at or above carrying capacity.

Limited information exists on stock- or population-specific migration patterns and ocean distributions of chum salmon. Maturing chum salmon in the North Pacific begin to move coastward in May and June and enter coastal waters from June to November (Neave et al. 1976, Fredin et al. 1977, Hartt 1980). No region-specific information on chum salmon migrations to Washington and Oregon has been reported.

Whether the large populations of chum salmon that once inhabited the Columbia River (Rich 1942) had oceanic distributions similar to Puget Sound chum salmon is unknown. As landings in coastal Oregon historically excluded landings on the Oregon side of the Columbia River (Henry 1953), these fish may have had a more southern distribution, like the present distribution of Columbia River coho salmon (Sandercock 1991), and may have returned northward along the Oregon coast.

A.3.2. Distribution




Figure A-75. Historical demographically independent chum salmon populations in the lower Columbia River ESU (Myers et al. 2002).

Chum salmon spawn from Korea and the Japanese island of Honshu, east around the rim of the North Pacific Ocean, to the Columbia River. In the Arctic Ocean, they range from the Laptev Sea in the Russian Federation to the Mackenzie River in Canada.

Chum once were widely distributed in Columbia River tributaries below Celilo Falls (Figure A-75). Some chum historically passed over Celilo Falls on the Columbia River to spawn in the Umatilla and Walla Walla rivers (Nehlsen et al. 1991).

The size and distribution of the Columbia River chum population dropped dramatically in the 1950s (NMFS] 1997), though the runs were still high compared to today. Estimates of chum escapement in 1951 included 3,000 fish to the Lewis River; 1,200 fish to the Chinook and Deep rivers and Crooked and

Jim Crow creeks; 7,500 to Grays River; 3,000 to Skamokawa Creek; 1,000 to the Elochoman River; 1,000 to the Cowlitz River; 600 to the Kalama River; 1,000 to the Washougal River; and 2,700 chum to the Abernathy/Mill/Germany Creek area. These escapement estimates document that natural populations of chum salmon were present in basins throughout the lower Columbia River as recently as the 1950s.

Chum populations in many lower Washington tributaries continued declining into the 1970s. Between 1961–66, the Mayfield fish-passage facility on the Cowlitz River reported collecting only two adult chum (Thompson and Rothfus, 1969). Now, fewer than ten adults are usually collected each year at the Cowlitz Salmon Hatchery (Harza 1999). Chum populations also declined in the Lewis and Kalama rivers. In 1973, WDFW estimated the spawning population in the Lewis and Kalama basins as only a few hundred fish. According to a 1973 report the most dense observed chum spawning observed occurred in side channels and upwelling areas in the lower 6 miles (9.7 km) of the EF Lewis River (WDFW 1973). These declining chum populations illustrate that chum salmon distribution within the lower Columbia River was becoming localized by the 1970s; only a small portion of the historical lower Columbia River distribution possessed natural chum salmon populations.

Near Bonneville Dam, chum salmon return to Hardy and Hamilton creeks and to the lower reaches of Lawton, Good Bear and Duncan creeks. Chum salmon spawn primarily in the lower reaches of Hardy and Hamilton creeks. Annual escapement to these streams near Bonneville Dam averaged about 1,000 fish from 1967–1971. Bryant (1949) noted that a few chum spawned near the mouth of Woodward Creek in 1944. WDFW (1951) reported that chum use the lower portion of Gibbons, Walton, St. Cloud, Duncan, Woodward, Hardy, and Hamilton creeks.

Small numbers of chum salmon also return to other historical spawning tributaries in the lower Columbia River. Aside from the Grays River and Hamilton and Hardy creeks, chum salmon have been observed in Cowlitz, Lewis, Elochoman, Kalama, and Washougal rivers, and in Skamokawa, Germany, and Abernathy creeks. Biologists have monitored chum salmon populations in several of these river systems since 1998 and report that the populations remain extremely low (Uusitalo 2001). Monitoring was expanded in 2000-2002 to include repeat surveys in over 60 tributary streams. Significant spawning populations have been monitored for several years in the mainstem Columbia near Ives Island and Multnomah Falls in the lower Gorge and more recently chum spawning has been monitored in the mainstem Columbia at several spring seeps along the Washington shore near the I-205 Bridge. Some chum salmon may also return to areas above Bonneville Dam. In 1998 and 1999, about 195 and 135 chum salmon, respectively, were observed ascending the fish ladder at the dam (Keller 2001, NMFS 2000).

A.3.3. Genetic Diversity

While many streams in the lower Columbia River support small populations of chum salmon, large enough numbers to conduct a meaningful allozyme analysis have only been found in two regions, Grays River and just downstream of Bonneville Dam (Hamilton and Hardy creeks). Since 1992, collections of several hundred spawning adults have been made from these sites. Spawning has been observed recently in the mainstem Columbia River at the Pierce/Ives Island complex and in seep areas on the Washington shoreline near the I-205 Bridge. Chum adults and juveniles from these mainstem areas have been intermittently collected for genetic analysis from 1998 to 2001. Additionally, small numbers of chum also were collected in the Chinook and Cowlitz rivers in 2000.

The genetic analysis clearly separated the samples of spawning chum into three groups: the Grays River, the below-Bonneville area (Hamilton and Hardy creeks, Ives Island, and the I-205 seeps) and the Chinook River (Sea Resources Hatchery origin). When sampling occurred, the Sea Resources Hatchery was propagating a non-Columbia chum stock from Southwest Washington, but has since switched to Grays River stock. The maximum p-value between the Grays River collection and any other collection

was 0.0001, showing good separation between this population and all others. There also were several high p-value comparisons among the Hardy and Hamilton Creek collections; however, there was no clear distinction among the below-Bonneville (F-test analysis) collections.

Statistical analysis results indicate that the Grays River and below-Bonneville populations are reproductively isolated to a large extent, but that there is no such evidence for isolation among the below-Bonneville areas. Similarities between the collections from the I-205 seeps and the more upstream collections likely indicate opportunistic colonization of a new area. Thus, there appear to be two Columbia chum groups: Grays River and below-Bonneville mainstem and tributaries, which agree with the GDU designations of Phelps et al. (1995).

The genetic samples also showed no apparent differences in age structure of the three aggregations, with 3-year old fish predominating (Keller 2001). These findings resembled findings from scale analysis for chum salmon returning to the Columbia River in 1914, which also indicated that 3-year old fish constituted the majority of the run, 70.4% (Marr 1943). Although scale samples from more recent returns in 2001 and 2002 show a relatively even split between age 3 and age 4 spawning chum (Figure A-76).




Figure A-76. Age composition of natural spawning chum salmon.

A.3.4. ESU Definition

During the proposed listing process for chum salmon in the Pacific Northwest, NMFS received comments stating that chum salmon in the region represent one ESU (Fed. Reg., V64, N57, March 25, 1999, p. 14509). However, the NMFS Biological Review Team felt justified in separating Pacific Northwest chum salmon into 4 ESUs: Puget Sound/ Straight of Georgia, Hood Canal Summer-Run, Pacific Coast, and Columbia River. NMFS defined the Columbia River chum salmon ESU as including all naturally spawning populations in the Columbia River and its tributaries in Washington and Oregon (Fed. Reg., V64, N57, March 25, 1999, p. 14508). More recently, the specific historical populations have been identified by Myers et al. (2003) (Figure A-75).

A.3.5. Life History Diversity

Runs to the Grays River, Hamilton Creek, and Hardy Creek return to the Columbia River basin in October and November (the peak is mid-November). This run time resembles that of chum salmon in rivers along the Washington coast (WDF et al. 1993). Small differences, however, do exist in the timing of spawning for these lower Columbia River populations. Barin (1886) observed that dog salmon (chum) appeared in the Clackamas River by November and spawned soon after. Peak spawning activity for chum salmon in the Grays River and Hamilton and Hardy creeks differs by about a month (November 8 and December 8 or 19, respectively), providing considerable geographic and temporal isolation (Keller 2001). The different spawn timing for these populations suggest that there are likely differences in time of emergence for these chum populations; however, time of emergence data are not available for naturally produced Grays River chum salmon. The spawn timing differences among these naturally producing chum populations supports the genetic analysis that suggests that the Grays River and below Bonneville area (Hamilton and Hardy creeks) are different stocks. Because of limited research focused on chum salmon in the lower Columbia, no other life history differences have been documented among chum salmon populations.

A.3.6. Abundance

The historical chum run size in the Columbia River has been estimated at nearly 1.4 million fish per year. Annual escapements to Washington waters of the lower Columbia mainstem and tributaries declined to an average of 3,000 after 1955 (WDFW 2001). The chum returns remained relatively stable at low levels from 1956-2000, but there were significant increases in returns to Washington waters during 2001-2002 as indicated in index area peak counts in Grays River, Hardy Creek, and Hamilton Creek areas (Figure A-77).

Today, chum salmon are limited almost exclusively to habitats downstream of Bonneville Dam, with the majority of spawning occurring on the Washington side of the Columbia River. Chum spawning returns to the Grays River and Hamilton Creek have been monitored annually since 1944 and returns to Hardy Creek since 1957. Chum spawning in the mainstem Columbia River near the Pierce/Ives Island complex, Multnomah Falls, and I-205 Bridge has been monitored in recent years, and mainstem Columbia River chum spawning population estimates have been made using multiple methods (Rawding and Hillson 2003, Van der Naald et al. 2003, WDFW 2003). Chum salmon also return to spawn in several other lower Columbia tributaries, however most past years' chum surveys have not included tributaries other than the index streams. Beginning in 2000, BPA funded the expansion of WDFW chum surveys to include lower Columbia tributaries in addition to the index stream surveys.




Figure A-77. Chum escapement fish/mile peak counts for combined Washington index tributaries.

The Grays River Index chum monitoring areas include the mainstem Grays River, West Fork Grays River, and Crazy Johnson Creek. Grays River chum are considered depressed by WDFW due to chronically low spawning escapement (WDFW 2002). Average fish-per-mile values in the survey indices show a sharp decline in spawning escapement beginning in about 1955, with an increase beginning in 2001. Average fish-per-mile values from 1955-2000 ranged from a low of 6 fish in 1958 to a high of 521 in 1959 (WDF et al. 1993). The past two years have exceeded the 1959 count with 759 in 2001 and 1,587 in 2002 (Figure A-78).

The Bonneville (Lower Gorge Tributary) chum monitoring index area includes Hardy Creek, Hamilton Creek, and man-made Hamilton Spring Channel. The Bonneville area (Lower Gorge Tributary) chum population is considered depressed due to chronically low spawning escapements (WDFW 2002). Average fish per mile in the Bonneville area survey indices also display a sharp decline beginning in 1955 (based on Hamilton Creek counts). During 1955-2000, Hamilton Creek counts have ranged from a low of 4 in 1979 to a high of 892 in 1963. The 2001 and 2002 fish/mile counts in Hamilton Creek improved to 987 and 888 respectively (Figure A-78). During 1957-2000 the Hardy Creek counts ranged from a low of 1 in 1979 to a high of 636 in 1992. The 2001 and 2002 fish/mile counts in Hardy Creek improved to 711 and 416, respectively (Figure A-78).

Monitoring of lower Columbia chum returns has expanded in recent years to include mainstem spawners and over 60 non-index stream tributaries since. The lower Gorge area includes spawning in the mainstem Columbia near Ives Island and near Multnomah Falls. Mainstem Columbia spawning also occurs further downstream near I-205 Bridge near the Washington shore. Abundance estimates, distribution, and genetic research of mainstem spawners are currently a focus of federal and state agencies in an effort to determine if they are an independent population. The Bonneville mainstem and tributary spawning populations are also components of a reintroduction and low flow salvage plan. An effort is underway to reintroduce chum to Duncan Creek, just downstream of Hardy Creek, and also to 'salvage' adult chum in the mainstem Columbia and in the lower reaches of Hamilton and Hardy creeks during years when flows are too low for fish to spawn successfully or to access the spring-seep spawning areas.




Figure A-78. Fish-per-mile index area counts in Grays River, Hamilton Creek, and Hardy Creek.

Non-index area tributary monitoring includes chum spawning counts in watersheds from the Big White Salmon River downstream to the Chinook River. In 2002, WDFW used the spawning survey data to estimate chum spawning populations for index and non-index areas of the lower Columbia mainstem and Washington tributaries. The total spawning population estimate was 19,403, including 13,850 in index areas and 5,553 in non-index areas (Figure A-79). When including the commercial fishery catch of 14 chum salmon, hatchery escapement of 309, and a Bonneville Dam count of 188 chum, the total minimum chum return to the Columbia River in 2002, excluding Oregon tributary spawning, which is considered to be low, is estimated to be 19,914 fish.

The vast majority of 2002 chum spawning occurred in the Grays River and lower gorge tributaries, and in the mainstem Columbia between I-205 Bridge and Bonneville Dam. However, notable spawning occurred in the Washougal, Lewis, and Chinook river basins, and in Skamokawa, Germany, and Abernathy creeks. The improved chum returns in the past two years has provided a unique opportunity to assess abundance (and presence and absence) in an expanded area of the Lower Columbia basin. This information can assist in discovery of areas which have maintained some capacity to produce chum under current conditions.




Figure A-79. Distribution of Lower Columbia chum spawning populations in 2002.

A.3.7. Productivity

Little is known about the chum salmon production potential of subbasins in the lower Columbia River. Historically, many lower Columbia subbasins were capable of producing chum salmon runs in the thousands. Most watersheds have been negatively affected to some degree by human activity. The primary causes of habitat degradation, and hence salmon productivity, include urban development, dam construction and operation, channelization, riprapping, and timber harvest. For example, in the Cowlitz and Lewis rivers, dam construction has blocked chum salmon access to the majority of the productive habitat within the basin; the upper sections of these basins possess most of the productive salmon habitat. In the Grays River basin, the largest producer of chum salmon, habitat productivity and stability have been reduced as a result of logging road construction, timber harvest, and dike construction in the lower river. The productivity of the Grays River population was also reduced by the loss of Gorley Springs spawning channel that was destroyed in a 1998 flood. Mainstem Columbia chum productivity in the Ives and Pierce islands area is affected by flow operations at Bonneville Dam.

Chum salmon fecundity data are variable. In North America, literature-reported individual fecundity ranged from 2,018 to 3,977 eggs per female. No fecundity data are available for wild chum salmon in the lower Columbia River.

Chum salmon production is affected by the differential losses chum salmon experience during each stage of their life history; the magnitude of these losses varies geographically and temporally and is a reflection of complex interactions between biota and environment. Reported average egg to fry survival in natural streams can be quite variable. For example, Levanidov (1964) and Beacham and Starr (1982) reported egg to fry survival ranging from 6.1-14.2%. Meanwhile, Bakkala (1970) reported annual chum salmon egg to fry survival ranged from 0.1-34.4% and multi-year means ranged from 1.5-27.6%. In controlled stream environments, such as the spawning channel in Abernathy Creek, mean egg to fry survival can be as high as 82.1% (Bakkala 1970). Reported values for mean fry to adult survival range from 0.8-2.8% (Parker 1962, Wolcott 1978). Most mortality suffered by chum salmon in the marine environment occurs within the first few months. Fishing mortality further reduces the number of adults escaping to natal streams to spawn; however, for lower Columbia River chum salmon, fishing mortality has been a minor factor limiting production in recent years (see Chapter 3). Harvest in the lower Columbia River mainstem has been <100 chum/year since 1992. Retention of chum in tributary recreational fisheries is prohibited.

A.3.8. Hatchery Production

The number of hatchery chum salmon produced in Washington is generally small relative to the number naturally produced, and very small compared to the number of hatchery chum salmon produced annually in other areas such as Japan (2 billion) or Alaska (over 450 million; Salo 1991, McNair 1996). In the early 1900s, hatchery managers in the lower Columbia River made little effort to collect chum salmon as the stock declined, primarily because of their low market value in the commercial fishery. The majority of eggs were collected at the Lewis River Hatchery (up to 750 females being spawned in any one year). However, transfers of hatchery chum from outside the Columbia River basin were substantial. During 1913-18, some 30 million chum fry (predominantly from the Chehalis River) were released throughout the Columbia River, including sites above Celilo Falls, the Methow and Walla Walla rivers. At that time, hatchery practices emphasized releasing unfed fry and the success of many of these transfers, especially those far upriver, is doubtful.

Later introductions of non-native chum to lower Columbia tributaries also were generally unsuccessful. For instance, eyed chum eggs of non-local origin were introduced into Spring Channel, a tributary to Hamilton Creek, in the 1970s with no apparent increase in adult production. Several attempts also were made to augment natural chum production in Grays River with releases from the Grays River Hatchery.

Releases from 1982 to 1991 included juveniles resulting from small numbers of adults trapped at Grays River Hatchery and chum of Hood Canal and Japanese origin. Hatchery releases have failed to produce significant adult returns (WDF et al. 1993). The average total number of released juvenile hatchery chum salmon is well below the releases of other salmonid species (Figure A-80).




Figure A-80. Total historical hatchery releases of chum salmon into the lower Columbia basins.

There are currently three hatchery programs in operation in the lower Columbia (Table A-33). Releases of chum fry from the Grays River Hatchery into the Grays River continues at a level of 300,000 per year. In 2002, natural spawn returns to the Grays River basin increased substantially to about 10,000 adults. The hatchery fish were otolith marked, so it will be possible to estimate the proportion of hatchery fish in the return based on samples collected annually on the spawning grounds. Sea Resources Hatchery currently releases 147,500 Grays River stock chum per year into the Chinook River. A new chum hatchery program has commenced at the Washougal hatchery and is part of a multi-faceted lower Gorge natural spawning chum maintenance and enhancement project. The project is aimed at restoration of natural chum in Duncan Creek and also utilizes the Washougal hatchery to support chum production in the mainstem Columbia near Ives Island during years when mainstem Columbia flows are not adequate to fully support spawners, and in Hamilton and Hardy creeks in years when mainstem and/or tributary flows may compromise adult entry into the streams or passage to the most productive spawning areas.

Historical hatchery returns of chum salmon in the lower Columbia River have generally been below 500 fish per hatchery, except for 3 years in the early 1990s at the Sea Resources Hatchery and in 1999 at the Grays River Hatchery (Figure A-81).

Table A-33. Current (2003 brood) chum juvenile hatchery release goals for Washington lower Columbia tributaries.

Basin	Brood Source	Release Goal
Columbia & Tribs near Bonneville Dam	Washougal Hatchery	100,000
Grays	Grays River Hatchery	300,000
Chinook	Grays River Hatchery	147,500
<i>Lower Columbia Total</i>		547,500




Figure A-81. Chum salmon hatchery returns in the lower Columbia River by hatchery, 1957–2002.

Chum salmon have been the primary species raised by the Sea Resources Hatchery, on the Chinook River. This hatchery had a return of 3,000 fish in 1993, the largest hatchery chum salmon return ever documented in the lower Columbia River. Until recently, the Sea Resources Hatchery raised chum using Bear River (Willapa Bay) stock. This non-native stock has now been replaced with local stocks from nearby Grays River (Keller 1999). The Sea Resources Hatchery final returns of the Bear River stock in 1997 and 1998 were 11 and 17 chum, respectively (Keller 1999). In 1999, 60,000 Grays River-stock chum fry were released into the Chinook River from the Sea Resources Facility resulting in 600 three-year-old adults that returned in 2002.

A.3.9. Fishery

Chum Harvest Over Time

Chum salmon once were very abundant in the Columbia River Basin with commercial landings ranging from 1 to 8 million pounds (80,000 to 650,000 fish) in most years before the early 1940s. Chum salmon were harvested in significant numbers in mainstem Columbia River commercial fisheries until their

decline in the early 1950s. Chum were harvested in late fall with most caught in November. Corresponding with the decline in salmon returns, late fall commercial fisheries were reduced. December has been closed to commercial salmon fishing since 1949 and November commercial fisheries have been closed or minimized since 1959. Commercial chum landings gradually diminished during the 1940s and 1950s to less than 50,000 pounds annually by 1959 (Figure A-82). Now there are neither recreational nor commercial fisheries for chum salmon in the Columbia River (ODFW and WDFW 1995). Some chum are taken incidentally in the gill net fisheries for coho and Chinook salmon, but commercial landings have been 500 pounds or less since 1993 (Figure A-82).

NMFS' biological opinions limit the incidental impact of Columbia River fisheries targeting other species to 5% of the annual return of chum listed under the ESA. Since Columbia River chum salmon were listed in 1999, fisheries impacts have remained below the ESA limit.

Few chum are landed in ocean fisheries south of the Strait of Juan De Fuca in Washington. Most landings occur in Canada and Alaska ocean fisheries. There is no specific information on the ocean distribution of Columbia River chum, but it is suspected they migrate similar to Puget Sound stocks, moving to the high seas of the Pacific until they mature and then migrate directly back to the Columbia River. The mature salmon would be present along the coasts of Oregon and Southern Washington in the fall after the ocean seasons have closed, and not present in the chum fisheries farther north.




Figure A-82. Commercial landings of chum salmon in the Columbia River from 1938–2002.




Figure A-83. Fishery exploitation rates over time. Significant Columbia commercial harvest until 1950s. Ocean interception is rare. Columbia commercial harvest steadily decreased since 1950s. Currently, no target commercial or sport fisheries for chum.

Current Chum Harvest

Columbia River chum are harvested incidental to coho, Chinook, and sturgeon during the late fall commercial season (Table A-34). Recent commercial landings have been small, ranging from 0 to 128 chum since 1993, when management measures were implemented to protect late wild coho and chum. According to ESA management limits, Columbia River fall fisheries salmon management requires an incidental impact rate of less than 5% for Columbia River chum salmon; the season structure of the commercial fishery has resulted in less than ESA limits for the annual harvest since Columbia River chum salmon were listed as threatened in 1999. Directed chum salmon ocean commercial fisheries are limited to southern British Columbia and Washington and are managed under agreements resulting from the Pacific Salmon Treaty. Contribution of lower Columbia River chum stocks to these fisheries is expected to be minimal. Generally, most mainstem commercial fisheries are closed before the primary chum salmon migration time. Mainstem Columbia and Washington tributary sport fishing regulations require release of all chum caught while fishing for other species. Chum are not normally encountered in treaty Indian fall fisheries upstream of Bonneville Dam.

Table A-34. Example of current chum harvest.

Fishery	Harvest	Comment
Ocean	<1%	High seas migration likely avoids Northern chum fisheries
Columbia River	1.5%	Incidental to commercial coho fisheries
Tributary	1%	Incidental to steelhead salmon fisheries
<i>Total</i>	<i>2.5%</i>	<i>No directed Columbia Basin fisheries</i>

Chum Harvest Management Details

Directed chum salmon ocean commercial fisheries occur in Alaska, British Columbia, and Washington (Table A-35) and are managed under agreements resulting from the Pacific Salmon Treaty (PST). Although there is very little specific information on the ocean distribution of Columbia River chum salmon, given the timing and distant location of the ocean fisheries that target chum, the contribution of stocks of lower Columbia River chum to these fisheries is expected to be minimal.

Table A-35. Preliminary 2002 chum salmon harvest in ocean fisheries managed under PST.

Fishery	Total 2002 Harvest
ALASKA	
SE Alaska District 104 purse seine	75,218
SE Alaska District 101 drift gill net	144,920
SE Alaska District 106 drift gill net	112,541
SE Alaska District 108 drift gill net	2,017
SE Alaska District 111 drift gill net	231,966
CANADA	
Johnstone Strait	648,000
Strait of Georgia	225,000
Fraser River	100,530
West coast Vancouver Island	554,000
WASHINGTON	
Strait of Juan de Fuca Treaty Indian	1,303
San Juan Islands/Point Roberts Treaty Indian	59,314
San Juan Islands/Point Roberts non-Indian	49,952
Total PST Harvest	2,204,761

Late fall commercial fisheries are regulated by the Columbia River Compact and are focused on harvest of late stock hatchery-produced coho destined for Washington lower Columbia River facilities, and on harvest of white sturgeon remaining on the annual commercial allocation. The Compact exercises time, area, and gear regulations to target hatchery coho and sturgeon while minimizing impacts to chum and late wild Clackamas coho. Clackamas coho are a later-timed late coho which begin entry into the Columbia in late October and early November, similar to Columbia River chum entry time. The commercial season typically closes by late October, unless significant late hatchery coho remain in the river and/or sturgeon harvest allocation is not yet attained. Commercial fisheries extending into the end of October or early November are restricted to larger mesh size to catch sturgeon and avoid chum and coho, or closed in the very lowest part of the lower Columbia where the chum and Clackamas coho presence would be highest. The Compact staff monitors the incidental catch of chum throughout October and recommends preventive regulations earlier in the season, if chum begin to be intercepted earlier than normal. The management strategies employed since 1993 have enabled access to coho, Chinook, and sturgeon while minimizing chum harvest (Figure A-84). Although the most significant reduction in chum harvest occurred in the 1950s, another significant reduction occurred in the 1990s resulting from late fall commercial management changes (Figure A-85).

Late Fall (mid September - November) Harvest of Chinook, Coho, Sturgeon, and Chum (1993-2001 average)




Figure A-84. Average harvest of Chinook, coho, sturgeon, and chum in late fall (mid September to November), 1993-2001.

**Chum Harvest in the Columbia River
Late Fall Commercial Fishery, 1970-2002**




Figure A-85. Late fall commercial harvest of chum salmon in the Columbia River, 1970–2002.

Recreational harvest impacts of chum salmon in the lower Columbia River is minimal. Targeted salmon fisheries in the Grays River were estimated to harvest about 5-10% of the wild chum run prior to 1995. WDFW’s salmon catch record system was originally designed to track Chinook and coho harvest; pink, sockeye, and chum salmon were combined in one category so direct chum salmon catch estimates are unavailable. Retention of chum salmon in the mainstem Columbia River and the tributaries has been prohibited since 1992 in Oregon and since 1995 in Washington; Washington tributaries are closed to chum salmon fishing. Current chum salmon interception rates in Washington tributary recreational fisheries are estimated to be less than 5% with a hooking mortality estimate of 8.6%; these estimates result in a tributary sport fishing mortality rate of less than 1% from 1995 to the present (WDFW 2003).

In a biological assessment of incidental impacts of 2002 Columbia River Fall Fisheries on ESA-listed salmon and steelhead, the *U.S. v. Oregon* TAC estimated a Columbia River chum run size of 2,400 fish and a mainstem Columbia harvest of 38 chum. A WDFW estimate of 1 % mortality rate in Washington tributary fisheries results in a combined Columbia basin harvest impact estimate of 2.58 % of the total chum return in 2002 (Table A-36).

Table A-36. Harvest related mortality estimates for ESA-listed chum salmon in Columbia River basin fisheries during August–December, 2002.

Fishery	Columbia River Chum
Mainstem salmonid sport fishery	0
Mainstem commercial salmon/sturgeon fishery	35
Select Area fall commercial fisheries	3
<i>Total mainstem harvest</i>	38
Tributary sport fisheries	24
Run size at Columbia River mouth	2,400
Harvest/mortality	2.58%

A.3.10. Assessment of Current Status and Limiting Factors

Listing Status

The BRT established by NMFS to examine the status of chum, concluded that the Columbia River ESU is presently at significant risk. The BRT believes the current abundance is probably only 1% of historical levels and the ESU has undoubtedly lost some (perhaps much) of its original genetic diversity. The NMFS chum status review goes on to state:

Although the current abundance is only a fraction of historical levels, and much of the original inter-population diversity has presumably been lost, the total spawning run of chum salmon to the Columbia River has been relatively stable since the mid 1950s and total natural escapement for the ESU is probably at least several thousand per year. Taking all of these factors into consideration, about half of the BRT members concluded that this ESU is at significant risk of extinction; the remainder concluded that the short-term extinction risk was not as high, but that the ESU is at risk of becoming endangered.

Lower Columbia chum salmon, including all naturally spawning populations in the Columbia and its tributaries in Washington and Oregon, were officially listed as threatened on March 25, 1999 (Fed. Reg., V64, N57, p. 14508). NMFS assessed the status of the Lower Columbia River chum salmon ESU and reaffirmed the threatened listing on June 28, 2005 (Fed. Reg., V70, N123, p.37160). Subsequently, the current critical habitat was designated for this ESU on September 2, 2005 (Fed. Reg., V70, N123, p.52630).

Current Status

Extinction risks were estimated to be very high in nine of eleven populations of chum salmon occurring in whole or in part in Washington. All six of the Oregon-only populations are believed to be functionally extirpated (Figure A-86). The majority of the current chum production is occurs in two populations – Washington’s Grays River (viability = moderate) and the Columbia River mainstem and small tributaries in the lower Gorge (viability = high). Current numbers of highly viable populations (0 in both the Coast and Cascade strata, 1 in the Gorge stratum) and average viability scores (Coast = 0.4, Cascade = 0.4, Gorge = 1.8) fall well below the standards established by the WLC-TRT for viability (2 populations per strata at or above 3.0, strata average viability of at least 2.25).

Population viability attributes reflect very low abundance and productivity (Table A-37). The Columbia River historically produced hundreds of thousands of chum salmon (Figure A-82) but only a few thousand remain. The BRT concluded that current abundance is probably only 1% of historical levels. Grays and lower gorge populations generally include 1,000 or more spawners per year. However, remaining abundance of remaining populations is typically fewer than 100 spawners per year. Index counts of chum salmon in significant Washington production areas have remained at consistently low levels since 1950 with the exception of a spike in returns in 2001-2002 due to better-than-average freshwater and ocean production patterns around 2000 (Figure A-77).




Figure A-86. Current status of historical demographically-independent lower Columbia chum populations.

Table A-37. Status and recovery goals for lower Columbia River chum populations (relative to listing baseline).

Population	Contribution	Baseline viability				Obj.	Prod. target	Abundance		
		A&P	S	D	Net			Historical	Baseline	Target
Coast										
Grays/Chinook ^{C,G}	Primary	VH	M	H	M ¹	VH	0% ⁴	10,000	1,600	1,600
Eloch/Skam ^C	Primary	VL	H	L	VL ²	H	>500%	16,000	<200	1,300
Mill/Ab/Germ	Primary	VL	H	L	VL	H	>500%	7,000	<100	1,300
Youngs (OR) ^C	Stabilizing ²	-- ³	-- ³	-- ³	VL	VL	-- ³	-- ³	-- ³	-- ³
Big Creek (OR) ^C	Stabilizing ²	-- ³	-- ³	-- ³	VL	VL	-- ³	-- ³	-- ³	-- ³
Clatskanie (OR)	Primary ¹	-- ³	-- ³	-- ³	VL	H	-- ³	-- ³	-- ³	-- ³
Scappoose (OR)	Primary ¹	-- ³	-- ³	-- ³	VL	H	-- ³	-- ³	-- ³	-- ³
Cascade										
Cowlitz (Fall) ^C	Contributing	VL	H	L	VL	M	>500%	195,000	<300	900
Cowlitz (Summer) ^C	Contributing	VL	L	L	VL	M	>500%	na	na	900
Kalama	Contributing	VL	H	L	VL	M	>500%	20,000	<100	900
Lewis ^C	Primary	VL	H	L	VL	H	>500%	125,000	<100	1,300
Salmon	Stabilizing	VL	L	L	VL	VL	0%	na	<100	--
Washougal	Primary	VL	H	L	VL ²	H+	>500%	18,000	<100	1,300
Clackamas (OR) ^C	Contributing	-- ³	-- ³	-- ³	VL	M	-- ³	-- ³	-- ³	-- ³
Sandy (OR)	Primary	-- ³	-- ³	-- ³	VL	H	-- ³	-- ³	-- ³	-- ³
Gorge										
L. Gorge (WA/OR) ^{C,G}	Primary	VH	H	VH	H ¹	VH	0% ⁴	6,000	2,000	2,000
U. Gorge (WA/OR)	Contributing	VL	L	L	VL	M	>500%	11,000	<50	900

¹ Increase relative to the interim plan.

² Reduction relative to the interim plan.

³ Addressed in the Oregon Management Unit plan.

⁴ Improvement increments are based on abundance and productivity, however, this population will require improvements in spatial structure or diversity to meet recovery objectives.

^C Designated as a historical core population by the Technical Recovery Team.

^G Designated as a historical legacy population by the Technical Recovery Team.

Current Limiting Factors

Net Effect of Manageable Factors

Declines in status of Columbia River chum salmon result from the combined impacts of human activities involving freshwater habitat, estuary habitat, dam construction and operation, fishing, fish hatcheries, and ecological factors such as predation. Impacts of each factor are compounded across the salmon life cycle to drive most populations to current very low levels. Net effects of quantifiable and potentially manageable impacts translate into an estimated 71-99% (average 94%) reduction in abundance and productivity of Columbia River chum populations in Washington (Figure A-87). Thus, current fish numbers represent only 1-29% of the historical production potential in the absence of potentially manageable impacts. Total reductions would be even greater if all human impacts could be effectively quantified. No single factor accounts for the majority of the reduction in fish numbers and the significance of specific factors varies from population to population (Figure A-87, Table A-38).




Figure A-87. Net effect and relative contribution of potentially manageable impact factors on chum salmon in Washington lower Columbia River subbasins. Net effect is the approximate reduction from historical fish numbers as a result of manageable factors included in this analysis

Table A-38. Productivity improvements and impact reduction benchmarks consistent with recovery of Washington lower Columbia River chum populations if reductions are distributed across factors in proportion to their baseline impact.

Population	Productivity	Baseline impacts (listing date reference)							Impacts at target					
	Target	Trib	Est	Dams	Harv	Hat	Pred	Δ	Trib	Est	Dams	Harv	Hat	Pred
<u>Coast</u>														
Grays/Chinook	0%	0.80	0.25	0.00	0.05	0.11	0.03	0%	0.80	0.25	0.00	0.05	0.11	0.03
Eloch/Skam	>500%	0.90	0.25	0.00	0.05	0.03	0.03	50%	0.45	0.13	0.00	0.03	0.01	0.02
Mill/Ab/Germ	>500%	0.90	0.25	0.00	0.05	0.03	0.03	50%	0.45	0.13	0.00	0.03	0.01	0.02
<u>Cascade</u>														
Cowlitz (Fall)	>500%	0.96	0.25	0.00	0.05	0.02	0.03	50%	0.48	0.13	0.00	0.03	0.01	0.02
Cowlitz (Summer)	>500%	0.96	0.25	0.00	0.05	0.02	0.03	50%	0.48	0.13	0.00	0.03	0.01	0.02
Kalama	>500%	0.90	0.25	0.00	0.05	0.01	0.03	50%	0.45	0.13	0.00	0.03	0.00	0.02
Lewis	>500%	0.90	0.25	0.00	0.05	0.01	0.03	50%	0.45	0.13	0.00	0.03	0.01	0.02
Salmon	0%	0.98	0.25	0.00	0.05	0.01	0.03	0%	0.98	0.25	0.00	0.05	0.01	0.03
Washougal	>500%	0.96	0.25	0.00	0.05	0.01	0.03	50%	0.48	0.13	0.00	0.03	0.01	0.02
<u>Gorge</u>														
Lower Gorge	0%	0.40	0.25	0.30	0.05	0.01	0.03	0%	0.40	0.25	0.30	0.05	0.01	0.03
Upper Gorge	>500%	0.97	0.25	0.96	0.05	0.01	0.03	50%	0.49	0.13	0.48	0.03	0.00	0.02

Fisheries

Current fishing impacts on chum salmon are very low and provide no significant opportunity for increasing their numbers through additional regulation of fisheries. No sport or commercial fisheries target chum salmon. Impacts of 5% or less at listing and 3% or less at present are incidental to fisheries for other species (Figure A-88). Historical fishing rates were much greater; these have been steadily reduced to current levels.




Figure A-88. Approximate chum fishery exploitation rates over time and allocation of current rates among fisheries.

Hatcheries

Historical and current hatchery influences on chum are minimal. Hatchery chum salmon have been released into only 4 of 10 Washington populations (Table A-39). Hatchery fish do not comprise a substantial fraction of any naturally spawning chum population and all originate from local wild populations (category 1 brood types). Current chum hatchery programs are focused on reintroduction (Chinook River) and conservation (Duncan Creek).

Inter-specific hatchery predation impacts on juvenile chum range from 0% in basins without significant releases of coho and spring Chinook to a high of 11% in the Cowlitz basin where large hatchery programs are underway.

The Columbia River Hatchery Scientific Review Group (HSRG) recently completed a review of hatchery programs affecting salmon and steelhead populations in the Lower Columbia River. The objective of the review was to change the focus of the Columbia River hatchery system from a harvest supplementation/hydropower mitigation focus to an ecosystem-based approach founded on the idea that harvest goals are sustainable only if they are compatible with conservation goals. Individual population reviews and recommendations can be found at http://www.hatcheryreform.us/mfs/reports/appendix/welcome_show.action.

Table A-39. Presumed reductions in wild population fitness as a result of natural hatchery spawners and survival as a result of interactions with other hatchery species for Washington lower Columbia River chum populations.

Population	Annual releases ^a	Hatchery fraction	Fitness category	Assumed fitness	Fitness impact	Interacting releases ^g	Interspecies impact
Coast							
Grays/Chinook ^b	447,500 ^b	0.25	1	0.9	0.025	40,000	0.002
Eloch/Skam	0 ^c	0.25	1	0.9	0.025	120,000	0.006
Mill/Ab/Germ	0 ^d	0.25	1	0.9	0.025	0	0
Cascade							
Cowlitz	0 ^e	0	--	--	0	2,189,500	0.109
Kalama	0 ^e	0	--	--	0	680,000	0.034
Lewis	0 ^e	0	--	--	0	745,000	0.037
Salmon	0 ^e	0	--	--	0	0	0.001
Washougal	0 ^e	0	--	--	0	120,000	0.006
Gorge							
Lower Gorge	100,000 ^f	0.05		0.9	0.005	0	0
Upper Gorge	0 ^e	0	--	--	0	1,420,000	0.071

^a Annual release goals.

^b Releases include 300,000 in the Grays River to supplement natural production and 147,500 to restore a Chinook River population.

^c Hatchery chum salmon have not been released in the basin since 1983.

^d There is currently no chum salmon hatchery program in Mill, Abernathy, or Germany Creek; hatchery chum salmon have not been released in Abernathy Creek since 1991 or Germany Creek since 1983.

^e There are no records of hatchery chum releases in the basin.

^f A hatchery program recently began at the Washougal Hatchery utilizing Hardy Creek chum for brood stock; releases are planned for Duncan Creek to enhance current chum returns. Additional releases may occur in Hardy and Hamilton Creeks, and in the mainstem Columbia near Ives Island when low flows limit adult access to spawning areas.

^g Includes steelhead and spring Chinook.

Tributary Habitat

EDT analyses suggest that stream degradation has substantially reduced the habitat potential for chum salmon in all Washington lower Columbia River subbasins where analyses have been completed (Figure A-89). Declines in habitat quantity and quality for chum salmon have reduced current productivity potential to only 10-30% and current equilibrium numbers to only 0-11% of the historical template. Substantial stream habitat improvements would be necessary to reach optimum conditions (i.e. PFC) for chum salmon in all pertinent subbasins. Restoration of optimum habitat quality would be expected to increase habitat capacity by 1,000 to 47,000 adult chum per subbasin.

Chum salmon rely on the lower and middle mainstem stream reaches of large streams and rivers where channel instability, low habitat diversity, and sedimentation consistently limit habitat suitability. More detailed descriptions of stream habitat conditions and effects on fish in each subbasin may be found in Volume II of the Technical Foundation.




Figure A-89. Current, optimal, and historical subbasin productivity and equilibrium population size inferred for chum salmon from stream reach habitat conditions using EDT.

Dams

Direct dam effects on Washington lower Columbia chum were limited to the Gorge populations. Assumed dam-related reductions were 20% in the lower Gorge population and 96% in the upper Gorge population. These impacts included assumed passage mortality at Bonneville Dam of 20% for juveniles and an additional 50% for adults. Bonneville Dam inundated an assumed 90% of the chum habitat in the Wind River. Lower Gorge assessments included an assumed 20% reduction in chum productivity in

the Columbia River mainstem as a result of Bonneville Dam operations. Dam operations in the Cowlitz and Lewis River also have the potential to affect downstream habitat conditions for chum but the significance of this impact is unknown.

Mainstem and Estuary Habitat

Chum salmon rear and migrate through critical mainstem and estuary areas soon after emergence and emigration from tributary streams. Mainstem and estuary habitat impacts were assumed to account for approximately a 20-40% reduction in productivity of chum.

Predation

Potentially manageable predation mortality on chum salmon was assumed to average 20% to 25% depending on travel distance from the subbasin to the ocean. Chinook salmon rates were used in the absence of specific chum data. Pikeminnow and tern management is projected to reduce salmonid predation by approximately 50%. Tern predation is almost entirely an artifact of recently established colonies on dredge spoil islands in the estuary but the current rate (9%) is less than half that observed prior to downstream translocation of the Rice Island colony (20%). Pikeminnow predation was greatest for populations that originate in Bonneville Reservoir tributaries (5%), pass the pikeminnow gauntlet in Bonneville Dam forebay and tailrace, and travel the entire 145-mile length from Bonneville to the Estuary. Predation rates by seals and sea lions on adult Chinook were assumed to average 3%.

Summary Assessment

1. Human activities including fishing, hatchery operation, alteration of stream, river, and estuary habitats, hydropower development and operation, and potentially manageable predation have collectively reduced productivity of chum populations to 0-8% of historical levels.
2. Current fishing impacts on chum salmon are very low and these provide limited opportunity for increasing their numbers through additional regulation of fisheries.
3. Existing chum hatchery programs pose no significant risk of reduced wild productivity as a result of interbreeding with potentially less-fit hatchery fish. Inter-specific hatchery predation impacts on juvenile chum range from 0% in basins without significant releases of coho and spring Chinook to a high of 11% in the Cowlitz basin where large hatchery programs are underway.
4. Recovery efforts will require significant improvements in stream habitat quantity and quality. Stream habitat degradation accounts for large declines in chum salmon numbers in all populations.
5. Significant degradation has occurred in estuary and mainstem habitats assumed to be critical to chum salmon life history.

Hydropower impacts on chum salmon are poorly quantified but significant impacts result from operational effects on chum spawning habitat in the mainstem downstream from Bonneville Dam, passage mortality of adults and juveniles at Bonneville Dam, and the inundation by Bonneville Reservoir of lower tributary reaches in the Columbia River Gorge. Cowlitz and Lewis dams have not blocked significant amounts of chum habitat but flood control has altered habitat-forming processes in lower subbasin reaches favored by chum



A.4. Steelhead

Along with cutthroat trout, steelhead/rainbow trout (*Oncorhynchus mykiss*) has the greatest diversity of life history patterns of any Pacific salmonid. Variability in degrees of anadromy and plasticity of life history types between generations is common. Generally, anadromous forms of *O. mykiss* are referred to as steelhead, and resident forms are known as rainbow trout, though some interior subspecies of resident fish are known as redband trout (Behnke 2002). In coastal areas, anadromous and resident forms usually do not co-occur; they are most often separated by a natural or man-made barrier to migration (Busby 1996). Where anadromous and resident forms do co-occur, it is possible for the progeny of individuals of one life history form to exhibit a different life history form (Mullan et al. 1992, Busby 1996). Shapovalov and Taft (1954) reported *O. mykiss* maturing in fresh water and spawning before their first ocean migration, a life history variation that is also reported in cutthroat trout.

In the lower Columbia basin, migrating adult steelhead can occur in the Columbia River year-round, but peaks in migratory activity and differences in reproductive ecotype lend themselves to classifying steelhead into two races: summer and winter steelhead (Figure A-90, Figure A-91).

- Summer steelhead return to fresh water from May to November (Figure A-90), and enter the Columbia River in a sexually immature condition, requiring several months in fresh water to reach sexual maturity and spawn.
- Winter steelhead enter fresh water from November to April (Figure A-91); they are close to sexual maturation and generally spawn shortly after arrival in their natal streams.

Some rivers have both summer and winter steelhead, while others have only one race. Where both runs occur in the same stream, summer steelhead tend to spawn higher in the watershed than do winter forms, perhaps suggesting that summer steelhead tend to exist where winter runs do not fully utilize available habitat (Busby 1996, Withler 1966, Roelofs 1983, Behnke 1992).

In rivers where both winter and summer forms occur, they are often separated by a seasonal hydrologic barrier, such as a waterfall. Coastal streams are predominantly winter steelhead, whereas interior subbasins are dominated by summer steelhead. Historically, winter steelhead may have been excluded from interior Columbia River subbasins by Celilo Falls (Busby 1996).

Initial downstream migration of juveniles generally occurs at either of two life stages: parr (2.36-3.94 in [60-100 mm]), or smolts (5.91-7.87 in [150-200 mm]). Migration of parr leads to rearing in additional areas downstream. Within primary areas of steelhead production, the evidence suggests that capacity for steelhead is generally limited by summer rearing habitat for parr.

A.4.1. Life History and Requirements

Upstream Migration Timing

In the lower Columbia basin, summer steelhead return to fresh water from May to October, and enter the Columbia in a sexually immature condition, requiring several months in fresh water to reach sexual maturity and spawn. Winter steelhead enter fresh water from November to April, and return as sexually mature individuals that spawn shortly thereafter.

Spawning

Steelhead spawn in clear, cool, well-oxygenated streams with suitable gravel and water velocity. Adult fish waiting to spawn or in the process of spawning are vulnerable to disturbance and predation in areas without suitable cover. Cover types include overhanging vegetation, undercut banks, submerged vegetation, submerged objects such as logs and rocks, deep water, and turbulence. Spawning occurs earlier in areas of lower elevation and where water temperature is warmer than in areas of higher elevation and cooler water temperature. Spawning occurs from January through May, and precise spawn timing is related to stream temperature. Bovee (1978) reported a spawning temperature for steelhead of 39-55°F (4-13°C), with an optimum of 46°F (8°C).

Steelhead spawn in areas with water velocities of 1-3.62 ft/sec (30-110 cm/sec) but prefer velocities around 2 ft/sec (60 cm/sec) (Bovee 1978). Female steelhead bury their eggs at a depth of 2-12 in (51-305mm) in redds that occupy up to 60 ft² (5.57m²). More than one redd may be constructed by each female in a season. Spawning sites typically require gravel (0.5-4.5 in [12.5-114mm] diameter) and well-aerated flow. Adult steelhead, unlike salmon, do not necessarily die after spawning but return to the ocean. However, repeat spawning is not common among steelhead migrating several hundred miles or more upstream from the ocean. Researchers have reported that the incidence of repeat spawning increases from north to south (Sheppard 1972, Barnhart 1986). Females have a higher survival rate after spawning than do males (Barnhart 1986). Spawning males usually mate with more than one female and remain in the spawning stream longer than do females (Jones 1974, Barnhart 1986).

Incubation and Emergence

Steelhead are spring spawners, so they spawn at a time when temperatures are typically cold, but increasing. Their spawning time must optimize avoidance of competing risks from gravel-bed scour during high flow and increasing water temperatures that can become lethal to eggs as the warm season arrives. Steelhead eggs hatch in 35–50 days depending on water temperature. Following hatching, alevins remain in the gravel 2 to 3 weeks until the yolk-sac is absorbed (Barnhart 1986).

As with other salmonids, steelhead eggs show a distinct thermal range at which survival is optimum, and survival decreases at temperatures above or below that optimum. Incubation experiments by Kwain (1975) established that survival of steelhead eggs was near 100% at 44.6-50°F (7-10°C), was only about 25% at 59° F (15°C), and was about 50% at either 37 or 41°F (3 or 5°C). These data suggest that, while steelhead eggs have a narrower range of temperatures for optimal survival than either Chinook or coho salmon, their rate of survival drops less at temperatures above or below optimum (Beacham and Murray 1990). While both steelhead and Chinook eggs start dropping in survival as temperatures fall below 41°F (5°C), Chinook egg survival drops to near 0% at 37°F (3°C), while steelhead egg survival drops to about 50% at that temperature. Survival of coho eggs remains near 100% down to 36°F (2°C).




Figure A-90. Summer steelhead life history.




Figure A-91. Winter steelhead life history.

Excessive amounts of fine sediment can reduce survival of salmonid eggs, rearing density of parr, and even survival of overwintering presmolts. Kondolf (2000) points out that although grains smaller than 0.39 in (10 mm) can reduce egg-to-fry survival, some streams with successful natural reproduction have egg-to-fry survivals of considerably less than 50%. Such occurrences indicate that egg-to-fry survival often is not the limiting factor to populations for which juveniles must rear at least a full year in fresh water.

Fry emergence is principally determined by the time of egg deposition and the water temperature during the incubation period. Fry emergence may occur from May through August in the Yakima River subbasin (YIN et al. 1990). In the lower Columbia, emergence timing differs slightly between steelhead races and among Washington subbasins. The different emergence times between races may be a function of spawning location within the watershed (and hence water temperature) or a result of genetic differences of the races. Generally, emergence occurs from March into July, with peak emergence time generally in April and May.

Freshwater Rearing

Following emergence, fry usually move into shallow and slow-moving margins of the stream, where they may aggregate in small schools of up to 10 individuals (Barnhart 1986) in waters 3-14 in (8 to 36 cm) deep (Bovee 1978). Fry tend to occupy shallow riffle habitats, but will occupy pool type habitats during periods of low flow (Barnhart 1986). As they grow, they inhabit areas with deeper water, a wider range of velocities, and larger substrate. Also as they grow older, fry cease schooling behavior and adopt and defend individual territories. Newly emerged fry are sometimes preyed upon by older juvenile steelhead (Barnhart 1986). When water temperatures fall below 39° F (4°C), fry become inactive and conceal themselves in the substrate (Chapman and Bjornn 1969, Barnhart 1986). Bjornn et al. (1977) found significant reductions in juvenile steelhead where gravel substrate was embedded in fine sediment. Sedimentation affects not only the availability of interstitial habitat, but also the productivity of invertebrates, a major food component of juvenile steelhead (Reiser and Bjornn 1979).

Excess fine sediment can reduce rearing density for salmonid parr. Newly emerged fry can occupy the voids of substrate made up of 0.8-2 in (2-5 cm) diameter gravel, but presmolts need cobble (>3 in (7.5 cm)) and boulder-sized rock to provide interstitial spaces they can occupy. Density of juvenile steelhead in summer and winter was reduced by more than half when enough sand was added to fully embed the large cobble substrate in an experimental stream (Bjornn et al 1977). Thompson and Lee (2000) found that probability of moderate to high densities of steelhead parr in Idaho streams decreased as the percentage of watershed with unconsolidated lithology increased ($P>0.05$). They deduced that this type of lithology was prone to sedimentation that could reduce parr survival.

Several studies have found that, although the stock-recruitment relationships show strong density dependence in survival over the length of a full generation from steelhead spawners to their mature offspring, the life stage at which the density dependence shows up is not the spawning-to-fry survival but rather parr-to-smolt survival. For example, Cramer et al. (1985) found from 6 years of data in the Rogue River that subyearling steelhead abundance was a positive linear function of spawner abundance. This finding remained the same after 2 more years of data were added (ODFW 1994). On the Keogh River in British Columbia, Ward and Slaney (1993), found from extensive sampling of all life stages since 1976 that the relationship of eggs-to-fry was linear, while the relationship between fry and smolts showed strong density dependence. Bjornn (1978) found that abundance of yearling steelhead migrants from the Lemhi River over 12 years approached an asymptote at which more age 0 steelhead did not produce more yearling migrants. Thus, carrying capacity of a stream for steelhead is determined by competition for space among parr. This density dependent mechanism affects overall abundance

(Bjornn 1978, Everest et al. 1987, Ward and Slaney 1993), however, it appears to have little affect on steelhead growth (Bjornn 1978, Reeves et al. 1997).

Snorkel observations of rearing steelhead consistently show that they defend individual territories associated with the substrate, and are seldom found in schools (Everest and Chapman 1972, Hillman et al. 1987; Don Chapman Consultants 1989). By manipulating both the stocking density and the density of prey that rainbow trout eat in laboratory stream channels, Slaney and Northcote found that subyearling rainbow defended smaller territories and had fewer aggressive encounters as prey availability increased. Further, as prey density increased, the proportion of fry emigrating from the channels decreased. These behavioral traits lead to displacement of juvenile steelhead when space or food is limiting.

A distinct downstream movement of steelhead presmolts occurs in most streams in the fall, and this could be interpreted to indicate a shortage of winter habitat in their summer rearing area. Studies demonstrate that steelhead presmolts will migrate from an area in the fall where habitat for winter refuge is in short supply, but these fish typically find appropriate winter habitat farther downstream (Bjornn 1978, Tredger 1980, Leider et al. 1986).

Solazzi et al. (2000) found that construction of dammed pools and alcoves and adding woody debris to the constructed units substantially increased the number of steelhead smolts produced during the 4–5 years after restoration treatments in two streams. Solazzi et al. (2000) found that treatments did not increase the number of presmolts in the stream, so they deduced that the increased number of smolts must have resulted from higher survival of presmolts to smolts. While this may have been the case, the increased number of smolts also may have reflected a greater fraction of presmolts remaining in the treatment areas, rather than migrating downstream in search of winter habitat. The proportion of presmolts that migrate downstream in fall decreases as habitat complexity increases in the summer rearing areas.

Research has established that downstream movement of presmolts in the fall is not an indicator of increased mortality. In the Lemhi Basin, Bjornn (1978) found that 30-85% of steelhead presmolts in Big Springs Creek migrated downstream in the fall to overwinter in the Lemhi River, and that presmolt-to-smolt survival ranged from 6-41% for those that migrated, compared to 8-17% for those that stayed in Big Springs Creek. Similarly, Leider et al. (1986) found that most steelhead smolts originating from Gobar Creek had migrated downstream to the Kalama River as presmolts and reared successfully through the winter there. In Nuaitch Creek of British Columbia, Tredger (1980) estimate that 69% of steelhead smolts were actually produced downstream in the Nicoloa River where the fish had migrated to as fry or parr. Thus, evidence indicates that overwintering habitat is important to juvenile steelhead and such habitat may be equally valuable whether it is in the natal stream reach or in some reach downstream.

Observations indicate that juvenile steelhead habitat preferences in order of importance may be depth, velocity, and cover. In a study where effects of cover were held constant, Beecher et al. (1993) compared depth and velocity preferences of steelhead parr 3-8 in (75-200 mm) in a Washington stream that was uniformly lacking in cover; large boulders accounted for less than 1% of surface area and there was no large woody debris (LWD). The stream was believed to be fully seeded with juveniles, because goals for wild spawner escapement were met. Beecher et al. (1993) found that steelhead parr strongly avoided shallow habitats, but once depth was sufficient, velocity preference influenced habitat selection. The strongest effect that depth had on rearing distribution was that parr completely avoided areas with depths < 6 inches. Beecher et al. (1993) found the highest number of parr at depths of 1.6-2.5 ft (0.5-0.75 m), but parr showed the greatest preference for depth > 2.5 ft (0.75 m). Parr avoided depth < 0.8 ft (0.25 m) and velocities < 0.7 ft/sec (21 cm/sec). Beecher et al. (1993) found that most parr were observed at velocities of 0.9-1.1ft/sec (27.4-33.2 cm/sec), but velocities most preferred were

0.7-0.9 ft/sec (21.3-27.1 cm/sec). Preference of steelhead parr for these depths and velocities was also found in an Idaho stream by Everest and Chapman (1972), and was confirmed in an experimental setting by Fausch (1993). Additionally, Beecher et al. (1993) found a significant relationship between parr density and joint preference for depth and velocity combined in a fully-seeded Washington stream.

Fausch (1993) installed replicate experimental structures in a natural stream to evaluate steelhead preference for three types of cover; 1) velocity, 2) lateral cover, and 3) overhead cover. Fausch found that parr selected structures located adjacent to swifter velocities (>0.7 ft/sec [21 cm/sec]) and within 2.9 ft (0.9 m) of natural overhead cover. The preference for overhead cover was stronger than that for velocity cover or lateral cover, but preferences were additive when all were available. Overhead cover was located in or above the water, and presumably provided protection from air or land-based predators. Installation of experimental structures did not attract new fish to the area, but fish present organized around structures with consistent preferences as described. Shirvell (1990) found with experimental placement of rootwads that steelhead parr used velocity and shading created by rootwads, but were not attracted specifically to rootwads. The experiments of Shirvell (1990) concluded that steelhead preference for velocity ranked first, depth second, and light intensity (shade) third. These findings were still consistent with those of Beecher et al. (1993), because depths steelhead used were greater in the Shirvell study than the minimum depth that steelhead strongly avoided in the Beecher et al. (1993) study. Densities of steelhead parr in the stream studied by Shirvell (1990) averaged 1.7 parr/yd² (2 parr/m²) at mean depth of 22 in (56 cm) and mean fish length of 48 in (124 cm).

Findings of steelhead micro-habitat preferences during snorkel surveys in natural streams were consistent with those in experimental settings. Don Chapman Consultants (1989) found that steelhead parr in Washington's Wenatchee River generally selected stations where adjacent velocities were 6-8 times their nose velocity. Stations chosen by parr increased in depth and velocity with fish size. The combined result of steelhead seeking cover from velocity and seeking stations adjacent to higher velocity, as described by Don Chapman Consultants (1989) and Fausch (1993), is that steelhead are often found in riffles or cascades behind boulders. Don Chapman Consultants (1989) found that steelhead concentrated in high gradient reaches ($>5\%$) and usually stationed individually behind boulders where surface turbulence provided cover.

Ward and Slaney (1993) found that placement of boulders resulted in about one steelhead parr rearing per boulder, where none had reared previously. Dambacher (1991) found in the Umpqua River Basin, Oregon, "*Stream channels with relatively high ($0.02/m^2$) and low abundances ($<0.02/m^2$) of age >1 steelhead were separated, with some overlap, by the relative amount of large boulder substrate.*" Although boulders (a form of substrate) can create velocity conditions that steelhead prefer, it appears that it is the velocity patterns, and not the boulders, that influence steelhead habitat selection. Snorkel surveys in the Wenatchee River by Don Chapman Consultants (1989) found that steelhead, when forced by sharp flow reductions to move, selected new stations with similar velocity patterns, but different substrates. Don Chapman Consultants (1989) concluded that associations with specific types of substrate were coincidental rather than causal.

Dambacher (1991) found that streams with greater average riffle depth also had greater densities of steelhead ($R^2 = 0.69$). Similarly, Bisson et al. (1988) found a positive correlation of age 1+ steelhead use with habitat depth over the depth range of 0.4-1.6 in (10-40 cm). In smaller streams where riffles were too shallow, Dambacher (1991) found that age >1 steelhead showed strong electivity for pools, rather than riffles. Dambacher (1991) concludes, "*Stream size (as described by mean riffle depth) apparently creates an upper limit on density of age >1 steelhead rearing during the summer in stream channels of presumably good habitat quality.*" Roper (1995) found in the South Fork (SF) Umpqua River and Jackson Creek that steelhead parr preferred riffles in the lower reaches and pools in the uppermost reach. In nine tributaries sampled in that study, steelhead tended to be more in pools than in riffles. Roper

(1995) concluded that depth or other physical factors may be more important to steelhead preference than habitat selection for pools or riffles.

The preference of steelhead to station themselves where velocities are substantially higher on each side of their focal point results in higher densities as channel roughness (frequency of large boulders) increases. Johnson (1985) performed snorkel surveys of parr densities in a number of western Washington rivers. The published data show over a 10-fold variation between reaches in average parr densities within riffles, and much of that variation was related to whether boulders were the dominant substrate type. All of the reaches with higher than 3 parr/100m² (4 of 18 reaches) had boulders as dominant substrate, and the lowest density (1.5 parr/100m²) in riffles dominated by boulder substrates was equal to the highest density observed in any other unit type. Parr densities in boulder-dominated riffles averaged about five times greater than in riffles with other substrate types as dominant.

Johnson et al. (1993) found that densities of steelhead parr within scour pools tended to increase with increasing structure complexity resulting from wood. They scored wood complexity from 1 to 5, with 1 being little to no wood present, 5 being a large amount of stable complex wood present. The average number of fish/pool during summer generally increased with increasing complexity from wood, and ranged from 2.6 parr/pool in 91 pools lacking complexity to 7.5 parr/pool in 27 pools with a complexity score of 4. Although there were fewer fish in pools with a habitat complexity of 5 compared to those rated as 4, the decrease in fish abundance at the highest pool complexity is likely to be an artifact of sampling bias. A different method of estimating fish abundance had to be used in the pools with the highest cover complexity, because both observation and capture of fish was obstructed. Recent studies have demonstrated that fish abundance in highly complex habitats is usually underestimated by conventional techniques. In winter, differences between complexity scores were not as consistent, and showed a gross indication of increasing fish numbers with greater wood complexity.

The USFWS (1988) studied winter habitat use by juvenile salmonids in the Trinity River. They found steelhead densities were greatest in areas containing cobble and boulder substrates, generally at shallow depths of 0.5-1.5 ft (0.2-0.6 m) with slow current (0-1.2 ft/sec [0-36.6 cm/sec]). Densities of steelhead in areas with high amounts of woody debris, but silt and sand substrate, had less than 1/10 of the steelhead densities found in side channels with cobble and boulder substrate. During winter, the USFWS (1988) found, "*Focal points were nearly always located underneath cobbles or boulders.*" After 2 years of additional sampling for the same study in the Trinity River, USFWS (1990) concluded, "*For steelhead, by far the most important criterion of habitat utilization is the presence of cobbles from 6 to 12 inches in diameter free of sand or silt.*" The preference of steelhead to overwinter in the interstices of cobbles was also reported by Bjornn (1971), Bustard and Narver (1975), and Hartman (1965).

Bjornn and Reiser (1991) reported from studies with steelhead and Chinook in experimental stream enclosures, that more fish remained in pools with a combination of deep water, undercut bank, large rocks, and a bundle of brush than in pools with less cover. The number of yearling steelhead remaining in the stream section increased from 8 to 36 (>4-fold) when brush, large rock, and undercut banks were added. Thus, cobble or boulder substrate in semi-protected water is important as winter habitat for steelhead, either in their natal stream or in a larger channel downstream.

Juvenile Migration

Steelhead exhibit a great deal of variability in smolt age and ocean age. Most steelhead smolt at age 2, though British Columbia and Alaska populations exhibit a significant degree of age 3 smolting, and hatchery fish tend to smolt at age 1 (Busby 1996). Age at smolting tends to be younger in the southern part of the geographic range of steelhead.

Growth rate determines the size and age of smolts, and each of these has a strong influence on survival to maturity. Evidence from several studies shows that faster growing juveniles smolt at a younger age

and that smolt-to-adult survival increases as smolt size increases. Ward and Slaney (1993) found that mean smolt age became younger in a linear relationship to increasing size of age 0 steelhead in mid summer ($R^2 = 0.45$; $P < 0.05$). A similar trend for steelhead to reach smolting at a younger age as growth rate increased was found in the Rogue River (Cramer et al. 1985). Ward and Slaney (1988) found that steelhead smolting at a younger age tended to spend more years at sea than those that smolt at an older age (47% of age 2 female smolts stayed 3 years in the ocean, but only 20% of age 4 smolts stayed 3 years in the ocean). This trend was much stronger for females than males, and allowed a portion of females that grew quickly in fresh water to invest more years for growth in the ocean. Since fecundity is related to female size, females that spent longer time in the ocean produced more eggs (eggs increased by factor of 1.5 from 2-salt to 3-salt female). Additionally, fish that reached a smolt size earlier avoided an additional year of potential mortality in fresh water. A further advantage of earlier age at smolting is the reduced competition for space among the fish remaining in fresh water. Cramer (1986) found that the percentage of age 1 smolts was highly correlated ($R^2 = 0.64$) to growth during spring in the year of smolting, which in turn was highly and positively correlated ($R^2 = 0.71$) to stream temperature during February–April. Cramer (1986) found this same relationship was repeated in the following year to determine what proportion of the remaining cohort would smolt at ages 2 or 3. Ward and Slaney (1988) estimated that smolt-to-adult mortality rate dropped by more than half with each successive year in fresh water, probably as a function of fish size.

Size at smolting also has a strong influence on smolt-to-adult survival. Ward et al. (1989) studied wild steelhead from the Keogh River, BC, and showed that larger smolts survived at a higher rate. Size at smolting was highly correlated to age at smolting. Ward et al. (1989) compared the length at ocean entry for surviving adults (as determined from scales) to the length frequency distribution of smolts (same brood) passing a counting fence, and were able to reconstruct the relationship between smolt length and smolt-to-adult survival. That relationship showed that survival increased about 4-fold as smolt length increased from 6.3-8.7 in (160 mm- 220 mm). However, a portion of the increased mortality at smaller smolt size would also have resulted from the increased tendency of those smaller (and younger) smolts to remain an extra year in the ocean. The classic study by Shapovalov and Taft (1954) also showed for wild steelhead in Waddel Creek, California, that larger smolts survived at a higher rate to maturity. Hatchery managers that raise steelhead are well aware that smolt-to-adult survival of hatchery steelhead increases dramatically as smolt size increases.

Available evidence indicates that a combination of the temperature regime and the size of a stream containing rainbow/steelhead trout (*O. mykiss*) determines whether the population will be predominantly anadromous or resident. Several large river basins on the West Coast have large populations of both resident and anadromous life-history types of *O. mykiss*, but in each case, the types occupy different portions of the basin. Further, breeding experiments indicate that the tendency to be anadromous or resident is an inherited trait, and that resident and anadromous fish in the same basin often breed as independent populations (Zimmerman and Reeves 1999, NMFS 1999). The separation of the primary rearing distributions of resident and anadromous *O. mykiss* within the same basin consistently occurs where there are strong differences in temperature regime. Resident rainbow occur upstream from anadromous forms in areas that are cooler in spring and summer. Resident trout that are larger than steelhead parr will competitively displace juvenile steelhead. The spatial patterns of stream temperature in basins where both the resident and anadromous forms are abundant are consistent with the theory that resident populations will prevail in streams where summer conditions are consistently favorable for growth and survival.

Baltz et al. (1987) studied the physical features that distinguished distribution of four fish species, including rainbow trout, in a 492 ft (150 m) reach of the Pit River, California, where sharp gradient of temperature occurs because of the inflow of cool water from a tributary. Their data show that rainbow were typically holding in water about 63°F (17°C), as were other species, but when river temperatures

warmed to >68°F (20°C), rainbow held positions with stream water that averaged about 64°F (18 °C) while other species (sucker, northern pikeminnow, and hardhead) were at about 68°F (20°C). No rainbow were found in portions of the reach where temperature exceeded 68°F (20°C) on any of four sampling dates. These data suggest that temperatures of 64°F (18°C) or more would stimulate migration of *O. mykiss* out of the area, and if such temperatures were likely to occur at some time in most summers, then natural selection would probably favor anadromy over residency in that stream section.

Migrating smolts are particularly susceptible to predation because they may pass through areas of low cover and high predator concentration (Larsson 1985). Streamflow is important in facilitating downstream transport of outmigrating fish. Along with environmental cues such as photoperiod and temperature, flow is believed to be an important priming factor that triggers migratory behavior once a state of physiological readiness is achieved (Groot 1982). Flow may also influence the rate at which individuals move downstream, although some research indicates that flow may be a secondary factor to photoperiod, as faster-migrating individuals tend to occur at the peak of a run, regardless of low flow patterns that may exist at the time (Bjornn and Reiser 1991). Further, temperature influences the timing of freshwater migration by influencing the rate of growth and physiological development, and by affecting the responsiveness to smolts of other environmental stimuli (Groot 1982). Because of these relationships between migration behavior and flow or temperature, alteration of thermal and flow regimes can influence timing and rates of migration.

While it is likely that dissolved oxygen levels near saturation are required by smolts during the physiologically stressful period of outmigration, supersaturation of dissolved gases (especially nitrogen) has been found to cause gas bubble disease in outmigrating salmonids (Ebel and Raymond 1976). Reiser and Bjornn (1979) hypothesize that steelhead appear to be more susceptible to gas bubble disease because they seem to be less able to detect and avoid supersaturated waters (Stevens et al. 1980).

In the lower Columbia River, emigration of steelhead smolt generally occurs from March to June, with peak migration usually in April or May.

- On the Grays, Cowlitz, Lewis, and Washougal rivers, winter steelhead emigration is from April to May, with peak movement in early May.
- In the Kalama basin, emigration of summer and winter steelhead occurs from March to June, with peak migration from mid-April to mid-May.
- In the Lewis River, summer steelhead smolt emigration occurs from March through May, with peak migration in early May.
- In the Washougal River, summer steelhead smolt emigration generally occurs from April to May.

The dominant age class of emigrating steelhead smolts in the lower Columbia River is age 2.

- In the Grays River, juvenile rearing for the majority of wild winter steelhead lasts 2 years.
- Based on three years of data on the Cowlitz River, 91.1% of winter steelhead smolts resided for 2 years and 8.9% resided for 3 years before their emigration to salt water (Tipping et al. 1979, Tipping 1984).
- On the Toutle River in the Cowlitz basin, emigrating winter steelhead smolts were 86.5% age 2 and 13.5% were age 3 (Schuck and Kruse 1982).
- In the Kalama basin, winter and summer steelhead freshwater rearing primarily lasts 2 years (82.4%) before emigration, but some juveniles reside for 1 (6.2%) or 3 (11.4%) years prior to emigration (Loch et al. 1985).

- In the Lewis River, most winter steelhead juveniles rear for 2 years before emigration (83%), while others do not emigrate until age 3 (17%; Lavoy and Fenton 1983).

Lower Columbia River steelhead average smolt size was estimated at 6.3 in (160 mm). Emigrating steelhead smolts captured from the Kalama River and Gobar Creek ranged in length from 5.4-6.6 in (137.1-167.8 mm).

Estuary Rearing and Growth

Stream-type salmonid populations in the lower Columbia River include winter and summer steelhead. In general, stream-type juvenile salmon reach the lower mainstem and estuary at a relatively large size (> 100mm) and commonly spend less time than ocean-type salmonids rearing in the lower mainstem and estuary. Stream-type juvenile salmonids actively migrate through the lower Columbia River mainstem and estuary. Stream-type salmon are oriented to water column habitats and are typically found throughout the near-surface water column (i.e. top 6 m); they tend to avoid low-velocity areas and are not associated with any specific substrate type.

Juvenile steelhead were present in the Columbia River estuary from February to July of each year of sampling by Bottom et al. (1984); steelhead abundance was greatest in May and relatively low for other months (Bottom et al. 1984). Juvenile steelhead constituted 5% of the total juvenile salmonid catch (Bottom et al. 1984). Steelhead juveniles were distributed throughout the freshwater, estuarine, and marine regions of the estuary; they were most frequently associated with water column habitats (Bottom et al. 1984). Juvenile steelhead moved through the estuary more rapidly than other salmonids; based on catch data, they were present in the estuary for the shortest duration of any of the salmonid group (Bottom et al. 1984). Winter steelhead have been found to migrate at an average rate of 3.3 km/hr, traveling 134-143 km in 32-90 hours (Durkin 1982, Dawley et al. 1986 as cited in USACE 2001). Migration rate of many hatchery groups of juvenile steelhead increased through the estuary (Bottom et al. 1984). As with other salmonids, juvenile hatchery steelhead released further upstream in the basin migrated at a faster rate than juveniles released lower in the system (Bottom et al. 1984).

Steelhead in the Columbia River estuary consumed a relatively even proportion of *Corophium salmonis* (amphipod), *Corbicula manilensis* (bivalve), and adult *Diptera* (Bottom et al. 1984).

Ocean Migrations

In the ocean, steelhead migrate north and south along the continental shelf. Steelhead migrational patterns are generally believed to extend further out in the ocean than other salmonids; however, limited CWT recovery data is available to conclusively confirm this belief. Individuals grow rapidly in the ocean and their size at maturity depends primarily on how long they reside in salt water. Like other anadromous salmonids, steelhead smolt-to-adult survival can be dramatically affected by changes in ocean conditions. High variation in ocean survival between years is the norm for anadromous salmonids, and steelhead populations show some central tendency around a 10-fold range between smolt years (Cramer et al. 2003). Trends in ocean survival have been the driving force in years of low adult returns, where corresponding numbers of outmigrating smolts from previous years have been largely the same as years of higher ocean productivity (Cramer et al. 2003).

A.4.2. Distribution

Historically, steelhead were present throughout the lower Columbia River basins (Figure A-92, Figure A-93). Winter steelhead were distributed throughout most lower Columbia River tributaries from the Grays to the Wind rivers, while summer steelhead were found in the Kalama, NF Lewis, EF Lewis,

Washougal, and Wind River basins. Steelhead continue to be produced naturally in most areas where native steelhead were found, although the abundance of most wild populations is thought to be low.

Spatial separation generally occurs in systems that have both summer and winter steelhead; summer steelhead are usually distributed within headwater areas of the basin, while winter steelhead spawn throughout the lower reaches. The headwater areas are often inaccessible to winter steelhead because of natural barriers that are not passable during the high winter water flows common during winter steelhead migration. These barriers are often passable during the lower flow conditions encountered by summer steelhead during upstream migration. Even in systems that do not have both summer and winter races of steelhead, summer steelhead generally use the upper river reaches, while winter steelhead generally spawn in the lower reaches.




Figure A-92. Historical demographically independent summer steelhead populations in the Lower Columbia River DPS (Myers et al. 2002).



Winter Steelhead

Figure A-93. Historical demographically independent winter steelhead populations in the Lower Columbia River DPS (Myers et al. 2002).

A.4.3. Genetic Diversity

Multiple methods have been used to characterize West Coast steelhead genetic diversity; allozyme electrophoresis, DNA variations, and chromosomal karyotypes (Busby et al. 1996).

Allozyme frequencies have shown two genetically distinct groups of steelhead in Washington, Oregon, and Idaho; a coastal and an inland group (Allendorf 1975, Utter and Allendorf 1977, Utter et al. 1980, Okazaki 1984, Schreck et al. 1986, Reisenbichler et al. 1992). The geographic separation for the two groups appears to be the Cascade Crest, although some uncertainty remains in identifying the boundary. Based on genetic data alone (i.e. no geographical consideration), Leider et al. (1995) suggested that the boundary on the Columbia River is between the Wind and Big White Salmon Rivers. Similar differences have been identified between steelhead from interior and coastal regions of British Columbia (Huzyk and Tsuyuki 1974, Parkinson 1984).

Coastal steelhead have been further segregated into distinct groups; however, not all studies have delineated the same groupings. Hatch (1990) reported evidence for a north-south cline in allele

frequencies for basins larger than 350 km². He suggested Cape Blanco as a geographic feature that limited straying in populations to its' north and south. Reisenbichler et al. (1992) observed that wild coastal steelhead clustered into a north coast group (from the mouth of the Columbia River south to coastal streams just north of the Umpqua River) and a south coast group (from the Umpqua River in central Oregon to the Mad River in northern California). During ESA status reviews, NMFS analyses (Busby et al. 1993, 1994) suggested three genetic population groups: Oregon coast north of Cape Blanco, Cape Blanco to the Klamath River basin inclusive, and south of the Klamath River basin. Leider et al. (1995) expanded the database for genetic data on the Washington coast; samples from certain geographic areas tended to be more similar to each other than they were to samples from other areas and established the following groupings: north Puget Sound (Stillaguamish River and basins north), south Puget Sound, Olympic Peninsula, southwest Washington, and the lower Columbia River (Kalama, Wind, and Washougal Rivers).

Reisenbichler and Phelps (1989) found some genetic variation among steelhead in nine northwest Washington (primarily Olympic Peninsula) basins. Genetic differences among steelhead populations in adjacent drainages were substantially smaller than those reported by Parkinson (1984) for steelhead populations in British Columbia. Reisenbichler and Phelps (1989) and Reisenbichler et al. (1992) suggested that the lower degree of variation in Washington drainages is a result of introgression of hatchery fish into naturally spawning populations. The use of hatchery steelhead in Washington has been extensive and most hatchery steelhead in Washington originated from two primary stocks (Chambers Creek and Skamania stock). However, it is possible that the Reisenbichler and Phelps (1989) study did not cover a large enough geographic area and is not comparable to research in British Columbia (Hatch 1990). Conversely, Phelps et al. (1994) found significant differences between all pairs of populations from Puget Sound and the lower Columbia River. Analyses of the Columbia River data suggests that summer run steelhead in the Wind and Washougal rivers are outliers in the analyses. For example, the Wind River sample contained an allele at a frequency of 15% that was not found in steelhead in any other sample.

In their investigation of genetic variation in steelhead from Idaho to northern California, Reisenbichler et al. (1992) observed that most hatchery populations were significantly different from wild fish. Phelps et al. (1994) also noted statistically significant differences between samples of hatchery and natural steelhead from Puget Sound and the lower Columbia River. These data suggest that at least some native population structure remains. Phelps et al. (1994) also investigated the effect of rainbow trout stocking on steelhead populations in Washington. Because there were large genetic distances between four widely used rainbow stocks and all steelhead populations sampled, Phelps et al. (1994) concluded that there has been little, if any, permanent genetic effect on steelhead populations from widespread rainbow trout stocking.

Using mtDNA analysis, Buroker (unpublished) examined 23 major river systems from Alaska to California and found no evidence for strong geographic structuring of populations because the most common clonal types were widely distributed. Thorgaard (1983) examined chromosomal karyotypes in steelhead from Alaska to Central California. The most common chromosome number was a 58-chromosome karyotype; however, two geographic regions were characterized by steelhead with 59-60 chromosomes: Puget Sound/Strait of Georgia and Rogue River/northern California. In contrast to the allozyme electrophoresis studies, Thorgaard (1983) did not find differences between coastal and inland steelhead populations.

Allozyme frequencies have been unable to demonstrate differences between distinct winter and summer runs of steelhead in the same drainage. Allendorf (1975) and Utter and Allendorf (1977) found that summer and winter steelhead of a particular coastal stream tended to genetically resemble one another more than they resembled populations in adjacent basins with similar run timing. Further allozyme studies support this conclusion in a variety of geographic regions (Chilcote et al. 1980, Schreck

et al. 1986, Reisenbichler and Phelps 1989, Reisenbichler et al. 1992). Shreck et al. (1986) found allele frequencies to be similar for summer- and winter-run steelhead in geographically proximate streams in the Columbia River system. However, in more recent studies, the summer-run stocks have had some extent of hatchery introgression and may not represent the indigenous population. Also, interpretation of the results may be complicated by difficulties in determining run timing of sampled fish. Using chromosomal variability, Thorgaard (1983) was unable to demonstrate a difference in winter- and summer-run steelhead from two systems with limited hatchery introductions (Quinault River in Washington and Rogue River in Oregon). The chromosome number differed between the two systems but was similar in summer and winter steelhead within each river system. Reisenbichler et al. (1992) caution that the absence of difference in allozyme frequencies among groups of fish does not always provide a reliable basis for concluding that these groups are genetically homogeneous. The lack of evidence for genetic differentiation of steelhead within drainages observed in the aforementioned allozyme electrophoresis studies does not rule out the possibility that genetic differences exist in traits affecting survival. For example, genetic differences have been found between sympatric summer- and winter-run steelhead in number of vertebrae, gill rakers, parr marks, rate of maturation in salt water, and level of storage fat in juveniles and adults (Smith 1969).

No clear determination has been made regarding the genetic variation between steelhead and rainbow trout.

A.4.4. DPS Definition

Steelhead found in the lower Columbia River in Washington (as delineated by this recovery plan) fall into three separate DPSs defined by NMFS (Busby et al. 1996):

- Southwest Washington includes steelhead from the Grays and Elochoman rivers, and Skamokawa, Mill, Abernathy, and Germany creeks,
- Lower Columbia includes steelhead from the Cowlitz, Kalama, Lewis, Washougal, and Wind Rivers and Salmon and Hardy creeks, and
- Middle Columbia includes steelhead from the Little White Salmon and Big White Salmon rivers.

On March 19, 1998 NMFS issued a formal notice listing the Lower Columbia steelhead DPS as threatened under ESA. The listed DPS includes only naturally spawned populations of steelhead (and their progeny) residing below naturally and man-made impassable barriers (e.g., impassable waterfalls and dams). The populations that have been identified as comprising the Lower Columbia DPS are shown in Figure A-92 and Figure A-93. More detail on these populations is reported by Myers et al. (2003).

A.4.5. Life History Differences

Adult summer steelhead generally migrate in the lower Columbia River from May through November. Although limited age data are available, dominant age class of returning adults is 2.2 (i.e. 2 years in freshwater and 2 years in the ocean). Lower Columbia stocks of summer steelhead may spawn as early as January through early June in the year following their entry into fresh water. Wild steelhead fry emerge from April–July, depending on spawn timing and water temperature. Juvenile steelhead generally rear in fresh water for 2 years; juvenile emigration occurs from March–June, with peak migration usually in early May. NF Lewis, EF Lewis, Washougal, and Wind summer steelhead conform to these general life history strategies.

Kalama summer steelhead differ slightly in the timing of the adult run and spawning; their adult migration usually occurs over a shorter period (normally from June–October), and spawning occurs from mid-January through April, earlier than other lower Columbia stocks. Although considerable

hatchery summer steelhead releases have occurred in the Kalama basin, Kalama summer steelhead wild stock appears to have retained genetic traits of considerable adaptive value relative to the transplanted hatchery stock (Leider et al. 1995). Also, Kalama summer steelhead have been observed spawning with Kalama winter steelhead; thus, genetic material has been shared to some extent among the steelhead races. These genetic differences of Kalama summer steelhead compared to other lower Columbia summer steelhead may provide some explanation of the different life history strategies of Kalama summer steelhead.

Winter steelhead adult migration timing in the lower Columbia River is generally from November through April. Although, Chambers Creek stock early-run winter steelhead arrive from November to December and spawn earlier than the listed late-run winter steelhead. Although limited age data are available, dominant age class of returning adults is 2.2. Winter steelhead spawn timing for lower Columbia stocks is usually from March through early June, with less time spent in fresh water before spawning than summer steelhead. Wild steelhead fry emerge from April to July, depending on spawn timing and water temperature. Juvenile steelhead generally rear in fresh water for 2 years; juvenile emigration occurs from March to June, with peak migration usually in early May. Winter steelhead in the basins of tributaries to the Grays, Elochoman, and Cowlitz rivers, as well as the lower Columbia Gorge tributaries, conform to these general life history strategies. Most of these basins are in the lower portions of the lower Columbia River basin and have had considerable hatchery influence from Elochoman and Cowlitz stocks.

Winter steelhead in the Kalama, NF Lewis, EF Lewis, Washougal, and Wind basins were identified as a distinct stock partially based on run timing, but the specific run timing for each stock was not provided. Most data suggests that the run timing for these stocks is similar to other lower Columbia River winter steelhead. All of these basins are in the upper portions of the lower Columbia River basin; most stocks have had substantial influence from hatchery stocks from either the Elochoman and Cowlitz or the Skamania Hatchery stock. Kalama winter steelhead are known to spawn from early January to early June; this is an earlier and longer spawning period than other lower Columbia River winter steelhead. Limited escapement surveys suggest that Salmon Creek winter steelhead spawn timing may be earlier than most lower Columbia River winter steelhead.

A.4.6. Abundance

Summer Steelhead

Summer steelhead abundance is naturally quite variable, with variation in ocean conditions believed to be the major driving force in the fluctuating sizes of Pacific salmonid runs and escapement (Pearcy 1992, Beamish and Bouillon 1993, Lawson 1993). Poor ocean conditions during the 1990s resulted in decreased steelhead abundance throughout the lower Columbia River basin. During the early 1980s, steelhead abundance in the Lower Columbia River DPS (including the Upper Willamette DPS) was estimated at approximately 80,000 summer steelhead, although 75% of this estimate was thought to be of hatchery origin (Light 1987). Nehlsen et al. (1991) identified 19 stocks in the Lower Columbia River DPS at some risk of extinction or of special concern. The following designations were given to lower Columbia River stocks covered in this recovery plan:

- high risk of extinction—Cowlitz River summer steelhead, NF Lewis River summer steelhead, and Washougal River summer steelhead
- moderate risk of extinction—Wind River summer steelhead
- special concern—EF Lewis River summer steelhead

Historical (pre-1960) abundance estimates of steelhead populations in the lower Columbia River are scarce. Most summer steelhead stocks in the lower Columbia River are at low abundance levels compared to estimated historical levels as a result of hydro projects, habitat degradation from human activities in the basin (development, logging, etc.), and possible hatchery impacts.

Adequate long-term data is not available for most stocks in the lower Columbia River to address population trends, although available data indicates negative population trends and low abundance compared to historical estimates. Because most of the data sets are short-term, any determinations of population trends may be heavily influenced by short-term climate effects. Of the summer steelhead stocks identified by WDFW in the lower Columbia River in 2002, two were considered depressed (Kalama and Wind) and the status of the remaining three (NF Lewis, EF Lewis, and Washougal) was unknown (Table A-40).

In a recent status report of steelhead in the Lower Columbia River ESU (unpublished), the TRT indicated that, of the six historical summer steelhead populations, not one population could be conclusively identified as naturally self-sustaining. Some degree of natural production was documented in three of six summer steelhead populations. The Ecosystem Diagnosis and Treatment (EDT) model used expected historical habitat conditions to estimate historical steelhead abundance in the Lower Columbia River ESU. A total historical abundance estimate of summer steelhead was 7,294 fish.

All summer steelhead populations in the Washington portion of the lower Columbia River are below WDFW’s natural escapement goals. The Kalama summer steelhead escapement goal of 1,000 fish has not been met since 1995 (Figure A-94). Although spawning escapement estimates of summer steelhead in the Lewis, Wind, and Washougal systems are not available, snorkel index escapement counts exist. Index snorkel counts are estimated to represent 25-70% of natural escapement in each stream (WDFW 1997). Based on these escapement index counts, summer steelhead escapement goals for the EF Lewis River (814 fish), Washougal River (1,210 fish), and Wind River (957 fish) have not been met for at least a decade and likely longer. An escapement goal has not been set for the NF Lewis summer steelhead stock.

Table A-40. Lower Columbia River steelhead stock status as determined by SASSI 2002.

Basin	Stock	Winter Steelhead	Summer Steelhead
Grays River	Grays	Depressed	n/a
Elochoman River	Elochoman/Skamokawa	Depressed	n/a
	Mill	Unknown	
	Abernathy/Germany	Depressed	n/a
Cowlitz River	Mainstem Cowlitz	Unknown	n/a
	Coweeman	Depressed	n/a
	NF Toutle/Green	Depressed	n/a
	SF Toutle	Depressed	n/a
Kalama River	Kalama	Healthy	Depressed
Lewis River	NF Lewis	Unknown	Unknown
	EF Lewis	Unknown	Unknown
Lower Columbia Gorge	Salmon	Unknown	n/a
	Hamilton	Unknown	n/a
Washougal River	Washougal	Depressed	Unknown
Wind River	Wind	Unknown	Depressed




Figure A-94. Counts of total wild and hatchery summer steelhead trapped at Kalama Falls Salmon Hatchery trap, 1976–2001.

Winter Steelhead

Winter steelhead abundance is naturally quite variable, with variation in ocean conditions believed to be the major driving force in the fluctuating sizes of Pacific salmonid runs and escapement (Pearcy 1992, Beamish and Bouillon 1993, Lawson 1993). Poor ocean conditions during the 1990s resulted in decreased steelhead abundance throughout the lower Columbia River basin.

Like steelhead populations throughout their range, lower Columbia winter steelhead have experienced declines in abundance during the past several decades. Wild winter steelhead of native origin exist in the Grays River, Skamokawa Creek, Elochoman River, Mill Creek, Abernathy Creek, Germany Creek, Cowlitz River, Coweeman River, Toutle River, Kalama River, Lewis River, Salmon Creek, Washougal River, Hamilton Creek, and Wind River systems (WDFW 1993).

During the early 1980s, winter steelhead abundance in the Lower Columbia River ESU (including the Upper Willamette ESU) was estimated at approximately 150,000, although 75% of this estimate was thought to be of hatchery origin (Light 1987). Nehlsen et al. (1991) identified 19 stocks in the Lower Columbia River ESU at some risk of extinction or of special concern. The following designations were given to lower Columbia River stocks covered in this recovery plan:

- high risk of extinction—Wind River winter steelhead,
- moderate risk of extinction—Cowlitz River winter steelhead, Washougal River winter steelhead, and
- special concern—Coweeman River winter steelhead, Toutle River winter steelhead, Kalama River winter steelhead, Lewis River winter steelhead.

No estimates of historical (pre-1960s) abundance specific to the Southwest Washington ESU are available. Nehlsen et al. (1991) identified three stocks within this ESU at risk of extinction or of special concern; all identified stocks were within the lower Columbia River portion of this ESU. Winter steelhead in the small lower Columbia River tributaries (including Mill, Abernathy, and Germany creeks) were designated with a moderate risk of extinction, and winter steelhead in the Grays and Elochoman rivers were designated as special concern.

Historical (pre-1960) abundance estimates of steelhead populations in the lower Columbia River are scarce. The largest steelhead population in the lower Columbia River was thought to be in the Cowlitz basin. WDF and WDG (1949) estimated that the steelhead spawning escapement past the Mayfield Dam site was 11,000 fish; considering harvest, the total run size was estimated at 22,000 fish. Naturally spawning populations of steelhead still exist in the lower mainstem Cowlitz, Coweeman, and Toutle River basins, although loss of habitat as a result of dam construction and the 1980 Mt. St. Helens eruption has significantly contributed to decreased abundance. Most other steelhead stocks in the lower Columbia River also are at low abundance levels compared to estimated historical levels as a result of hydro projects, habitat degradation from human activities in the basin (development, logging, etc.), and possible hatchery impacts.

Long-term data adequate to address population trends for most stocks in the lower Columbia River is not available, although available data indicates negative population trends and low abundance compared to historical estimates. Because most of the data sets are short-term, any determinations of population trends may be heavily influenced by short-term climate effects.

In the early 1990s, only two steelhead stocks in the lower Columbia River showed an increasing trend: the South Fork Toutle River and the Kalama River winter steelhead (Figure A-95) (Busby et al. 1996). The increasing trend observed in Toutle/NF Cowlitz River winter steelhead stock was likely a result of continued rebuilding of the stock after the 1980 Mt. St. Helens eruption; abundance of this stock was still considered low.

Also in the early 1990s, only two steelhead stocks in the lower Columbia River were considered healthy: SF Toutle River winter steelhead and Kalama River winter steelhead (WDF et al. 1993). In 2002, WDFW identified 16 winter steelhead stocks on the Washington side of the lower Columbia River; one was healthy (Kalama winter steelhead), eight were depressed, and the status of the remainder was unknown because of a lack of escapement data (Table A-40).

In a recent report on the status of steelhead in the Lower Columbia River ESU (unpublished), the TRT indicated that, of the 17 historical winter steelhead populations, not one population could be conclusively identified as naturally self-sustaining. Some degree of natural production was documented in 9 of 14 winter steelhead populations. The EDT model used expected historical habitat conditions to estimate historical steelhead abundance in the Lower Columbia River ESU. The total historical abundance estimate of winter steelhead based on the EDT model was 18,243 fish. However, WDFW has estimated there were historically 20,000 winter steelhead in the Cowlitz system alone (Hymer et al. 1992, WDFW 1997). Estimates for winter steelhead escapement in Washington tributaries based on redd counts are presented in Table A-41. No redd count data are available for the Cowlitz River. Recent year escapements for all winter steelhead populations in the Washington portion of the lower Columbia River are below WDFW's natural escapement goals. However, the most recent year (2002) return of winter steelhead to the lower Columbia basins was improved for most populations (Figure A-96); also the North Toutle escapement was a post-eruption high.




Figure A-95. Counts of total wild and hatchery winter steelhead trapped at the Kalama Falls Salmon Hatchery trap, 1976–2001.




Figure A-96. Total winter steelhead escapement (Grays River, Skamokawa Creek, Abernathy Creek, Germany Creek, Coweeman River, NF Toutle River, and SF Toutle River combined).

Table A-41. Escapement of winter steelhead in lower Columbia Washington tributaries. Escapement estimates are from redd count data, and represent all natural production, including naturally spawning hatchery winter steelhead.

	Grays	Elo- choman	Skamo- kawa	Aber- nathy	Ger- many	Green	NF Toutle	SF Toutle	Cowee- man	Kalama	EF Lewis	Total
1977										774		774
1978										694		694
1979										371		371
1980										1025		1025
1981										2150		2150
1982										869		869
1983										532		532
1984										943		943
1985						775		1807		632		3214
1986								1595		1081	282	2958
1987						402		1650	889	1155	192	4288
1988						310		2222	1088	1269	258	5147
1989						128	18	1371	392	588	140	2637
1990						86	36	752	522	419	102	1917
1991	716	166		280		108	108	904		1128	72	3482
1992	1224	278	304	246		44	322	1290		2322	88	6118
1993	1086	378	258	88	216	84	165	1242	438	992	90	5037
1994	704	230	208	58	108	128	90	632	362	853	78	3451
1995	426	132	92	34	42	174	175	396	252	1212	53	2988
1996	203	52	112	16	40		251	150	44	853		1721
1997	158	64	128	64	46		183	388	108	537	192	1868
1998	546	100	208	146	90	118	137	374	314	438	250	2721
1999	300	90	200	78	110	72	129	562	126	562	276	2505
2000						124	238	790	290	941	207	2590
2001						79	185	334	284	1085	79	2046
<i>Average</i>	<i>596</i>	<i>166</i>	<i>189</i>	<i>112</i>	<i>93</i>	<i>103</i>	<i>180</i>	<i>642</i>	<i>246</i>	<i>993</i>	<i>139</i>	<i>3138</i>

A.4.7. Productivity

Summer Steelhead

As with other salmonids, steelhead productivity is highly influenced by ocean conditions. Steelhead in the ocean appear to follow a counterclockwise migration pattern in waters east of 167° East longitude, primarily within 27 feet of the surface. Poor ocean conditions during the 1990s resulted in decreased steelhead productivity throughout the lower Columbia River basin. Abundance may have been further decreased with increased predation by Caspian terns in the lower Columbia River during the late 1990s, although there is no basis to determine how the recent estimated tern predation differs from historic losses of juvenile salmonids.

Historically, steelhead production in Washington basins of the lower Columbia River was thought to be high, although most steelhead production was likely from the winter race. The production potential of most lower Columbia River basins is substantially reduced from historical conditions as a result of habitat degradation. Most habitat degradation has resulted from human activity, such as development or logging, although considerable habitat loss occurred in the Cowlitz basin, especially the NF Toutle River, as a result of the 1980 Mt. St. Helens eruption. Major hydro projects in the Cowlitz and Lewis basins have blocked access to approximately 80% of the historical steelhead spawning and rearing habitat within both basins.

The NPCC's smolt production model was applied to many systems within the lower Columbia River in the early 1990s to estimate steelhead production potential. The potential summer steelhead smolt production estimate was 62,273 smolts for the Wind River basin. In the Kalama basin, WDW estimated that potential winter and summer steelhead smolt production was 34,850; the NPCC's model estimated 64,860 (the NPCC model is generally optimistic and can overestimate production potential). From 1978–84, the number of naturally produced steelhead smolts migrating annually from the Kalama ranged from 11,175 to 46,659.

In a recent status report of steelhead in the Lower Columbia River ESU (unpublished), the TRT compared the potential historical habitat available to steelhead to the potential current available habitat. For the entire Lower Columbia River ESU (including Oregon basins), 63% of the historical available steelhead habitat is available today. Most basins have over half the historical habitat still available and some basins still have the majority of the historically available habitat.

Winter Steelhead

As with other salmonids, steelhead productivity is highly influenced by ocean conditions. Steelhead in the ocean appear to follow a counterclockwise migration pattern in waters east of 167° East longitude, primarily within 27 feet of the surface. Poor ocean conditions during the 1990s resulted in decreased steelhead abundance throughout the lower Columbia River basin.

Historically, steelhead production in Washington basins of the lower Columbia River was thought to be high. For example, total run size for steelhead in the Cowlitz River was estimated to be greater than 20,000 fish and, based on preliminary information developed in the process of Lewis hydro relicensing, 10,000 or more may have been produced in the Lewis basin. The production potential of most lower Columbia River basins is substantially reduced from historical conditions as a result of habitat degradation resulting mostly from human activity, such as development or logging. However, considerable habitat loss occurred in the Cowlitz basin, especially the NF Toutle River, as a result of the 1980 Mt. St. Helens eruption. Major hydro projects in the Cowlitz and Lewis basins have blocked access to approximately 80% of the historical steelhead spawning and rearing habitat within both basins.

The NPCC’s smolt production model was applied to many systems within the lower Columbia River in the early 1990s to estimated steelhead production potential. Smolt production estimates were 45,300 winter steelhead smolts in the Grays River; 63,399 winter steelhead smolts in the Cowlitz River; 38,229 winter steelhead smolts in the Coweeman River; and 135,573 winter steelhead smolts in the Toutle River. In the Kalama basin, WDW estimated that potential winter and summer steelhead smolt production was 34,850; from 1978–84, the number of naturally produced steelhead smolts migrating annually from the Kalama ranged from 11,175 to 46,659.

In a recent status report of steelhead in the Lower Columbia River ESU (unpublished), the TRT compared the potential historical habitat available to steelhead to the potential current available habitat. For the entire Lower Columbia River ESU (including Oregon basins), 63% of the historical available steelhead habitat is available today. Most basins have over half of the historical habitat still available; some basins still have the majority of the historically available habitat (e.g. Columbia Gorge tributaries winter steelhead [100%], Kalama River winter steelhead [92%], SF Toutle River winter steelhead [89%], and Salmon Creek winter steelhead [88%]). Some notable exceptions—where very little historical habitat remains available to steelhead—include Cispus River winter steelhead (0%), Tilton River winter steelhead (0%), Upper Cowlitz River winter steelhead (2%), and NF Lewis River winter steelhead (22%).

A.4.8. Hatchery Production

Summer Steelhead

Hatchery releases of summer steelhead occur in the Elochoman, Cowlitz, NF Toutle (released at mouth of Green River), SF Toutle, Kalama, NF Lewis, EF Lewis, and Washougal rivers. Approximately 1 million summer steelhead smolts are released annually within the lower Columbia River ESU (Table A-42).

Table A-42. Current (2003 brood) summer steelhead smolt release goals.

Basin	Brood Source	Release Goal
Washougal	Skamania Hatchery	60,000
NF Lewis	Merwin Hatchery	175,000
NF Lewis	Skamania Hatchery	50,000
EF Lewis	Skamania Hatchery	25,000
Kalama	Skamania Hatchery	30,000
Kalama	Kalama Wild	60,000
NF Toutle	Skamania Hatchery	25,000
SF Toutle	Skamania Hatchery	25,000
L Cowlitz	Cowlitz Trout Hatchery	500,000
Elochoman	Merwin Hatchery	30,000
Lower Columbia Total		980,000

The NMFS status review of West Coast steelhead identified several steelhead broodstocks that have been widely used and have the greatest potential to affect native steelhead populations because of their broad distribution and extensive incorporation in artificial propagation programs (Busby et al. 1996). Among these broodstocks is the Skamania summer steelhead stock. This stock was developed in the late 1950s at the Skamania Hatchery from Washougal and Klickitat river summer steelhead. The Skamania Hatchery is located about one mile from the mouth of the WF Washougal River. Skamania summer steelhead stock has been released throughout Washington, Idaho, Oregon, California, Indiana,

Rhode Island, and North Carolina (Crawford 1979, CDFG 1994). In many cases, Skamania summer steelhead have been introduced to provide angling opportunities where summer steelhead did not naturally exist. However, in the Columbia River, Skamania summer steelhead have been released in basins having endemic summer steelhead populations.

In a recent report of steelhead in the lower Columbia River ESU (WLC-TRT unpublished), the fraction of hatchery fish in the escapement over the last 4 years was calculated for some lower Columbia River basins. For the entire lower Columbia ESU (including Oregon basins), the hatchery fraction of spawners was 24%. For Washington basins, the highest hatchery fractions were observed in the Kalama River summer steelhead (35%) and the Wind River summer steelhead (21%). The hatchery fractions were not calculated for NF Lewis, EF Lewis, and Washougal summer steelhead because of a lack of data.

Most summer steelhead programs (i.e. Kalama, Lewis, Washougal, and Wind River basins) have released fewer than 300,000 juveniles annually during the past 20 years (Figure A-97). The Cowlitz summer steelhead hatchery program released 500,000 or more juveniles and, when including fingerling plants in lakes to provide recreational fishery opportunity, the 2002 releases totaled almost 1.5 million.




Figure A-97. Hatchery releases of summer steelhead in the lower Columbia River by basin, 1982–2002.

Hatchery summer steelhead are widespread and escape to spawn naturally throughout the region. Though hatchery-origin fish contribute substantially to natural production, wild summer steelhead are purported to be reproductively isolated from hatchery fish by spatial and temporal differences. There is overlap, however, between summer and winter steelhead spawn time (WDFW 1993) and some stocks appear to have had substantial hatchery contribution to wild spawning (e.g., Kalama winter and summer steelhead). Nehlsen et al. (1991) identified several stocks from the Lower Columbia and Southwest Washington ESUs as a special concern because of hatchery influence. The impacts of hatchery fish on wild stocks have been studied in the Kalama basin. Skamania hatchery summer steelhead have in the past comprised around 75% of total summer spawning escapement in the Kalama system (WDFW 1993). Even though 40% of returning naturally produced adults are estimated to have at least one parent of hatchery origin, the wild stock has retained genetic traits of considerable adaptive value relative to hatchery stock (Leider et al. 1995, WDFW 1993).

Adult summer steelhead returns to the lower Columbia hatchery facilities are highly variable and are dependent on variable smolt to adult survival as well as variable sport fishery harvest rates. The returns to the hatchery racks only represent a part of the hatchery returns, as the tributary sport fisheries are significant for hatchery steelhead, typically harvesting 50% or more of the fish returning to the rivers (WDFW 2003).

- Hatchery summer steelhead returns to the Cowlitz Salmon Hatchery have been variable, ranging from about 250 to 3,500 and returns to the Cowlitz Trout Hatchery have ranged from about 1,000 to 3,000 (Figure A-98). Although the period for the return data set is short, the trend in returns through the latter part of the 1990s appears to be decreasing. Summer steelhead returns to the North Toutle Hatchery are low, primarily due to the program being developed to rely on releases from other stations as well as an effective tributary sport harvest.
- Hatchery summer steelhead returns to the Kalama River were variable in the latter part of the 1990s (Figure A-98). Although the period for the return data set is short, the trend in returns appears to be decreasing.
- In the Lewis basin, hatchery returns also have been variable in recent years. Summer steelhead hatchery returns to the Lewis River basin have typically ranged between 1,000 and 3,000 fish (Figure A-99).
- In the Washougal basin, returns to the Skamania Hatchery are usually between 500 and 3,000 summer steelhead, though there is little consistency in returns from year to year.




Figure A-98. Hatchery returns of summer steelhead to the Cowlitz and Kalama River basins by hatchery, 1996–2002.




Figure A-99. Hatchery returns of summer steelhead to the Lewis and Washougal River basins by hatchery, 1995–2002.

Winter Steelhead

Hatchery releases of winter steelhead occur in the Grays, Elochoman, Cowlitz, Tilton, Coweeman, Kalama, NF Lewis, EF Lewis and Washougal rivers, and in Salmon Creek. The current (2003 brood) goal is to release 1.2 million winter steelhead smolts and 350,000 subyearling winter steelhead into lower Columbia River tributaries (Table A-43).

Table A-43. Current (2003 brood) Winter steelhead smolt release goals.

Basin	Brood Source	Release	
		Yearling	Subyearling
Washougal	Skamania Hatchery	60,000	
Salmon Creek	Skamania Hatchery	20,000	
NF Lewis	Merwin Hatchery	100,000	
EF Lewis	Skamania Hatchery	90,000	
Kalama	Kalama Falls Hatchery	45,000	
Kalama	Kalama Late Wild	45,000	
Coweeman	Elochoman Hatchery	20,000	
L. Cowlitz	Cowlitz Trout Hatchery	300,000	
L. Cowlitz	Cowlitz Trout Hatchery (late winter)	352,500	
Upper Cowlitz	Cowlitz Trout Hatchery (late winter)	37,500	250,000
Tilton	Cowlitz Trout Hatchery (late winter)		100,000
Elochoman	Elochoman Hatchery	60,000	
Elochoman	Elochoman (late wild)	30,000	
Grays	Elochoman Hatchery	40,000	
Lower Columbia Total		1,200,000	350,000

Hatchery winter steelhead are widespread and escape to spawn naturally throughout the region. Though hatchery-origin fish contribute substantially to natural production, wild winter steelhead are purported to be reproductively isolated from hatchery fish by spatial and temporal differences. The

listed populations of winter steelhead in the ESU are generally late-run; returning and spawning later than the early returning Chambers Creek stock (a Puget Sound stock), commonly used for many of the hatchery programs. (The Chambers Creek Hatchery broodstock originated from Chambers Creek (Tacoma, Washington) in the 1920s. Hatchery winter steelhead from Chambers Creek have been released throughout the lower Columbia River.) There is also overlap, however, between summer and winter steelhead spawn time (WDFW 1993) and some stocks appear to have had substantial hatchery contribution to wild spawning (i.e., Kalama winter and summer steelhead).

The Beaver Creek Hatchery (on a tributary of the Elochoman River) formerly produced approximately 400,000-500,000 winter steelhead smolts annually. The hatchery has utilized broodstock from the Elochoman and Cowlitz rivers and Chambers Creek. Smolts from the Beaver Creek Hatchery have been planted throughout the lower Columbia River. Beaver Creek Hatchery was closed in 1999 due to Mitchell Act funding shortfalls.

In a recent status report of steelhead in the lower Columbia River ESU (WLC-TRT unpublished), the fraction of hatchery fish in the escapement over the last 4 years was calculated for some lower Columbia River basins. For the entire lower Columbia ESU (including Oregon basins), the hatchery fraction of spawners was 24%. For Washington basins, the highest hatchery fraction was observed in the Coweeman River winter steelhead (50%). The winter steelhead hatchery fraction was estimated as <2% for a number of stocks: SF Toutle River (2%), NF Toutle River (0%), Kalama River (0%), EF Lewis (0%), and Washougal River (0%).

- In the Elochoman basin, approximately 100,000 winter steelhead have been released annually, although the 2000 releases totaled approximately 350,000 juveniles (Figure A-100). The current Elochoman program includes Elochoman Hatchery (Beaver Creek) origin winter steelhead smolt releases as well as a Elochoman late wild winter steelhead smolt releases. Annual winter steelhead releases to the Grays River, Skamokawa Creek, and other lower Columbia tributaries (Mill, Germany and Abernathy creeks) have generally been less than 50,000 juveniles annually. Winter steelhead are no longer released into Skamokawa, Mill, Germany and Abernathy creeks.
- The Cowlitz Trout Hatchery, located on the mainstem Cowlitz at RM 42, has two programs producing winter steelhead: an early winter stock derived from Cowlitz River and Chambers Creek stock and a late winter Cowlitz stock used to reintroduce natural production in the upper Cowlitz. Smolts from the Cowlitz Trout Hatchery have been planted throughout the region. In some cases, the influence of hatchery winter steelhead is pronounced. For example, Cowlitz River “wild” winter steelhead are almost all the progeny of Cowlitz Hatchery winter steelhead (WDF 1993). Total winter steelhead annual releases to the Cowlitz River have considerably exceeded releases to other lower Columbia River basins. Releases into the Cowlitz basin were generally over 1 million annually throughout the 1990s (Figure A-101).
- In the Kalama basin, Gobar Pond (four miles up Gobar Creek at RM 19.5) has been utilized as an acclimation site for hatchery steelhead before their release. Yearling hatchery winter steelhead from the Cowlitz or Beaver Creek hatcheries have been released into Gobar Pond for subsequent release to the Kalama basin. Approximately 100,000 hatchery winter steelhead smolts are released in the Kalama River basin annually, except for a release of approximately 300,000 smolts in 2000 (Figure A-101). The Kalama Falls Hatchery is continuing a research program to investigate the effectiveness of using naturally produced late-run winter steelhead for broodstock to replace non-listed, early-run winter steelhead from Beaver Creek.




Figure A-100. Hatchery releases of winter steelhead to the Grays, Elochoman, and lower tributaries basins, 1982–2002.




Figure A-101. Hatchery releases of winter steelhead to the Cowlitz and Kalama River basins, 1982–2002.

- In the Lewis basin, a net pen system has operated on Merwin Reservoir since 1979; annual winter steelhead smolt production has averaged 35,000 fish. The source of the broodstock is from the Merwin Dam trap. Merwin Hatchery (just downstream from Merwin Dam) has produced winter steelhead since the early 1990s and hatchery fish are released primarily within

the Lewis River basin. Releases in the EF Lewis have averaged about 100,000 juveniles annually, while releases to the NF Lewis have been slightly higher (Figure A-102). The 2003 releases were approximately 80,000 smolts into the EF Lewis and 100,000 into the mainstem Lewis from the Island Boat Launch in the lower river. An additional 90,000 Skamania stock winter steelhead were released into the EF Lewis River in 2003.

- Approximately 20,000 winter steelhead also are released annually into Salmon Creek near Vancouver.
- In the Washougal River basin, winter steelhead annual release numbers have been variable, with up to about 180,000 fish released in the late 1980s and approximately 60,000 fish released in recent years.




Figure A-102. Hatchery releases winter steelhead to the Lewis and Washougal River basins, 1981–2002.

In recent years, winter steelhead hatchery returns to most Washington hatcheries in the lower Columbia basin have been below 1,000 fish (Figure A-103). Notable exceptions include the Cowlitz Salmon and Cowlitz Trout Hatcheries where hatchery returns were >5,000 fish during 1996 and 1997. For the four hatcheries with returns below the 1,000 fish level, hatchery returns seemed to mirror one another (i.e. experienced increased or decreased production during same years). The two Cowlitz basin hatchery returns also tracked annually with each other, but did not mirror the returns from the other lower Columbia River hatcheries.




Figure A-103. Hatchery returns winter steelhead in the lower Columbia River by hatchery, 1995–2002.

A.4.9. Fishery

Summer Steelhead Harvest Over Time

Historically, steelhead were harvested in Columbia non-Indian fall commercial gillnet fisheries along with salmon. From 1938-74, steelhead catch ranged from 4,000 to 239,800 (Figure A-104). Non-Indian commercial steelhead harvest has been prohibited since 1975.




Figure A-104. Harvest of summer steelhead in the Columbia River from 1938–2002.

Commercial harvest rates were highest when the Columbia River was open 270 days per year (pre-1943). Summer steelhead commercial harvest was reduced beginning in 1965 when the summer commercial seasons (June and July) were closed to protect depressed summer Chinook populations. The Columbia and tributary recreational fisheries began increasing in effort and total harvest in the 1960s. After 1975, when non-Indian commercial take of steelhead was prohibited, the harvest impacts of lower Columbia steelhead were almost entirely from recreational fisheries (although incidental catch and release mortality of summer steelhead can occur in lower Columbia River fall gill net fisheries). The treaty Indian commercial fishery became more significant after 1968 following federal court decisions clarifying treaty Indian fishing rights. Most treaty Indian steelhead harvest occurs in September during the fall salmon season. The sport harvest of summer steelhead in the lower Columbia tributaries can be significant in years of high production (Figure A-105). Release of wild steelhead in the mainstem Columbia and Washington tributaries was implemented in 1984.

Lower Columbia River steelhead were listed as threatened under the federal ESA in 1998. This ESU includes all naturally spawned summer and winter steelhead in the Columbia River basin and tributaries between the Cowlitz and Wind rivers (inclusive) in Washington and the Willamette and Hood rivers (inclusive) in Oregon, excluding steelhead in the upper Willamette River above Willamette Falls.




Figure A-105. Summer steelhead sport catch harvest (combined wild and hatchery) in Washington lower Columbia basin tributaries.

Current Summer Steelhead Harvest

The Columbia River summer steelhead run is comprised of populations from lower and upper river tributaries. The lower river component of the run is primarily hatchery-produced, derived from Skamania stock, and tends to be earlier timed (May-June) than upriver stocks. The upriver summer steelhead run has historically been separated into A and B groups based on run timing. Group A steelhead include early-returning Skamania stock which pass Bonneville Dam prior to July and are primarily destined for Bonneville Pool tributaries. Group A also includes non-Skamania stock that pass Bonneville Dam from late June through late August on their way to tributaries throughout the Columbia and Snake River basins. Group B steelhead return to the Clearwater and Salmon rivers in Idaho and pass Bonneville Dam from late August through October. In recent years, Group A and B steelhead have not shown the bimodal migration timing peaks. To alleviate fisheries management problems that occur with overlapping runs, the *U.S. v. Oregon* TAC developed a new method in 1999 to assess the returns of Group A and B steelhead. The new index method defined three index stocks: Skamania Index (all fish counted at Bonneville Dam from April 1 to June 30), Group A Index (fish passing Bonneville Dam from July 1 to October 31 that are less than 30 in [78 cm] FL), and Group B Index (fish passing Bonneville Dam from July 1 to October 31 that are greater than or equal to 30 in [78 cm] FL).

Treaty Indian commercial and C&S fisheries in Zone 6 target summer steelhead. Since 1984, the commercial catch has been sampled to determine the percentage of hatchery and wild/natural fish for both Group A and B Index components. Harvest of wild fish in the treaty Indian commercial fishery is compared to the number of wild fish passing Bonneville Dam to determine and manage treaty Indian harvest impacts.

The majority of summer steelhead sport harvest occurs in the tributaries. Tributary harvest is limited to hatchery-marked steelhead only; the date at which this regulation became effective varies by tributary. Hatchery-only harvest restrictions on mainstem Columbia River sport fisheries have been in effect since 1984 to protect wild summer steelhead.

The non-Indian commercial handling of summer steelhead is limited by time and gear restriction. Large mesh gear (minimum of 8 in) is used to harvest Chinook and sturgeon while minimizing the capture of steelhead. Prior to 2002, large mesh gill nets were used to target spring Chinook and sturgeon while minimizing steelhead handling. In 2002, a live capture spring Chinook commercial tangle net fishery was established and resulted in significant steelhead handling in smaller 5.5-in mesh gill net gear. Mesh size was further reduced to avoid capture of steelhead by gilling (instead tangling the fish) and improve survival of released steelhead. Treaty Indian fishery impacts to wild summer steelhead are limited to a maximum of 15% according to a *U.S. v. Oregon* Fall Management Agreement and ESA requirements.

Impacts from mainstem Columbia sport harvest occur primarily during summer months. The tributary sport fishery harvest rates of hatchery summer steelhead are variable, generally ranging from 30-70 %, with the highest rates in the tributaries where there is the most hatchery production (Cowlitz, Lewis, Kalama). The wild steelhead impacts also vary by tributary (3-6%) with the highest impacts in the tributaries with the most hatchery production. Distribution and estimated total harvest exploitation of hatchery and wild steelhead is illustrated in Table A-44.

Treaty Indian commercial and subsistence harvest of Wind River steelhead occurs in the Bonneville Pool, with most harvest occurring during the fall commercial seasons that target fall Chinook. The treaty Indian fisheries are limited to a 15% harvest rate on wild Group B Index steelhead headed to the Snake River, which imposes fishery regulations that result in harvest limitations on other wild steelhead populations, including Wind River steelhead. The harvest of Wind River steelhead by the treaty Indian fall commercial fishery is likely lower than the wild steelhead stocks which pass through the entire treaty Indian fishery from Bonneville Dam to McNary Dam.




Figure A-106. Steelhead fishery exploitation rate over time. Columbia commercial harvest significant until prohibited in 1975. Popular sport fish in mainstem Columbia and tributaries with significant catch since 1950s. Selective fisheries implemented for summer steelhead beginning in 1984.

Table A-44. Example of lower Columbia steelhead harvest exploitation and distribution under current management.

Fishery	H	W	Comment
Ocean	<1%	<1%	High seas migration results in negligible ocean harvest
Columbia River	15%	2%	Sport and incidental commercial impacts
Tributary	55%	5%	Harvest rate varies by tributary
Total Exploitation	70%	7%	Wind River wild approx. 12%, including treaty Indian harvest

Generally, steelhead are not caught in commercial or recreational fisheries in the ocean. Although mark and tag data indicate that high seas steelhead distribution and drift net fisheries overlap, ocean harvest is minimal because the ocean migration pattern of most steelhead is beyond the typical ocean fisheries.

Current harvest impacts for wild steelhead populations below Bonneville Dam are associated with release handling mortality in non-Indian shad, sockeye, and fall salmon commercial fisheries that target salmon and mainstem Columbia, and tributary sport fisheries that target hatchery steelhead and salmon. Wind River steelhead harvest impacts include retained harvest in the treaty Indian fishery above Bonneville Dam. Steelhead incidental capture and handling is minimized through time, area, and gear restrictions. In 1999, an estimated 100 steelhead non-retention mortalities occurred in the fall commercial fisheries.

Summer Steelhead Harvest Management Details

Annual fishery management planning relies on run forecasts to set annual harvest quotas and predict harvest impacts on ESA-listed stocks. Managers utilize numerous forecast methods to estimate annual steelhead runs; different methods are often appropriate for different components of the run and the individual run components can be added to obtain the total run estimate. For example, with the 2003-

2004 upriver summer steelhead run, the 1-salt return was predicted using the recent 5-year average, while the 2-salt return was predicted using a regression relationship between 1-salt and 2-salt returns. Independent estimates were made for the Group A and the Group B Index, and the wild and hatchery components of each.

Columbia River 2002 fall fisheries salmon management was guided by the following restrictions on steelhead harvest:

- Treaty Indian fall fisheries would be managed to limit impacts on wild Group B Index steelhead to 15% or less
- All non-Indian fisheries outside the Snake River basin will be managed for an upriver wild steelhead impact rate not to exceed 2% on Group B index steelhead
- Lower Columbia wild steelhead impacts are limited to 2%.

Summer steelhead sport fisheries exist on the mainstem Columbia River and within the tributaries. Mainstem harvest usually occurs between Tenasillahe Island and Bonneville Dam and few steelhead are caught below Tenasillahe Island. Summer steelhead enter mainstem Columbia River fisheries from March through October, but most of the catch occurs from late May through August. Generally, Group A summer steelhead comprise most of the mainstem Columbia River sport harvest annually (Figure A-107). Steelhead are also handled during warm water recreational fisheries in Columbia River pools from the mouth of the river to Priest Rapids Dam, although impact to steelhead is minor. Creel survey data from 1993-1996 in the area between Bonneville and McNary dams and in 1994 between McNary and Priest Rapids dams indicated only 1% of steelhead were caught by non-salmonid anglers.

Summer steelhead are native to the Kalama, Lewis, Washougal, and Wind basins. Hatchery smolts are released in the Elochoman, Cowlitz, Toutle, Kalama, Lewis, Washougal, and Little White Salmon basins for fisheries opportunity. All summer steelhead streams in Washington have substantial sanctuary water which is closed to fishing; these areas are located in the upper watersheds where an estimated 90% of the wild summer steelhead spawning occurs. Summer steelhead can also be taken incidentally in fall Chinook targeted fisheries; however, the interception rate for non-targeted species is expected to be 1% or less (WDFW 2001). WDFW recreational steelhead selective fisheries are managed to achieve a maximum 10% steelhead mortality for summer steelhead populations below Bonneville Dam. WDFW manages harvest impacts for Wind River summer steelhead to 4% or less because of adverse effects on productivity caused by the operation of Bonneville Dam, fisheries research activities, and mainstem harvest impacts.

Tributary harvest rates for Kalama River wild steelhead have been made by WDFW since 1976. The Kalama River is assumed to be representative of changes in wild steelhead harvest rates after the adoption of wild steelhead release regulations. Harvest rates for both summer and winter steelhead declined from more than 60% during harvest fisheries to less than 10% in the current wild steelhead release fisheries (Figure A-109).

The majority of lower Columbia-origin summer steelhead sport harvest occurs in the tributaries and most of the tributary harvest occurs on the Washington side of the lower Columbia (Figure A-108).




Figure A-107. Lower Columbia River sport catch by steelhead index group, 1969–2002.




Figure A-108. Lower Columbia River sport catch by area, 1969-2000.




Figure A-109. Wild steelhead harvest rates for summer and winter steelhead in the Kalama basin, 1976–2000. (Harvest for summer steelhead after 1984 and winter steelhead after 1991 is adult mortality as a result of hooking mortality in the wild steelhead release fisheries.)

WDFW (FMEP 2003) estimated tributary sport fishery encounter rates and mortality rates for wild summer steelhead in Washington tributary recreational fisheries affecting ESA-listed summer steelhead populations (Table A-45). These estimates include all types of recreational fisheries (for all species) in the Kalama, Lewis, Washougal, and Wind River watersheds.

Since 1984, returns of Group A and Group B summer steelhead have been enumerated at Bonneville Dam and sampled for wild and hatchery percentage. Group A total return (hatchery and wild) has ranged from 115,600 to 515,100 and the percentage of the run that is wild has ranged from 14% to 45%. The Group B total return (hatchery and wild) has ranged from 13,200 to 129,900 and the percentage of the run that is wild has ranged from 8% to 32%. The largest returns were recent, Group A in 2001 and Group B in 2002 (Table A-46).

Table A-45. Estimated take of ESA-listed steelhead in Washington tributary recreational fisheries.

Affected Stock	Anticipated Encounters*	Expected Mortality**
Kalama River summer steelhead	60%	5%
EF Lewis River summer steelhead	40%	3%
Washougal River summer steelhead (mainstem)	40%	3%
Wind River summer steelhead	<10%	1%

* Anticipated encounters are catch and released fish; the numbers represent the percentage of fish from a stock anticipated to be incidentally encountered by anglers of a particular fishery.

** Expected mortality is the hooking mortality of incidentally caught fish; expected mortalities are included in the anticipated encounters in terms of take.

Table A-46. Wild and hatchery contribution to Group A and Group B Index summer steelhead returns to Bonneville Dam, 1984–2003.

Year	Group A Index			Group B Index		
	% Wild	% Hatchery	Total Return	% Wild	% Hatchery	Total Return
1984	27	73	195,700	14	86	98,000
1985	18	82	281,500	32	68	40,900
1986	20	80	287,500	16	84	64,000
1987	45	55	238,300	31	69	45,000
1988	37	63	173,100	22	78	81,600
1989	30	70	193,100	16	84	77,600
1990	23	77	115,600	17	83	47,200
1991	26	74	234,100	22	78	28,300
1992	18	82	241,500	22	78	57,400
1993	21	79	136,700	12	88	36,200
1994	18	82	121,100	20	80	27,500
1995	14	86	180,000	14	86	13,200
1996	15	85	174,300	21	79	18,800
1997	15	85	208,300	11	89	36,600
1998	26	74	134,700	8	92	40,200
1999	32	68	176,400	17	83	22,100
2000	29	71	216,700	21	79	40,900
2001	27	73	515,100	14	86	86,400
2002*	27	73	323,100	25	75	129,900
2003**	25	75	279,600	18	82	64,700

* Preliminary

** Projected

Winter and spring treaty Indian commercial and C&S fisheries in Zone 6 targeting sturgeon can also harvest summer steelhead, but the majority of the treaty Indian summer steelhead harvest occurs during fall fisheries (Figure A-110). In some years, the Treaty Indian Tribes have instituted an 8 in (20 cm) minimum mesh size restriction to reduce the handle of steelhead in the fall fishery and maintain impacts to wild Group B Index steelhead below the ESA limit of 15%. Also, tribal harvest generally focuses on the peak of the fall Chinook run, thereby reducing the number of days needed to fish and minimizing potential impacts to steelhead. In 2001, fall treaty Indian fisheries harvested 29,200 steelhead, which is the largest harvest since 1992.

Since 1985, the commercial catch has been sampled to determine the percentage of hatchery and wild/natural fish for both Group A and B Index components. Harvest of wild fish in the treaty Indian commercial fishery is compared to the number of wild fish passing Bonneville Dam (Table A-47) to determine the percentage of the wild runs that are harvested. These data are used to regulate treaty Indian harvest of wild steelhead. Since 1985 the treaty Indian harvest rate has ranged from 2%-21% for Group A wild steelhead and from 11%-37% for Group B wild steelhead. However, recent year harvest rates have been less than 10% for wild Group A steelhead and less than 15% for wild Group B steelhead.




Figure A-110. Steelhead harvest in treaty Indian fisheries by season, 1970–2002.

Table A-47. Wild steelhead catch (in thousands of fish) in treaty Indian fisheries by index group, 1985–2001.

Year	Group A Index		Group B Index	
	Number	% of Wild Run	Number	% of Wild Run
1985	10.8	20.7	4.0	31.0
1986	7.8	13.8	2.7	26.7
1987	16.8	15.7	5.2	37.2
1988	11.0	17.1	4.2	23.4
1989	9.0	15.9	4.3	35.0
1990	4.3	16.0	1.9	21.5
1991	8.8	14.6	1.9	30.0
1992	7.2	16.2	3.3	26.3
1993	4.4	15.2	0.8	19.1
1994	2.2	10.3	1.0	18.6
1995	2.7	10.4	0.3	18.6
1996	2.3	8.9	1.4	34.8
1997	3.2	10.4	0.6	14.3
1998	3.1	8.8	0.5	15.6
1999	4.3	7.6	0.5	12.6
2000	2.3	3.7	1.0	11.4
2001	5.5	4.0	1.4	11.4
2002	2.4	2.0	1.1	3.4

The estimated harvest related mortality of listed steelhead ESUs for 2002 non-Indian fall fisheries is summarized in Table A-48 and for 2002 treaty Indian fisheries in Table A-49. This information is developed by the *U.S. v. Oregon* Technical Advisory Committee in an annual Biological Assessment and submitted to NMFS for reference when considering fisheries in their Biological Opinion. If fisheries are

determined to meet ESA harvest limits they are authorized with an Incidental Take Permit which is delivered to the State and Tribes prior to fisheries being set.

Table A-48. Harvest related mortality estimates for listed steelhead ESUs in proposed Columbia River basin fisheries* during August—December 2002.

Fishery	Upper Columbia		Snake River Wild	Lower Columbia Wild	Mid Columbia Wild	Total Wild	Total Listed
	Hatchery	Wild					
Mainstem salmonid sport fishery (below Bonneville)	763	15	204	20	179	419	1,182
Mainstem salmonid sport fishery (above Bonneville)	1,899	60	382	72	451	965	2,864
Mainstem commercial salmon/sturgeon fishery	28	3	126	0	35	164	192
Select Areas fall commercial fisheries	0	0	0	0	1	1	1
Wanapum tribal subsistence fishery	17	3	0	0	0	3	17
TOTAL Harvest/Mortality	2,704	81	713	92	665	1,552	4,256
Run Size at Columbia River Mouth	21,771	5,496	75,400	31,068	65,716	177,680	199,451
<i>Harvest/Mortality Rate (%)</i>	<i>12.4</i>	<i>1.5</i>	<i>0.9</i>	<i>0.3</i>	<i>1.0</i>	<i>0.9</i>	<i>2.1</i>

¹ Includes only those fisheries that have mortality of listed steelhead.

Table A-49. Estimated total harvest of steelhead in the 2002 proposed treaty Indian fall fisheries and incidental harvest by ESU.

ESU	Treaty Indian Fall Season Fisheries			
	Zone 6	Tributaries	Hanford Reach	Total
TOTAL HARVEST	29,150	1,710	100	30,960
Lower Columbia River Steelhead	27	3	0	30
Harvest Rate				0.96%
Mid Columbia River Steelhead	1,880	157	0	2,037
Harvest Rate				3.94%
Upper Columbia River Steelhead	1,725	0	100	1,825
Harvest Rate				7.51%
Snake River Steelhead	5,751	0	0	5,751
Harvest Rate				7.81%

Winter Steelhead

Winter steelhead are an important recreational fishery throughout their range. The vast majority of the harvest of winter steelhead occurs in the tributaries of the lower Columbia. In most areas, provisions for separating hatchery and wild fish were not in place until 1987. Since 1987, hatchery steelhead have been marked with an adipose fin clip. Regulations mandating the release of wild fish are in place.

Mainstem Columbia River harvest is typically small and incidental to spring Chinook fisheries. Since 2001 spring Chinook salmon fisheries in the lower Columbia have been extended due to implementation of selective fishing for spring salmon. The extended commercial and sport fisheries have increased handling of wild winter steelhead compared to the previous 25 years, but the total impact of lower Columbia steelhead listed under the ESA is limited to 2% or less of the annual return.

Generally, steelhead are not caught in commercial or recreational fisheries in the ocean. Although mark and tag data indicate that high seas steelhead distribution and drift net fisheries overlap, ocean harvest is minimal because the ocean migration pattern of most steelhead is seaward of the ocean salmon fisheries. Non-Indian commercial harvest of steelhead in the Columbia River has been prohibited since 1975. Mainstem Columbia sport fisheries have been regulated for selective harvest of adipose fin-marked hatchery fish and have required the release of wild steelhead since 1984. Some Washington tributary winter steelhead recreational fisheries were restricted to wild steelhead release in 1986. The remaining tributary winter steelhead recreational fisheries were restricted to wild steelhead release in 1992, with the exception of the South Fork Toutle, which began wild release regulations in 1994.

Current harvest impacts for wild steelhead populations below Bonneville Dam are associated with release handling mortality in non-Indian commercial fisheries that target salmon and mainstem Columbia and tributary sport fisheries that target hatchery steelhead and salmon. Wind River steelhead harvest impacts include retained harvest in the treaty Indian fishery above Bonneville Dam.

Winter Steelhead Harvest Over Time

Historically, steelhead were harvested in Columbia non-Indian winter commercial gillnet fisheries. From 1953-74, steelhead catch ranged from 2,400 to 23,400 (Figure A-111). Non-Indian commercial steelhead harvest has been prohibited since 1975. Commercial harvest rates were highest when the Columbia River was open 270 days per year (pre-1943). Commercial harvest of wild winter steelhead was lower than summer steelhead, because beginning in 1909, the season was closed from early March to late April. The Columbia and tributary recreational fisheries began increasing in effort and total harvest in the 1960s. After 1975, when non-Indian commercial take of steelhead was prohibited, the harvest impacts of lower Columbia steelhead were almost entirely from recreational fisheries. The treaty Indian commercial fishery became more significant after 1968 following federal court decisions clarifying treaty Indian fishing rights. Most treaty Indian steelhead harvest occurs in September during the fall salmon season.

Steelhead incidental capture and handling occurs in sturgeon and winter salmon fisheries; capture of steelhead is minimized through time, area, and gear restrictions. For example, the lower Columbia winter commercial gill net fishery was restricted to a 7 1/4 in minimum mesh size in 1970 to reduce steelhead handle. In 1975, the minimum mesh size restriction was increased to 8 in, concurrent with the prohibition of non-Indian commercial steelhead harvest. From 1975-90, a seasonal average of less than 500 steelhead were handled annually as a result of incidental capture in winter fisheries. Monitoring data for the same period indicates that steelhead immediate mortality from handling was about 17%. Monitoring in the 1990s by the Marine Mammal Observer Program suggests that steelhead immediate mortality from handling may be lower than 17%.

Limited numbers of winter steelhead are harvested annually in the treaty Indian winter commercial fishery (Figure A-111). Most harvest likely occurs in Bonneville Pool. The winter treaty Indian fishery targets sturgeon. The 2002 winter commercial gill net landings totaled 78 steelhead; all steelhead were caught in the Bonneville Pool.

Winter steelhead annual recreational harvest in the lower Columbia River and tributaries has exceeded commercial harvest since the 1950s (Figure A-111).




Figure A-111. Harvest of winter steelhead in the Columbia River from 1953–99.

Steelhead incidental capture and handling occurs in sturgeon and winter salmon fisheries; capture of steelhead is minimized through time, area, and gear restrictions. From 1975-1990, a seasonal average of less than 500 steelhead annually were handled as a result of incidental capture in winter fisheries. Monitoring data for the same period indicates that steelhead immediate mortality from handling was about 17%. Monitoring in the 1990s by the Marine Mammal Observer Program suggests that steelhead immediate mortality from handling may be lower than 17%. Recent year (2002-2003) handling of winter steelhead has increased in the winter/spring commercial fishery due to gear and seasonal structure changes associated with selective spring Chinook salmon fisheries.

Lower Columbia River steelhead were listed as threatened under the federal ESA in 1998. This ESU includes all naturally spawned summer and winter steelhead in the Columbia River basin and tributaries between the Cowlitz and Wind rivers (inclusive) in Washington and the Willamette and Hood rivers (inclusive) in Oregon, excluding steelhead in the upper Willamette River above Willamette Falls.

Current Winter Steelhead Harvest

Winter steelhead enter the Columbia River from November to May; the hatchery run peaks from December to January and the wild run peaks from March to April. Winter steelhead are destined

primarily for tributaries below Bonneville Dam; a few Bonneville Pool tributaries support winter steelhead runs.

Winter steelhead sport fisheries occur primarily in Columbia River tributaries. Hatchery-only harvest restrictions on mainstem Columbia River sport fisheries have been in effect since 1984 to protect wild steelhead. Release of all wild steelhead in recreational fisheries is now required basin-wide.

Limited numbers of winter steelhead are harvested annually in the treaty Indian winter commercial fishery. Most harvest likely occurs in Bonneville Pool. The winter treaty Indian fishery targets sturgeon.

Non-Indian commercial handling of wild winter steelhead occurs during winter and spring salmon seasons. Prior to 2002, large mesh gill nets were used to target spring Chinook and sturgeon while minimizing steelhead handle. In 2002, a live capture spring Chinook commercial tangle net fishery was established and resulted in a significant steelhead handle in smaller 5.5-in mesh gill-net gear. WDFW and ODFW are continuing to experiment with season structure and gear to avoid excessive impacts to winter steelhead during live capture spring salmon seasons. Treaty Indian fishery impacts to wild steelhead are limited to a maximum of 15% according to a *U.S. v. Oregon* Fall Management Agreement and ESA requirements.




Figure A-112. Steelhead fishery exploitation rate over time. Columbia commercial harvest significant until prohibited in 1975. Popular sport fish in mainstem Columbia and tributaries with significant catch since 1950s. Selective fisheries implemented for summer steelhead 1984-92.

Winter Steelhead Harvest Management Details

Winter steelhead enter the Columbia River from November to May; the hatchery run peaks from December to January and the wild run peaks from March to April. Winter steelhead are destined primarily for tributaries below Bonneville Dam; a few Bonneville Pool tributaries support winter steelhead runs.

Recent year (2002-2003) handling of winter steelhead has increased in the winter/spring commercial fishery because of gear and seasonal structure changes associated with selective spring Chinook salmon fisheries. The developing tangle net commercial fishery on the lower Columbia River has successfully targeted hatchery spring Chinook while releasing wild Chinook, with an estimated 10% catch and release mortality on wild spring Chinook. Because the impact to wild spring Chinook has been substantially reduced as a result of the reduction in gill net mesh size to 5.5 in maximum, the fishery remains open longer, creating more potential opportunities for encounters with wild winter steelhead.

The 2002 fishery was open much later in the year (i.e. into late March) than recent commercial seasons and the fishery timing coincided with the early part of the peak of the wild winter steelhead run. While the reduction in mesh size allowed for the release of Chinook, steelhead may be more susceptible to mortality resulting from injuries sustained during gill net entanglement.

Preliminary data from the 2002 winter season indicate that the steelhead catch greatly exceeded the preseason catch expectations because of an extremely large 2002 winter steelhead run, fishery timing, and gear employed. A total of 21,600 steelhead were handled during the 2002 fishery, of which 8,640 (40%) were marked and 12,960 (60%) were unmarked. It is possible that some steelhead were handled more than once. Preliminary monitoring results indicate that the immediate mortality rate for steelhead was 2.0%, which results in an immediate mortality estimate of 260 unmarked steelhead. A gear components study was conducted using test fishery and tangle net fishery data from 2000–02 to assess the effect of different mesh sizes on steelhead and Chinook condition and mortality after gear entanglement. Immediate and total mortality of steelhead was lowest with 5- or 6-in mesh, although, these mesh sizes also had the lowest sample size in the study (Table A-50). The 5.5-in mesh gill net had the lowest percentage of steelhead categorized as condition 1 (vigorous, not bleeding) after capture; the highest percentage of condition 1 steelhead after capture occurred with the 5-in mesh gill net (Table A-50).

For the 2003 winter season, the Columbia River Compact considered regulations requiring large mesh size early in the season to limit steelhead handle and small mesh size (less than 4.25-in) later in the season to promote high survival rates of released steelhead. Other considerations to reduce steelhead handle are reduced fishery effort during peak abundance time of winter steelhead and the use of large (>12 in) mesh excluder panels on the top 5-10 feet of the net.

Winter steelhead are native to all major and most minor basins in the lower Columbia River. Steelhead in tributaries downstream of the Cowlitz River are considered part of the SW Washington ESU and are not listed under the ESA. Winter steelhead sport fisheries occur mostly in Columbia River tributaries; fisheries are primarily in the Grays, Elochoman (both in the SW Washington ESU), Cowlitz, Toutle, Coweeman, Kalama, Lewis, Salmon, and Washougal basins.

Fisheries targeting winter steelhead are concentrated from December to February and close by March 15, except in the Cowlitz, Kalama, Lewis, and Washougal basins where winter steelhead fisheries extend through May 31. The closed periods in the tributaries are set to protect wild spawning steelhead. Winter steelhead are also taken incidentally in spring Chinook targeted fisheries from February–May; however, the interception rate for non-targeted species is expected to be 1% or less (WDFW 2001). Hatchery-only harvest restrictions on mainstem Columbia River sport fisheries have been in effect since 1984 to protect wild steelhead. In Washington, some tributary winter steelhead fisheries adopted wild steelhead release regulations in 1986; the remaining tributary winter steelhead fisheries adopted wild steelhead release regulations in 1992, except for the South Fork Toutle which went to wild steelhead release in 1994. Release of all wild steelhead in recreational fisheries is now required basin-wide.

WDFW recreational steelhead selective fisheries are managed to achieve a maximum 10% steelhead mortality for winter steelhead populations both above and below Bonneville Dam.

The estimated encounter and take of wild winter steelhead in Washington tributary recreational fisheries is summarized in Table A-51 (WDFW 2003).

Table A-50. Summary of catch rate, condition, and survival for steelhead captured in various spring fisheries, 2000–02^a.

	Mesh Size					
	3.5 ^b	4.0 ^c	5.0 ^d	5.5 ^c	6.0 ^d	SE ^d
Sample Size	105	93	7	45	13	9
CPUE ^e	0.44	0.52	0.34	0.75	0.56	0.40
Immediate Mortality ^f	15	12	0	9	1	0
Total Mortality ^g	NA	4 of 14 (28.6%)	0 (0%)	6 of 15 (40.0%)	2 of 13 (15.4%)	2 of 9 (22.2%)
Capture Condition ^h	1: 63%	1: 53%	1: 100%	1: 39%	1: 50%	1: 40%
	2: 3%	2: 3%	2: 0%	2: 14%	2: 0%	2: 20%
	3: 16%	3: 23%	3: 0%	3: 20%	3: 33%	3: 30%
	4: 0%	4: 4%	4: 0%	4: 7%	4: 8%	4: 0%
	5: 18%	5: 18%	5: 0%	5: 20%	5: 8%	5: 10%
Method of Capture ⁱ	Tooth tangle to max	Max to opercle	Max to opercle	Opercle to wedge	Wedge	Opercle to wedge

^a The information used in this table is pooled from various test and experimental fisheries conducted over three years. Many factors varied among the studies, including study protocol, personnel, and data collected.

^b CPUE and immediate mortality data from 2000 & 2001 test fishing and 2001 permit fishery.

^c CPUE and immediate mortality data from 2001 permit fishery and 2002 test fishing; total mortality from 2002 test fishing.

^d All data from 2002 test fishery.

^e CPUE standardized to 150-fathom net length; depth was not standardized and drift times and methods vary among studies.

^f Defined as fish that could not be recovered thus died on-board a vessel. (Note that data for 3.5- and 4.5-in mesh includes 3:1 hang ratios, which appears to cause excessive tangling and increased mortality.)

^g Data from 2002 test fishery; defined as total mortality after 48 hours and includes immediate mortality.

^h Standard condition ranking scale. Data for 3.5-in mesh from 2001 permit fishery and 2001 test fishery and 4.5- and 5.5-in mesh from 2001 permit fishery and 2002 test fishery.

ⁱ Data from 2002 test fishery and general observations.

Table A-51. Estimated take of ESA-listed steelhead in Washington tributary recreational fisheries.

Affected Stock	Anticipated Encounters ^a	Expected Mortality ^b
Coweeman River winter steelhead	30%	1%
SF Toutle River winter steelhead	38%	2%
Cowlitz River winter steelhead	70%	4%
Kalama River winter steelhead	70%	4%
Mainstem/NF Lewis River winter steelhead	70%	4%
EF Lewis River winter steelhead	40%	2%
Washougal River winter steelhead	40%	2%
Wind River winter steelhead	30%	1%
Salmon Creek winter steelhead	30%	1%
Other tributaries winter steelhead	30%	1%

^a Anticipated encounters are catch and released fish; the numbers represent the percentage of fish from a stock anticipated to be incidentally encountered by anglers of a particular fishery.

^b Expected mortality is the hooking mortality of incidentally caught fish; expected mortalities are included in the anticipated encounters in terms of take.

A.4.10. Assessment of Current Status and Limiting Factors

Listing Status

NMFS BRT concluded that the Southwest Washington steelhead ESU (which includes Columbia River populations downstream of the Cowlitz River and Grays Harbor and Willapa Bay tributary stocks) is not currently in danger nor is it likely to become endangered in the foreseeable future (Busby et al. 1996). Therefore, the Grays, Elochoman, Skamokawa, Abernathy, Mill, and Germany populations are not listed under the ESA. However, the BRT decision reflects the overall condition of the entire ESU and does not necessarily reflect the condition of each lower Columbia population within the ESU. All of the Columbia River populations in the Southwest Washington ESU were categorized as depressed by WDFW in 2002, with the exception of Mill Creek, which was listed as unknown.

The BRT concluded that the Lower Columbia steelhead ESU (which includes steelhead from the Cowlitz River upstream to the Wind River) is not presently in danger of extinction, but is likely to become endangered in the near future. Therefore, on March 19, 1998, NMFS issued a formal notice listing the Lower Columbia steelhead ESU as threatened under ESA (Fed. Reg., V63, N53, p.13347). NMFS assessed the status of the Lower Columbia River steelhead ESU and reaffirmed the threatened listing on June 28, 2005 (Fed. Reg., V70, N123, p.37160). Subsequently, the current critical habitat was designated for this ESU on September 2, 2005 (Fed. Reg., V70, N123, p.52630).

WDFW categorized Kalama steelhead status as healthy, while Coweeman, NF Toutle/Green, SF Toutle, and Washougal steelhead were categorized as depressed, and NF Lewis, EF Lewis, Salmon Creek, Lower Gorge (Bonneville) Tributaries, and Wind River steelhead status were categorized as unknown.

The overall status of lower Columbia steelhead populations is generally low, but sustained natural production has been maintained in most areas in which steelhead were historically present. The most notable exceptions include areas in the Cowlitz and Lewis rivers where hydro development has blocked passage, and areas of the NF Toutle drainage where habitat was devastated by the eruption of Mt. St. Helens in 1980.

Current Viability

Steelhead are widely distributed throughout lower Columbia River basins, most of which continue to support significant populations. Declining long term trends and habitat reductions have placed this ESU at significant risk of extinction. However, many lower Columbia River steelhead are substantially healthier than their salmon counterparts.

Summer Steelhead

Five of six summer steelhead populations in the lower Columbia ESU occur in Washington. Of these, two are at very high risk of extinction, two are at moderate risk, and one is at low risk (Figure A-113). The sole Oregon summer steelhead population in the ESU has been found by Oregon to also at very high risks of extinction. Wind River summer steelhead are a rare lower Columbia example of a population with a high current level of viability. Current numbers of highly viable populations (Cascade = 0, Gorge = 1) and average viability scores (Cascade = 1.25, Gorge = 1.50) fall well below the standards established by the WLC-TRT for viability (2 populations per strata at or above 3.0, strata average viability of at least 2.25).

Extinction risks of summer steelhead reflect current population values of abundance, spatial structure, and diversity attributes. Abundance and productivity of naturally-spawning summer steelhead are very low for North Fork and East Fork Lewis population but high in the Kalama and very high in the Wind. However, aggregate abundance of lower Columbia summer steelhead is thought to have declined several fold from historical numbers. Total wild numbers currently average around 1,000-2,000 summer steelhead per year. Historical numbers were believed to be on the order of 10,000-30,000 per year. Numbers, as typified by Kalama weir counts, have generally declined over the last 20 years except for periodic spikes during favorable ocean survival conditions (Figure A-113).

Except in the North Fork Lewis where dams have blocked passage into the upper basin, spatial structure of summer steelhead remains consistent with very high levels of viability. Most populations continue to have access throughout historical production areas and much of this forested moderate to high elevation habitat remains relatively intact. Diversity of many summer steelhead populations has been reduced by historical hatchery effects, although not to the degree as has been seen in Chinook and coho salmon. Diversity ranges from very low in the North Fork Lewis to high in the Wind River.

Winter Steelhead

Washington lower Columbia winter steelhead include six populations at very high risk of extinction, six at high risk, and five at moderate risk (Figure A-114). No Washington populations are at low or very low risk of extinction. In contrast, of the seven additional populations occurring only in Oregon, six are at moderate, low, or very low risk. Populations in the unlisted coast strata (3 populations at very high viability, strata average of 3.0) meet standards established by the WLC-TRT for viability (2 populations per strata at or above 3.0, strata average viability of at least 2.25). In other strata, current numbers of highly viable populations (Cascade = 0, Gorge = 0) and average viability scores (Cascade = 1.0, Gorge = 1.3) do not currently meet the standard.

Extinction risks of Washington summer steelhead reflect current population values of abundance, spatial structure, and diversity attributes (Table A-52). Abundance and productivity of naturally-spawning summer steelhead are low or very low for most Cascade and grge populations. Total historical abundance of Columbia River winter steelhead was thought number in the tens of thousands. Current returns of wild fish to Wasington streams generally number less than 3,000 with a significant component of hatchery fish in many areas. Long-term escapement trend data is not available for most

wild winter steelhead populations but recent escapement estimates in the subset of monitored streams has varied between 1,100 and 2,700 steelhead per year (Figure A-96).

Except in the upper Cowlitz and North Fork Lewis systems where dams have blocked passage into the upper basin, spatial structure of winter steelhead generally remains consistent with high or very high levels of viability. Most populations continue to have access throughout historical production areas. Diversity of many winter steelhead populations has been somewhat reduced by historical hatchery effects, although not to the degree as has been seen in Chinook and coho salmon. Diversity ranges from moderate to very high in all systems.




Figure A-113. Distribution of historical summer steelhead populations among lower Columbia River subbasins. Extinction risks are based on viability scores rather than modeled risks.




Figure A-114. Current status of historical demographically-independent lower Columbia winter steelhead populations

Table A-52. Recovery goals for lower Columbia River steelhead populations (relative to listing baseline).

Population	Contribution	Baseline viability				Obj.	Prod. target	Abundance		
		A&P	S	D	Net			Historical	Baseline	Target
Coast Winter										
Grays/Chinook	Primary	VH	VH	M	M ¹	H	0% ⁴	1,600	800	800
Eloch/Skam	Contributing	VH	VH	M	M ¹	M+	0% ⁴	1,100	600	600
Mill/Ab/Germ	Primary	H	VH	M	M ¹	H	0% ⁴	900	500	500
Youngs Bay (OR)	Primary	-- ³	-- ³	-- ³	VH	VH	-- ³	-- ³	-- ³	-- ³
Big Creek (OR)	Primary	-- ³	-- ³	-- ³	H	VH	-- ³	-- ³	-- ³	-- ³
Clatskanie (OR)	Primary	-- ³	-- ³	-- ³	VH	VH	-- ³	-- ³	-- ³	-- ³
Scappoose (OR)	Primary	-- ³	-- ³	-- ³	VH	VH	-- ³	-- ³	-- ³	-- ³
Cascade Winter										
Lower Cowlitz	Contributing	L	M	M	L	M	+5%	1,400	350	400
Upper Cowlitz ^{C,G}	Primary	VL	M	M	VL ²	H ¹	>500%	1,400	<50	500
Cispus ^{C,G}	Primary	VL	M	M	VL ²	H ¹	>500%	1,500	<50	500
Tilton	Contributing	VL	M	M	VL	L	>500%	1,700	<50	200
S.F. Toutle	Primary	M	VH	H	M	H+	+35%	3,600	350	600
N.F. Toutle ^C	Primary	VL	H	H	VL ²	H	+125%	3,600	120	600
Coweeman	Primary	L	VH	VH	L ²	H	+25%	900	350	500
Kalama	Primary	L	VH	H	L ²	H+	+45%	800	300	600
N.F. Lewis ^C	Contributing	VL	M	M	VL ²	M	>500%	8,300	150	400
E.F. Lewis	Primary	M	VH	M	M ¹	H	+25%	900	350	500
Salmon	Stabilizing	VL	H	M	VL ²	VL	0%	na	<50	50
Washougal	Contributing	L	VH	M	L ²	M	+15%	800	300	350
Clackamas (OR) ^C	Contributing ⁵	-- ³	-- ³	-- ³	M	M+	-- ³	-- ³	-- ³	-- ³
Sandy (OR) ^C	Primary	-- ³	-- ³	-- ³	L	VH	-- ³	-- ³	-- ³	-- ³
Cascade Summer										
Kalama ^C	Primary	H	VH	M	M ¹	H	0% ⁴	1,000	500	500
N.F. Lewis	Stabilizing	VL	VL	VL	VL	VL	0%	na	150	150
E.F. Lewis ^G	Primary	VL	VH	M	VL ²	H	>500%	600	<50	500
Washougal ^{C,G}	Primary	M	VH	M	M ¹	H	+40%	2,200	400	500
Gorge Winter										
L. Gorge (WA/OR)	Primary	L	VH	M	L ²	H	+45%	na	200	300
U. Gorge (WA/OR)	Stabilizing	L	M	M	L ²	L	0%	na	200	200
Hood (OR) ^{C,G}	Primary	-- ³	-- ³	-- ³	M	H	-- ³	-- ³	-- ³	-- ³
Gorge Summer										
Wind ^C	Primary	VH	VH	H	H ¹	VH	0% ⁴	na	1,000	1,000
Hood (OR)	Contributing ⁵	-- ³	-- ³	-- ³	VL	M	-- ³	-- ³	-- ³	-- ³

¹ Increase relative to the interim plan.

² Reduction relative to the interim plan.

³ Addressed in the Oregon Management Unit plan.

⁴ Improvement increments are based on abundance and productivity, however, this population will require improvements in spatial structure or diversity to meet recovery objectives.

⁵ OR analysis indicates a low probability of meeting their delisting objective of high viability for this population.

^C Designated as a historical core population by the Technical Recovery Team.

^G Designated as a historical legacy population by the Technical Recovery Team.

Current Limiting Factors

Net Effects of Manageable Factors

Declines in status of lower Columbia River steelhead result from the combined impacts of human activities involving freshwater habitat, estuary habitat, dam construction and operation, fishing, fish hatcheries, and ecological factors such as predation. Impacts of each factor are compounded across the salmon life cycle to drive most populations to current very low levels. Net effects of quantifiable and potentially manageable impacts translate into an estimated 67-100% (average 84%) reduction in abundance and productivity of lower Columbia River steelhead populations in Washington (Figure A-115). Thus, current fish numbers represent only 0-33% of the historical production potential in the absence of potentially manageable impacts. Total reductions would be even greater if all human impacts could be effectively quantified. No single factor accounts for the majority of the reduction in fish numbers and the significance of specific factors varies from population to population (Figure A-115, Table A-53).




Figure A-115. Net effect and relative contribution of potentially manageable impact factors on steelhead in Washington lower Columbia River subbasins.

Table A-53. Productivity improvements and impact reduction benchmarks consistent with recovery of lower Washington Columbia River steelhead populations if reductions are distributed across factors in proportion to their baseline impact.

Population	Productivity	Baseline impacts (listing date reference)							Impacts at target					
	Target	Trib	Est	Dams	Harv	Hat	Pred	Δ	Trib	Est	Dams	Harv	Hat	Pred
<u>Coast Winter</u>														
Grays/Chinook	0%	0.50	0.15	0.00	0.10	0.08	0.20	0%	0.50	0.15	0.00	0.10	0.08	0.20
Eloch/Skam	0%	0.40	0.15	0.00	0.10	0.34	0.20	0%	0.40	0.15	0.00	0.10	0.34	0.20
Mill/Ab/Germ	0%	0.40	0.15	0.00	0.10	0.01	0.23	0%	0.40	0.15	0.00	0.10	0.01	0.23
<u>Cascade Winter</u>														
Lower Cowlitz	+5%	0.70	0.15	0.00	0.10	0.49	0.24	1%	0.69	0.15	0.00	0.10	0.48	0.23
Upper Cowlitz	>500%	0.40	0.15	1.00	0.10	0.49	0.24	50%	0.20	0.08	0.50	0.05	0.25	0.12
Cispus	>500%	0.60	0.15	1.00	0.10	0.49	0.24	50%	0.30	0.08	0.50	0.05	0.25	0.12
Tilton	>500%	0.90	0.15	1.00	0.10	0.49	0.24	50%	0.45	0.08	0.50	0.05	0.25	0.12
N.F. Toutle	+125%	0.80	0.15	0.00	0.10	0.33	0.24	20%	0.64	0.12	0.00	0.08	0.26	0.19
S.F. Toutle	+35%	0.80	0.15	0.00	0.10	0.24	0.24	7%	0.74	0.14	0.00	0.09	0.22	0.22
Coweeman	+25%	0.50	0.15	0.00	0.10	0.12	0.24	13%	0.43	0.13	0.00	0.09	0.10	0.20
Kalama	+45%	0.50	0.15	0.00	0.10	0.02	0.24	25%	0.37	0.11	0.00	0.07	0.02	0.18
N.F. Lewis	>500%	0.10	0.15	0.92	0.10	0.49	0.24	50%	0.05	0.08	0.46	0.05	0.25	0.12
E.F. Lewis	+25%	0.50	0.15	0.00	0.10	0.48	0.24	9%	0.45	0.14	0.00	0.09	0.44	0.22
Salmon	0%	0.80	0.15	0.00	0.10	0.50	0.24	0%	0.80	0.15	0.00	0.10	0.50	0.24
Washougal	+15%	0.50	0.15	0.00	0.10	0.08	0.24	8%	0.46	0.14	0.00	0.09	0.08	0.22
<u>Cascade Summer</u>														
Kalama	0%	0.43	0.15	0.00	0.10	0.01	0.24	0%	0.43	0.15	0.00	0.10	0.01	0.24
N.F. Lewis	0%	0.40	0.15	0.50	0.10	0.47	0.24	0%	0.40	0.15	0.50	0.10	0.47	0.24
E.F. Lewis	>500%	0.70	0.15	0.00	0.10	0.26	0.24	50%	0.35	0.08	0.00	0.05	0.13	0.12
Washougal	+40%	0.40	0.15	0.00	0.10	0.30	0.24	21%	0.32	0.12	0.00	0.08	0.24	0.19
<u>Gorge Winter</u>														
L. Gorge	+45%	0.60	0.15	0.00	0.10	0.01	0.22	20%	0.48	0.12	0.00	0.08	0.00	0.18
U. Gorge	0%	0.60	0.14	0.15	0.10	0.01	0.30	0%	0.60	0.14	0.15	0.10	0.01	0.30
<u>Gorge Summer</u>														
Wind	0%	0.50	0.14	0.15	0.17	0.01	0.273	0%	0.50	0.14	0.15	0.17	0.01	0.27

Fisheries

Current fishing impacts on steelhead are low and provide limited opportunity for increasing their numbers through additional fishery regulation. The primary fisheries targeting steelhead occur in the Columbia River mainstem and tributaries (Figure A-116); these fisheries harvest primarily hatchery fish and wild fish mortality is incidental. Fishing rates on wild steelhead have been reduced from their historical peaks in the 1960s by over 90% following prohibition of commercial steelhead harvest in the mainstem (1975), hatchery-only retention regulations in the mainstem starting in 1986, and hatchery-only retention regulations in the tributaries during the late 1980s and early 1990s. Interception of steelhead in ocean salmon fisheries is rare.




Figure A-116. Approximate steelhead fishery exploitation rates over time and allocation of current exploitation rates among fisheries.

Hatcheries

With recent widespread changes in hatchery practices and substantial timing differences between many hatchery and wild stocks, hatchery influence is currently moderate to low for most Washington lower Columbia steelhead populations (Table A-54). Most steelhead hatchery programs are intended to mitigate for the loss of natural steelhead production by providing fish for harvest opportunity. Many steelhead hatchery programs were developed from out-of-basin transfers or from multiple stocks; this practice continues today, primarily with Skamania summer and winter steelhead stocks released throughout the lower Columbia. In most cases, brood stock mixing was limited to a few stocks and performed only during the initial years of establishing the hatchery program. After the hatchery program had been established, brood stock collection came from returning adults, aside from minimal outside brood stock usage during years of hatchery shortfalls. Inter-specific hatchery predation impacts on steelhead are not an issue because wild rearing areas of small juvenile steelhead are primarily in areas upstream of hatchery release sites.

The indexed potential for negative impacts of hatchery spawners on wild population fitness of winter steelhead was estimated to range from 0 to 38%. Hatchery releases of winter steelhead currently range from 0 to 652,500 per subbasin. Hatchery fish continue to comprise 77-100% of the natural winter steelhead spawners in the lower Cowlitz and Lewis basins where large hatchery programs are operated to mitigate for lost access to upper basin spawning areas. Hatchery fractions on wild population spawning grounds during wild spawning periods are much lower in other subbasins, ranging from 0 to 23%. Reintroduction attempts in the upper Cowlitz basin rely entirely on hatchery stock that was originally derived from fish blocked at the dams. Current winter steelhead hatchery broodstock are derived from a variety of sources ranging from entirely natural fish (category 1) to highly domesticated stock (category 4). Hatchery fractions are generally low where broodstock of poor fitness are present whereas, more robust broodstock are present where hatchery fractions are high. In the Lewis and Cowlitz basins, the high incidence of hatchery spawners suggests that the fitness of natural and hatchery fish is now probably quite similar and natural populations could collapse without continued hatchery subsidy under current habitat conditions.

The indexed potential for negative impacts of hatchery spawners on wild summer steelhead population fitness was estimated to range from 4 to 65%. Hatchery releases of summer steelhead currently range from 0 to 225,000 per subbasin. Hatchery fish continue to comprise 93% of the natural summer steelhead spawners in the lower NF Lewis where a large hatchery program is operated to mitigate for lost access to upper basin spawning areas. Hatchery fractions on wild population spawning grounds during wild spawning periods are lower in other subbasins, ranging from 21 to 35%. Current winter steelhead hatchery broodstock are derived entirely from natural fish in the Kalama (category 1) but are highly domesticated elsewhere (category 4). In the NF Lewis, the high incidence of hatchery spawners suggests that the fitness of natural and hatchery fish is now probably quite similar and the natural population could collapse without continued hatchery subsidy under current habitat conditions.

The Columbia River Hatchery Scientific Review Group (HSRG) recently completed a review of hatchery programs affecting salmon and steelhead populations in the Lower Columbia River. The objective of the review was to change the focus of the Columbia River hatchery system from a harvest supplementation/hydropower mitigation focus to an ecosystem-based approach founded on the idea that harvest goals are sustainable only if they are compatible with conservation goals. Individual population reviews and recommendations can be found at http://www.hatcheryreform.us/mfs/reports/appendix/welcome_show.action.

Table A-54. Presumed reductions in wild population fitness as a result of natural hatchery spawners for Washington lower Columbia River steelhead populations.

Population	Annual releases ^a	Hatchery fraction	Fitness category	Assumed Fitness	Potential impact	Interacting releases	Interspecies impact
<u>Coast Winter</u>							
Grays/Chinook	40,000	0.05	4	0.3	0.038	0	0
Eloch/Skam	90,000 ^b	0.09	4	0.3	0.065	0	0
Mill/Aber/Germ	0 ^c	0.06	4	0.3	0.040	0	0
<u>Cascade Winter</u>							
Lower Cowlitz	652,500 ^d	0.92	2	0.7	0.276	0	0
Upper Cowlitz	287,500 ^f	1.00	2	0.7	0.300	0	0
Cispus	-- ^f	1.00	2	0.7	0.300	0	0
Coweeman	20,000	0.23	4	0.3	0.161	0	0
S.F. Toutle	0 ^e	0.02	2	0.7	0.006	0	0
N.F. Toutle	0 ^e	0	2	0.7	0	0	0
Tilton	100,000 ^g	1.00	2	0.7	0.300	0	0
Kalama	90,000 ^h	0	1	0.9	0	0	0
NF Lewis	100,000	0.77	2	0.7	0.231	0	0
EF Lewis	90,000	0	--	--	0	0	0
Salmon	20,000	na	na	na	na	0	0
Washougal	60,000	0	--	--	0	0	0
<u>Gorge Winter</u>							
L Gorge	0 ⁱ	0.01	4	0.3	0.007	0	0
U Gorge	0 ^j	0	--	--	0	0	0
<u>Cascade Summer</u>							
Kalama	90,000	0.35	1	0.9	0.035	0	0
N.F. Lewis	225,000	0.93	4	0.3	0.651	0	0
E.F. Lewis	25,000	0.27	4	0.3	0.189	0	0
Washougal	60,000	0.25	4	0.3	0.175	0	0
<u>Gorge Summer</u>							
Wind	0	0.21	4	0.3	0.147	0	0

^a Annual release goals.

^b The Elochoman River winter steelhead hatchery program at the Beaver Creek Hatchery stopped releasing smolts in 1999; hatchery returns were expected to significantly diminish starting with the 2001 return. The Elokomina Salmon Hatchery started a 'wild' winter steelhead program in 2000 to replace the previous program with indigenous stock (30,000 smolts per year). An additional 60,000 hatchery fish are released per year for fisheries. An additional 30,000 summer steelhead are released each year.

^c There are no steelhead hatchery programs in Mill, Abernathy, or Germany Creek. Sporadic small releases of winter steelhead have been made from the former Beaver Creek Hatchery program.

^d Includes 300,000 hatchery stock and 352,500 late winter stock. An additional 500,000 summer steelhead are released per year.

^e 25,000 summer steelhead are also released in each of the North and South Toutle.

^f Includes 37,500 yearlings and 250,000 subyearlings of late run stock intended to restore an upper Cowlitz basin population.

^g Fingerling releases for reintroduction purposes.

^h Includes 45,000 each of hatchery and late wild stocks. The winter steelhead program changed focus in 1998 1999; only wild steelhead are collected for brood stock.

ⁱ There are no hatchery steelhead programs in the lower gorge tributaries; winter steelhead from the Skamania and Beaver Creek Hatcheries were sporadically released in the basins since 1958.

^j The Wind River winter and summer steelhead hatchery programs at the Carson NFH stopped releasing smolts in 1997; hatchery returns were expected to significantly diminish starting with the 1999 return.

Stream Habitat

EDT analyses suggest that stream degradation has substantially reduced the habitat potential for steelhead in all Washington lower Columbia River subbasins where analyses have been completed (Figure A-117). Declines in habitat quantity and quality for steelhead have reduced current productivity potential to 6-34% and equilibrium numbers to 10-60% of the historical template. Substantial stream habitat improvements would be necessary to reach optimum conditions (i.e. PFC) for steelhead in remaining subbasin. Restoration of optimum habitat quality would be expected to increase habitat capacity by 30 to 2,400 adult steelhead per subbasin.

Steelhead rely on the middle mainstem to upper stream reaches where a lack of habitat diversity, sedimentation, and flow consistently limit habitat suitability. More detailed descriptions of stream habitat conditions and effects on fish in each subbasin may be found in Volume II of the Technical Foundation.




Figure A-117. Current, optimal, and historical subbasin productivity and capacity inferred for steelhead from stream reach habitat conditions using EDT.

Dams

Dam impacts on Washington lower Columbia steelhead were estimated to range from 0 to 100% (Figure A-118). Dams on the Cowlitz have inundated or blocked access to 100% of the winter steelhead habitat based on EDT assessments. In the North Fork Lewis, 95% of the winter steelhead habitat and approximately 50% of the summer steelhead habitat has been inundated or blocked. Passage mortality

at Bonneville Dam was assumed to average 10% for juveniles and an additional 5% for adults based on a synthesis of the available literature. Steelhead generally spawn and rear in headwater and upper mainstem reaches of subbasins and are less subject to hydropower effects on downstream habitats than are chum and fall Chinook.




Figure A-118. Assumed dam impacts on Washington lower Columbia steelhead populations.

Mainstem and Estuary Habitat

Mainstem and estuary habitat impacts were estimated to account for approximately a 10-20% reduction in productivity of winter and summer steelhead. Steelhead migrate through mainstem and estuary areas soon after emigration from tributary streams. Residence time in estuary and mainstem habitats is relatively brief, but smoltification and transition from fresh to salt water is a critical life stage.

Predation

Potentially manageable predation mortality was assumed to average 20% to 25% depending on travel distance from the subbasin to the ocean. Pikeminnow and tern management is projected to reduce salmonid predation by approximately 50%. Tern predation is almost entirely an artifact of recently established colonies on dredge spoil islands in the estuary but the current rate (9%) is less than half that observed prior to downstream translocation of the Rice Island colony (20%). Pikeminnow predation was greatest for populations originating in Bonneville Reservoir tributaries (5%), passing the pikeminnow gauntlet in Bonneville Dam forebay and tailrace, and traveling the entire 145 mile length from Bonneville to the Estuary. Predation rates by seals and sea lions on adult steelhead added an assumed 3% mortality.

Summary Assessment

1. Human activities including fishing, hatchery operation, alteration of stream, river, and estuary habitats, hydropower development and operation, and potentially manageable predation have collectively reduced productivity of winter and summer steelhead populations to 0-40% of historic levels. Recovery efforts will require significant improvements in multiple areas because no single factor accounts for the majority of the reduction in fish numbers.

2. Current fishing impacts on steelhead are relatively low and provide limited opportunities for increasing numbers through additional regulation of fisheries. Fishing impacts occur almost exclusively in Columbia basin sport fisheries. Selective fishery regulations were implemented for steelhead prior to listing.
3. Reduced productivity of wild populations as a result of interbreeding with potentially less-fit hatchery fish is among the most significant of hatchery concerns for wild stock recovery although these negative effects are at least partially offset by the demographic benefits of additional spawners. Potential negative impacts increase with the proportion of hatchery spawners and the disparity between wild and hatchery fish. Potential fitness impacts among Washington lower Columbia steelhead populations range from 0 to 65%. Potential impacts are greatest in the Cowlitz and Lewis basins where dams block most of the available steelhead habitat. Inter-specific hatchery predation impacts on steelhead are not an issue because wild rearing areas of small juvenile steelhead are primarily in areas upstream of hatchery release sites.
4. Stream habitat conditions significantly limit steelhead in all Washington lower Columbia River subbasins where EDT analyses have been completed. Substantial stream habitat improvements would be necessary to reach optimum conditions (i.e. PFC) in most subbasins. The significance of stream habitat suggests that recovery may not be feasible without substantial improvements in habitat quantity and quality.
5. Estuary and mainstem habitats are important to steelhead life history with assumed habitat impacts of 10-20%.
6. Hydropower development in the Cowlitz and Lewis have blocked 50-95% of the summer and winter steelhead habitat. Mainstem dam passage affects upper Gorge populations although passage success for steelhead tends to be greater than among other salmon species.



A.5. Bull Trout

Bull trout (*Salvelinus confluentus*) are a distinct species (Cavender 1978) that were previously considered to be a single species with Dolly Varden (*Salvelinus malma*) because of their overlapping ranges, similar appearance, and lack of sufficient analysis to discern the two species. Several genetic studies of the genus *Salvelinus* confirm the distinction between the bull trout and Dolly Varden (Phillips et al. 1989, Crane et al. 1994), and in fact show they are more closely related to other char species than to each other (Phillips et al. 1989, Phillips et al. 1991).

Bull trout were historically distributed as far south as the McCloud River in northern California, and Jarbidge River in northern Nevada, north to British Columbia, and east to Montana and Alberta. They were widely distributed throughout the Columbia Basin as far east as western Montana. Their current range in the lower 48 states is limited to Montana, Idaho, Oregon and Washington, with a small population in northern Nevada. They are now extinct in northern California.

Bull trout are listed as threatened under the ESA and are under the jurisdiction of the USFWS. Bull trout are subject of a draft recovery plan, although the USFWS recently decided to delay finishing the recovery plan in lieu of a 5-year review of the bull trout listing. The overarching goal of the bull trout recovery plan is to ensure the long-term persistence of self-sustaining, complex interacting groups (or multiple local populations that may have overlapping spawning and rearing areas) of bull trout distributed across the species' native range. In the lower Columbia, bull trout were believed to be historically distributed in the some large subbasins including the Lewis River and Columbia River upper Gorge tributaries. Of the subbasins addressed by this plan, bull trout currently occur only in the upper Lewis River. Bull trout were reported in the White Salmon River as recently as 1989 but have not been observed since despite focused sampling efforts. In the USFWS bull trout recovery plan, the Lewis, White Salmon, and Klickitat rivers have been identified as core bull trout habitats for the Lower Columbia Recovery Unit.

A.5.1. Life History and Requirements

Bull trout exhibit resident, freshwater migratory, and anadromous life history patterns (Rieman and McIntyre 1993; Figure A-119). Resident and migratory forms are known to coexist in the same subbasin or even in the same stream. While it is unknown whether resident and migratory forms of bull trout can produce progeny exhibiting the alternate life history behavior, multiple life history forms of other char are known to give rise to one another (Rieman and McIntyre 1993).

Resident forms live out their lives in the tributary where they were born and in nearby streams. Freshwater migratory forms include both fluvial and adfluvial strategies (Fraley and Shepard 1989). The

fluvial form migrates between main rivers and tributaries. The adfluvial form migrates between lakes and streams. Anadromous forms have been reported (WDFW 1997) in certain coastal areas, probably occur in the Puget Sound drainages and in the Squamish River, and may have occurred historically as far south as the Puyallup River (McPhail and Baxter 1996). Confirming the existence of anadromous bull trout populations is difficult because of the geographic overlap with Dolly Varden and the difficulty in discerning between the two species. In the lower Columbia River, bull trout may exhibit resident or freshwater migratory life history patterns; anadromous bull trout have not been observed.

Bull trout are found primarily in cold streams. Researchers consistently find that water temperature is a principal factor influencing distribution of bull trout in many streams (Rieman and McIntyre 1993, Baxter and McPhail 1996). Fraley and Shepard (1989) observed that water temperature above 59°F (15°C) may limit bull trout distribution. Studies in the John Day basin found bull trout present when maximum summer temperatures were 16°C or below, and maximum densities occurred where temperature maxima were 12°C or below (Buchanan and Gregory 1997). Bull trout do not compete well with introduced salmonids in degraded habitats (McPhail and Baxter 1996).

Bull Trout




Figure A-119. Bull trout life history.

Migration and Spawn Timing

Adults typically spawn from August to November during periods of decreasing water temperatures. Water temperature may be the proximate cue that initiates reproductive behavior. 48°F (9°C) appears to be the threshold temperature above which no spawning will occur (McPhail and Baxter 1996). Upstream migration begins in April and peaks during high flows in May and June (Pratt 1992). Spawners migrate upriver slowly, mostly at night, and enter tributary streams from late July through September (Pratt 1992). The above statements are broad generalities, and in fact migration patterns within and across basins can be varied and complex.

Spawning

Bull trout have spawning habitat requirements that may be more specific than those of other salmonids (Baxter and McPhail 1996). Spawners prefer areas of groundwater infiltration (Fraleley and Shepard 1989, Baxter and McPhail 1996). Redds are relatively large, typically measuring about 1.5m x 2m (59 x 78 in) (Fraleley and Shepard 1989, McPhail and Baxter 1996). Redd site selection across years may be remarkably consistent, and superimposition of redds has been observed (Baxter and McPhail 1999). Baxter and McPhail (1996) suggest that maintaining the quality of specific areas where high redd concentrations occur across years may be critical for survival in some populations. Buchanan et al. (1997) reported redd densities in bull trout spawning areas in the Umatilla and Walla Walla basins of 1.4- 8.6 redds/km, in study sections 5-16 miles (7.5-25.5 km) long.

Redd building and courtship behaviors occur mainly at night but have been observed during the day (McPhail and Baxter 1996). Often, only a single male is involved in mating, but jacks have been observed to surreptitiously mate with a female with which they have not courted (McPhail and Baxter 1996). It is likely, however, that reports of jacks may include cases of smaller, resident males mating with larger migratory females in those areas where the ranges of the two different life history forms overlap (McPhail and Baxter 1996). Because of different maturation rates and potential overlap of different life history patterns, it is possible to have four or more year classes compose any spawning population and as many as 12 to 16 age combinations in any spawning year (Shepard et al. 1984).

Preferred spawning habitats include stream reaches with loose clean gravel and cobble substrates, and temperatures 41-48°F (5-9°C) in late summer and early fall (Fraleley and Shepard 1989, Goetz 1989). Optimal water depths for spawning are 12-24 in (30-60 cm) (Boag 1991, Baxter and McPhail 1996). Bull trout spawning areas are generally higher in the watershed than other salmonid species.

Some studies suggest that optimal water velocity for spawning bull trout are 0.33-1.6 ft/sec (10-49 cm/sec), with velocities greater than 2.3 ft/sec (70 cm/sec) unsuitable for spawning (Boag 1991, Baxter and McPhail 1996).

Repeat and alternate year spawning have been reported. Frequency of repeat spawning is not well documented (Fraleley and Shepard 1989, Pratt 1992, Rieman and McIntyre 1996). Fraleley and Shepard (1989) reported that 38-69% (average 57%) of adult sized bull trout stayed in Flathead Lake each year. These were presumed to be fish that skipped a year of spawning. Mushens and Post (2000) found that an average of only 13% (range of 9-17%) of spawners in Smith-Dorien Creek had skipped a year of spawning. These are adfluvial bull trout from lower Kananaskis Lake, Alberta.

There are also limited observations of post-spawning mortality rates, although it is generally presumed to be relatively low. Mushens and Post (2000) reported spawning related mortalities in Smith-Dorrien Creek of 0.7-5.2% for 4 years of observation.

Incubation and Emergence

Incubating and emergent bull trout require colder water than other salmonid species. Cool water during early life history results in higher egg survival and fry growth rates (Pratt 1992, McPhail and Murray 1979, Shepard et al. 1984). McPhail and Murray (1979) found that bull trout fry grew to larger sizes at lower temperatures, and reached largest size at 39°F (4°C). Goetz (1989) suggested optimum water temperatures for bull trout incubation of about 35-39°F (2-4°C).

Eggs are about 0.197-0.236 in (5-6 mm) in diameter (McPhail and Baxter 1996). Egg numbers deposited in redds increase with body size of females, and as few as 74 eggs to as many 6,753 eggs have been documented (see McPhail and Baxter 1996 and sources cited therein). Incubation is 100–145 days (Pratt 1992); development rate is temperature-dependent, but is also related to egg size, especially at low temperatures (Murray 1980, McPhail and Baxter 1996). Bull trout require around 350-440 thermal units (TUs) after fertilization to hatch (Weaver and White 1984, Gould 1987). McPhail and Murray (1979) investigated the relationship between egg survival and water temperature. They reported that at water temperatures of 46-50°F (8-10°C), 0-20% of eggs survived to hatching. At 42°F (6°C), 60-90% survived, and at 35-39°F (2-4°C), 80-95% survived.

The two major causes of egg mortality are siltation and freezing (McPhail and Baxter 1996). Weaver and White (1985) reported a negative relationship between intergravel fines and incubation survival in laboratory tests. Approximately 40%, 20%, and 1% of fertilized eggs survived to hatch when spawning substrate consisted of 20%, 30%, and 40% fines, respectively (fines defined as <0.37 in [9.5 mm] in diameter).

After hatching, juveniles remain in the substrate for up to 3 weeks before emerging from the gravel, and emergence may take place up to 200 days after the eggs have been deposited. Emergence is normally April–May, depending on water temperature and flow patterns (Pratt 1992, Ratliff and Howell 1992). Fraley and Shepard (1989) estimated that 50% of eggs survive to emergence of fry (Pratt 1992, Gould 1987). Size at emergence is usually around 1.0-1.1 in (25-28 mm).

Freshwater Rearing

In laboratory experiments, newly emerged fry did not fill their swim bladder for 3 weeks after emergence, were strongly bottom-oriented, and spent a great deal of time in the small spaces between pieces of gravel at the bottom of the water (McPhail and Baxter 1996).

Juvenile bull trout are associated with complex cover, including large wood, undercut banks, boulders, and pools (Fraley and Shepard 1989). In general, juvenile bull trout are associated with shallow water depths with good cover, near faster-flowing water that delivers food particles (Baxter and McPhail 1996). Fry stay close to the streambed; McPhail and Murray (1979) suggest this might be an adaptation to avoid being carried downstream before the fry are large enough to take up residence in a suitable feeding site. Mean distance above the stream bed increases as the fish get larger (Pratt 1984), although they tend to remain in the bottom 25% of the water column (McPhail and Baxter 1996). Age 0+ fish tend to hold in water depths of 0.1-1.5 in (2-40 mm) (Baxter and McPhail 1996, Baxter 1995, Tredger 1979, Ptolemy et al. 1977).

Fry frequently inhabit stream margins and side channels (Sexauer and James 1997). Martin et al. (1992) reported that age 0 bull trout densities in Mill Creek, Oregon, were highest in riffle- and cascade-type habitats and in the presence of woody debris. Goetz (1997) reported that age 0 bull trout were active at day while age 1 and older were most active during twilight. Paul (2000) found that age 0 bull trout were absent during their nighttime sampling in Smith-Dorrien Creek, Alberta.

Young bull trout exploit small pockets of slow water near higher velocity, food-bearing water (Shepard et al. 1984, Pratt 1992). These microhabitats may be created by cobble substrate, and Shepard et al.

(1984) found highest densities of juvenile bull trout in reaches with highest cobble substrate percentages in the Flathead River basin. Densities decline as the small spaces between gravel are filled with fine particles (Enk 1985). Where unembedded cobble substrate is not available, woody debris, turbulence, and undercut banks take on increasingly important roles in providing suitable habitat (Pratt 1984). In general, complex forms of cover and high stream complexity are favored by young bull trout (Baxter and McPhail 1996).

Significant shifts in habitat use from one day to the next have been observed (Goetz 1994, Sexauer 1994, Baxter and McPhail 1996). At night, most bull trout juveniles observed are not associated with cover.

Juvenile bull trout < 4.3 in (110 mm) most commonly consume aquatic insects and fish. Once they are approximately 4.3 in (110 mm) long, they may begin feeding on smaller fish (Pratt 1992), and may consume prey items that are large in relation to their own body size. McPhail and Baxter (1996) report observing a 1.8 in (45 mm) rainbow trout in the stomach contents of a 3.5 in (90 mm) bull trout. Cannibalistic behavior is common. Fish species identified in the stomachs of juvenile bull trout include mountain whitefish, sculpins, salmon fry, and trout, including other bull trout (McPhail and Baxter 1996). Boag (1987) reports increased piscivory in adfluvial populations, as larger fish prey more exclusively on fish as they move downstream into larger water.

Juvenile and adult bull trout densities are typically low, and the species may be more sensitive to environmental degradation than other salmonids (McPhail and Baxter 1996). Late summer densities of bull trout ages 1-3 in pool and glide habitats in Jack, Roaring, Brush, Canyon and Candle creeks tributary of the Metolius River, Oregon, were estimated to range from 2.0 to 20.6 fish per 100 square meters (Buchanan et al. 1997). In Mill Creek, juvenile bull trout densities were highest in plunge pools with woody debris (8.7 fish per 100 m²) and run habitat with woody debris (8.4 fish per 100 m²) (Martin et al. 1992). Fraley and Shepard (1989) and McPhail and Murray (1979) found juvenile bull trout densities to be higher in pools than in other habitat unit types. Carrying capacity of streams for juvenile bull trout is thought to be the major bottleneck in production (McPhail and Murray 1979, McPhail and Baxter 1996).

Paul (2000) found that densities of fluvial bull trout juveniles in Eunice Creek, Alberta over a period of 15 years fluctuated over two orders of magnitude (0.06 fish/100 m² to 19.57 fish/100m²). During the same time, rainbow varied only from 0.45 fish/100m² to 3.29 fish/100m². Few fish were over 9.8 in (250 mm). The study also investigated interactions of juvenile bull trout and their role in population dynamics. The author used the sum of fork length squared as a measure of effective density (Walters and Post 1993, Post et al. 1999), because consumption rate is an exponential function of body size, with fish consuming less per body weight as they increase in size. He found that survival of age 1 and 2 juveniles was highly correlated to effective density (FL²/m²). The model was significantly improved by adding mean discharge (April-October), with a positive effect. Survival of age 3 bull trout was best correlated to density of age 3 bull trout. Survival of age 2 bull trout was positively related to summer flow. Density effects reduce survival of older juveniles up to 60%.

Paul (2000) also found significant negative relationship of juvenile bull trout growth to effective density in a resident bull trout population in Prairie Creek, Alberta. Growth rate was depressed more by abundance of larger juveniles than by abundance of smaller juveniles. The author concluded that competition for food occurred among all juveniles age 1 or older. This differs from other salmonids that tend to partition food between age groups as the fish exploit different habitat zones.

Ratliff et al. (1996) found no apparent relationship between redd counts and densities of age 1-3 juveniles in five spawning tributaries of the Metolius River, suggesting density dependence mechanisms affect juvenile abundance. Paul (2000) used experiments with stream enclosures to demonstrate that growth of age 1 bull trout was density-dependent. Over 42 days of the experiment, there were no differences in survival among treatments, but highly significant (P<0.01) differences in growth. Further,

he found positive relationship between fish size at the beginning of experiment and survival to the end of experiment. Survival averaged 52% over 42 days for 4.33 in (110 mm) bull trout. He concluded that overwinter survival rate was determined by growth rate, which was determined by effective density. Larger fish at age survived better than smaller ones, and dominant age classes could be produced if larger fish were depleted, allowing rapid growth of age 0 fish and resulting in high overwinter survival. Survival rates through winter were based on bioenergetic modeling of lipid stores, which increase with body size. Thus, larger fish with more body fat withstand longer periods of starvation.

Juvenile Migration

Migratory juvenile bull trout typically rear in their natal streams for 2 to 3 years before migrating downstream. Although juveniles migrate in all months, most migration peaks in May and June. Migrating juveniles average about 8 in (200 mm) long. Juvenile bull trout migrate from the streams in which they are born to larger rivers and lakes throughout their range at ages 1, 2, and 3 (Pratt 1992).

Estuary Rearing/Ocean Migrations

Although bull trout are known to exhibit anadromous life history patterns, anadromy has not been observed in lower Columbia River bull trout. Thus, bull trout in the lower Columbia River do not have an estuarine rearing or ocean migration phase as part of their life cycle (Figure A-119).

Adults

Size and age at maturity vary depending if the fish is resident, freshwater migratory, or anadromous. At maturity, resident fish are generally smaller and less fecund than migratory fish (Fraleigh and Shepard 1989). Bull trout normally reach sexual maturity from age 4–7, and may live longer than 12 years.

Because bull trout can be resident, freshwater migratory, or anadromous, it is not a simple task to come up with generalized habitat requirements for adult bull trout, but some themes are common. Bull trout are a cold water species. Bull trout are seldom found in areas where water temperatures frequently exceed 59°F (15°C). Forms of cover favored by adult bull trout include deep pools. Usually, adult fish migrate into a stream during spring or early summer freshets and may reside in deep pools up to 2 months before spawning (Baxter and McPhail 1996). This tendency makes adult bull trout particularly vulnerable to poaching or overfishing (McPhail and Baxter 1996).

Resident populations of bull trout usually are separated from other populations by some physical or thermal barrier (McPhail and Baxter 1996). Where present, resident populations are typically found in headwater streams in mountainous areas, and in higher gradients than other forms. They are usually associated with deep pools and complex cover, and are much smaller than individuals that migrate into larger rivers during adulthood. Resident fish average about 7.87 in (200 mm) in length, and mature from 1 to 2 years earlier than migratory fish in the same geographic area (McPhail and Baxter 1996). Suitable overwinter sites are critical to the viability of stream resident populations. Research has found that the most suitable overwinter habitats are areas of groundwater upwelling and deep pools (McPhail and Baxter 1996).

Fluvial forms of bull trout live as adults in large rivers but return to small tributary streams to spawn. Fluvial individuals usually reach sexual maturity by age 5, and can attain large sizes. Individuals up to 35 in (900 mm) have been reported (Baxter 1995, McPhail and Baxter 1996). In many instances, fluvial bull trout densities in larger rivers are higher near the mouths of smaller spawning tributaries that deliver colder water to the system (Buckman et al. 1992). Fluvial adults are generally associated with deep pools and instream cover (Shepard et al. 1984). In some systems where water is more turbid, adult bull

trout are less associated with cover and more widely distributed, perhaps reflecting their status as top predators (Bishop 1975, McPhail and Baxter 1996).

Adfluvial bull trout live as adults in lakes and return to small tributaries to spawn. In some cases, spawning may occur in the lake outlet. Spawning migrations may be quite short, or as long as 124 miles (200 km) (Fraley and Shepard 1989). Evidently, lake dwelling adult bull trout may use different parts of lakes at different times of the year. In Flathead Lake, Montana, Goetz (1989) reported that bull trout forage in the littoral zone in fall and spring and move to deeper water in summer, likely because of temperature considerations. Sexual maturity is usually reached by age 5, and individuals can attain large sizes (up to 27.5 in [700 mm] long; McPhail and Murray 1979).

Beauchamp and van Tassell (1999) conducted a thorough diet study of an adfluvial bull trout population in Round Butte Reservoir on the Deschutes River. Kokanee, bull trout, rainbow trout, mountain whitefish, other salmonids, nonsalmonid fishes, and invertebrates were all important in adult bull trout diets, with diets changing seasonally and by size class of bull trout. Small bull trout (FL < 300mm) consumed primarily age 0 mountain whitefish in all seasons, as well as age 0 kokanee during summer; length of consumed prey items increased in a seasonal progression. Subadult bull trout (FL 300-450mm) ate age 0 kokanee during summer and fall and transitioned to age 1 kokanee during the winter and spring; intermediate sized mountain whitefish (~150mm) were an important food item during summer. Adult bull trout (\geq 450 mm) ate age 0 and age 1 kokanee from winter through summer but shifted to age 2 and age 3 kokanee during the fall; mountain whitefish were also an important prey item during the winter and spring. Fraley and Shepard (1989) sampled bull trout stomachs in Flathead Lake during November and January and found that kokanee composed 8.9% of the diet by weight, while various species of whitefish composed 48.1% and non-game fish composed 22.1%.

Beauchamp and Tassell (1999) used bioenergetics modeling to estimate the total predation impact by age 3 to 7 bull trout in Round Butte Reservoir. They estimated that the adult bull trout population annually consumed 11-49% of the available age 0 bull trout, 4-18% of the age 1 bull trout, 5-11% of the age 0 kokanee, 1-2% of the age 1 kokanee, and 13-74% of the age 2-3 kokanee. Larger bull trout ate larger prey, resulting in adults contributing the most to kokanee predation losses while subadults contributed the most to juvenile bull trout cannibalism losses.

A.5.2. Distribution

The Columbia River basin supports a total of 141 subpopulations of bull trout, 20 located in the lower Columbia River region downstream of the Klickitat River (Figure A-120). Of these 20 subpopulations, two are located in the Lewis River (Federal Register, Vol. 63, No. 111, June 10, 1998). Bull trout have never been reported in the Wind River above Shiphed Falls (RM 2.0) (Byrne et al. 2001). Bull trout have been reported in the Little White Salmon basin but never above Little White Salmon National Fish Hatchery. Byrne et al. (2001) conclude after sampling in the subbasin that bull trout are not present in Lava Creek of the mainstem Little White Salmon above the hatchery, but could not confirm absence of bull trout in Moss Creek due to equipment failure and extensive instream debris. Reports of White Salmon River bull trout are rare, and it is unclear where preferred spawning areas are located. It is doubtful that bull trout in the White Salmon system venture into the mainstem Columbia due to temperature considerations (WDFW 1998). Byrne et al. (2001) note that groundwater contributions in the canyon area of the White Salmon River and in Spring Creek could make these areas possible bull trout habitat. Bull trout populations were also suspected to historically inhabit the Cowlitz and Kalama subbasins, but the current distribution of bull trout in these subbasins is unknown.




Figure A-120. Distribution of historical bull trout populations among lower Columbia River subbasins.

A.5.3. Genetic Diversity

Genetic variability within populations is low, but genetic differences among populations are often marked. This suggests that many small populations have undergone genetic bottlenecks (McPhail and Baxter 1996). Genetic samples were taken from bull trout captured in Lake Merwin, Yale Lake, and Swift Reservoir in 1995 and 1996. Analysis showed that Lewis River basin bull trout were genetically similar to the Columbia River population (Spruell et al. 1998). Spruell et al. (2003) conducted microsatellite analysis and concluded that 'coastal' bull trout (west of the John Day River) were genetically distinguished from Snake River and Upper Columbia groups. Within the coastal population, however, some genetic variation was observed, primarily between drainages. Spruell et al. (1998) found that the Swift population was found to be significantly different from that in Yale Lake and Lake Merwin. This implies that there may have been biological separation of the upper and lower basin stocks prior to completion of Swift Dam in 1958.

A.5.4. ESU Definition

Because of widespread distribution, isolated populations, and variations in life history, bull trout populations are grouped by distinct population segments (DPS) rather than ESU. Bull trout are also grouped by recovery units, which serve as subsets of the distinct population segments. By examining distinct population segments, bull trout in most need of Federal protection become a listing priority. On June 10, 1998, the USFWS issued a final rule announcing the listing of bull trout in the Columbia and Klamath river basins as threatened under the ESA (Federal Register, Vol. 634, No. 111).

Within the Columbia River Basin distinct population segment of bull trout, the Lower Columbia River Recovery Unit includes the Lewis River and Klickitat River core areas in Washington. The Lewis River Core Area consists of the mainstem Lewis River and tributaries downstream to the confluence with the Columbia River, with the exclusion of the East Fork of the Lewis River. The Klickitat River Core Area includes the Klickitat River and all tributaries downstream to the confluence with the Columbia River.

A.5.5. Life History Diversity

Bull trout exhibit resident, freshwater migratory, and anadromous life history patterns (Rieman and McIntyre 1993). Resident and migratory forms are known to coexist in the same subbasin or even in the same stream. While it is unknown whether resident and migratory forms of bull trout can produce progeny exhibiting the alternate life history behavior, multiple life history forms of other char are known to give rise to one another (Rieman and McIntyre 1993).

In the lower Columbia River, bull trout may exhibit resident or freshwater migratory life history patterns (Table A-55); anadromous bull trout have not been observed. Confirming the existence of anadromous bull trout populations is difficult because of the geographic overlap with Dolly Varden and the difficulty in discerning between the two species.

Table A-55. Life history and population characteristics of bull trout originating in Washington portions of the lower Columbia.

Characteristic	Life History Form	
	Migratory	Resident
Number of extant populations	20 subpopulations	
Upstream spawning migration	April – September	April – September
Spawn timing	Early fall	Early fall
Spawning habitat type	Runs and tail-outs	Runs and tail-outs
Emergence timing	January – May	January – May
Natal area rearing	1-3 years	5-7 years
Downstream migration of juveniles	April - November	NA
Rearing habitat	Lake or large river	Headwater streams, higher gradient
Lake/river residence	2-6 years	NA
Age at spawning	4-12 years with annual or intermittent spawning ~10-40 in Cougar Creek, Yale Reservoir, Lewis River (1988-2003)	4-12 years with annual or intermittent spawning
Natural spawners	~100-900 in Rush/Pine Creeks, Swift Reservoir, Lewis River (1994-2003)	Unknown
Hatchery adults	None	None

A.5.6. Abundance

Status of bull trout is difficult to ascertain because of the lack of commercial harvest, hatchery production, and scarcity of data. The Lewis River bull trout population was classified as depressed because of chronically low numbers (WDFW 1998). Adfluvial populations exist in Yale and Swift reservoirs in the Lewis River system. No fish passage is in place at the dams impounding these

reservoirs; bull trout are thought to move downstream during spill events. Swift Reservoir bull trout spawn in Rush and Pine creeks. Cougar Creek is the only known spawning location for bull trout in Yale Reservoir; however, there may be potential for spawning in Ole Creek if flow is augmented. Bull trout in Merwin Reservoir are thought to be present due to spill from Yale Reservoir; however, there is no spawning population in Merwin Reservoir (WDFW 1998). WDFW and PacifiCorp have engaged in a program to relocate bull trout from the Yale tailrace back to Yale Reservoir (Table A-56).

Table A-56. Bull trout collected from the Yale tailrace (Lake Merwin) and transferred to the mouth of Cougar Creek (Yale Reservoir) or released back into Yale Reservoir (1995–2000).

Year	No. Collected in Yale Tailrace	No. Transferred to Mouth of Cougar Creek	No. Released Back into Yale Reservoir
1995	15	9	6
1996	15	13	2
1997	10	10	0
1998	6	6	0
1999	6	0	6
2000	7	7	0
Total	59	45	14

* not including recaptures

Historical information describing the abundance and distribution of bull trout in the Lewis River basin is limited. However, the number of bull trout spawners utilizing Cougar Creek has been documented annually since 1979. During this period, the number of adult spawners in Cougar Creek (based on annual peak counts) has ranged from 40 in 1979 to 0 in 1981 and 1982 (Figure A-121). The low number of spawners observed in the early 1980s may be related to impacts associated with the 1980 eruption of Mt. St. Helens.




Figure A-121. Annual peak counts of bull trout spawners observed in Cougar Creek, 1979–2003.

In addition to the survey work conducted in Cougar Creek, the U.S. Forest Service (USFS), WDFW, and PacifiCorp have collected distribution and abundance information on bull trout since the late 1980s. Bull trout collected at the head of Swift Reservoir have been marked with Floy (anchor) tags every spring since 1989 to facilitate mark and recapture counts in Rush and Pine creeks (i.e. the primary spawning tributaries for the Swift bull trout population; Lesko 2001). Between 1994 and 2003, the annual spawner population in Swift Reservoir has ranged from 101 to 911 fish (Figure A-122; Lesko 2001; personal communication, Dan Rawding and J. Weinheimer, WDFW, 2000).




Figure A-122. Spawning population estimate of bull trout in Swift Reservoir, 1994–2003 (source: Dan Rawding and John Weinheimer, WDFW).

A.5.7. Productivity

The two Lewis basin bull trout populations appear to maintain low but fairly stable production levels in the limited habitat available. The Yale Reservoir production is less certain on an annual basis because of dependence on only one known stream for spawning; thus, a catastrophic event to Cougar Creek would significantly change the productivity of Yale Reservoir bull trout. Additionally, the Yale Reservoir production could be reduced if a significant number are entrained at Yale Dam and displaced to Merwin Reservoir. The Swift bull trout production appears to be more stable, with both Pine and Rush Creek supplying spawning and rearing habitat.

A.5.8. Fishery

Sport fishing for bull trout was eliminated in the Lewis and White Salmon drainages in 1992. Hooking mortality may occur from catch and release of bull trout in fisheries targeting other fish, particularly the coho and kokanee fisheries in Merwin and Yale reservoirs (WDFW 1998). Incidental catch of bull trout is thought to be low, however. In the Lewis River system, incidental take of bull trout is thought to be higher above Swift Reservoir (WDFW 1998). WDFW has actively set fishery regulations to protect bull trout in reservoirs and tributaries in the Lewis River basin.

A.5.9. Assessment of Current Status and Limiting Factors

Listing Status

According to WDFW (1998), the bull trout populations in the Lewis River basin are considered at moderate risk of extinction. The bull trout in the coterminous United States was listed as threatened under the ESA on November 1, 1999 (64 FR 58910). Earlier rulemakings had listed distinct population segments of bull trout as threatened in the Columbia River, Klamath River, and Jarbidge River basins (63 FR 31647, 63 FR 42757, 64 FR 17110).

For listing purposes the range of bull trout was broken into distinct population segments. Bull trout occur in widespread, but fragmented habitats and have several life history patterns. In addition, threats are diverse and the population status and trends vary considerably throughout the range. By examining distinct population segments, bull trout in most need of Federal protection become a listing priority. Many of the actions intended to protect other declining salmonids may also help bull trout. Stream and habitat protection and restoration, reduction of siltation from roads and other erosion sites, and modification of land management practices to improve water quality and temperature are all important.

The Bull Trout Recovery Team has developed a draft recovery plan providing a framework for implementing recovery actions in the coterminous United States. Because bull trout are widely distributed over a large area and have differing threats, the U.S. Fish and Wildlife Service identified 27 recovery units based on large river basins and generally following existing boundaries of conservation units for other fish species described in state plans, where possible. Each recovery unit has its own individualized recovery strategy.

Bull trout in the Lower Columbia River Recovery Unit are included in the Columbia River Basin distinct population segment of bull trout. In the two core areas, local populations of bull trout exist in the Cougar, Pine, and Rush creeks (tributaries of the Lewis River) and the West Fork of the Klickitat River. No local populations have been identified in the White Salmon River, but that area contains core habitat and after migratory obstructions are addressed, could support bull trout that migrate from the Columbia River. Additional research is needed to determine if the Cowlitz and Kalama rivers are important for bull trout recovery.

Current Viability

Bull trout were federally listed as threatened in 1999. The USFWS has formulated a draft recovery plan, and identified 27 recovery units for bull trout. One of these is the Lower Columbia recovery unit, which has two core areas (the Lewis River and the Klickitat River). While no local populations have been identified within the White Salmon, the subbasin contains core habitat, and could support bull trout (USFWS 2002). Recent natural escapements in two upper Lewis River spawning areas currently average several hundred fish per year. The size of the Gorge population is unknown.

Recovery Planning Ranges

At present, recovery standards for bull trout have only been partially identified. However, USFWS has compiled a list of research criteria to gather the data necessary to assess whether management actions are resulting in the recovery of bull trout in the Lower Columbia recovery unit. USFWS (2002) identified the following recovery standards and research needs:

1. Distribution of bull trout in the Lower Columbia recovery unit is unknown and considered a research need. Until additional information is obtained, at a minimum, the existing local populations in the recovery unit need to be maintained.
 - a. USFWS (2002) states that “establishment of additional local populations . . . is essential for recovery.” Potential sites which have or could support bull trout if restored should be evaluated for possible reintroduction.
 - b. Factors that may limit potential for reintroduction should be identified
2. Estimated abundance of bull trout in the Lower Columbia recovery unit local populations is considered a research need.
 - a. A complete set of data is not available from which to make a reliable estimate of bull trout abundances in any of the local populations.
 - b. As more data is collected, population estimates will be conducted to more accurately reflect both migratory and resident life history forms.
3. Adult bull trout exhibit a stable or increasing trend for at least two generations at or above the identified abundance level (from criteria 2) within core areas.
 - a. The development of a standardized monitoring and evaluation program to accurately describe trends in bull trout abundance has been identified as a priority research need.
4. Barriers to bull trout migration in the Lower Columbia recovery unit need to be addressed.
 - a. Barriers that have been identified as primary impediments to recovery, and where connectivity must be reestablished are Swift 1 and 2 and Yale Dams on the Lewis River, and Condit Dam on the White Salmon River.

Summary Assessment

1. The historic distribution and abundance of bull trout in the lower Columbia region are unknown. Bull trout are known to exist in the Lewis drainage and some Gorge tributaries.
2. Hydropower development has negatively affected bull trout populations in the Lewis River system, where three hydroelectric dams block fish passage and eliminate connectivity of subpopulations.
3. The USFWS has recommended installing a means of fish passage at Condit Dam on the White Salmon River, although no bull trout are known to occupy that system now. Suitable habitat exists, and bull trout are believed to have existed in the White Salmon historically.
4. Fishing for bull trout is closed in Washington. Bycatch has been reported in the Lewis River watershed kokanee fishery but its impacts are believed to be very low.
5. There are no hatchery programs to produce bull trout. Interactions between bull trout and hatchery-produced salmonids have not been studied, and impacts are unknown.

A.6. References

- Abbe, T.B. and D.R. Montgomery. 1996. Large woody debris jams, channel hydraulics and habitat formation in large rivers. *Regulated Rivers: Research and Management* 12:201-221.
- Adams, T.N. and K. Sullivan. 1989. The physics of forest stream heating: A simple model. Weyerhaeuser Technical Report. 044-5002/89/1. Technology Center, Tacoma, Washington.
- Alderdice, D.F., and F.P.J. Velsen. 1978. Relation between temperature and incubation time for eggs of chinook salmon (*Oncorhynchus tshawytscha*). *Journal of the Fisheries Research Board of Canada* 35:69-75
- Allen, J.H. 1980. Life history notes on the Dolly Varden char (*Salvelinus malma*) in the upper Clearwater River, Alberta. Alberta Energy and Natural Resources, Fish and Wildlife Division, Red Derr, Alberta.
- Allen, K.R. 1969. Limitations on production in salmonids populations in streams, Pages 3-18. in T.G. Northcote, ed. *Symposium on Salmon and Trout in Streams*. H.R. MacMillan Lectures in Fisheries. Institute of Fisheries, University of British Columbia, Vancouver.
- Allendorf, F.W. 1975. Genetic variability in a species possessing extensive gene duplication: genetic interpretation of duplicate loci and examination in genetic variation in populations of rainbow trout. Doctoral dissertation. University of Washington, Seattle.
- Allendorf, F.W. and N. Ryman. 1987. Genetic management of hatchery stocks. Pages 141-159 in N. Ryman and F. Utter eds. *Population genetics and fishery management*. University of Washington Press, Seattle.
- Allendorf, F.W., and R.S. Waples. 1995. Conservation and genetics of salmonid fishes. Pages 238-281 In J.C. Avise and J.L. Hamrick, editors. *Conservation genetics: Case histories from nature*. Chapman and Hall, New York.
- Allendorf, F.W., et al. 1997. Prioritizing Pacific salmon stocks for conservation. *Conservation Biology* 11: 140-152.
- Altukhov, Y.P., and E.A. Salmenkova. 1994. Straying intensity and genetic differentiation in salmon populations. *Aquaculture and Fisheries Management* 25 (Suppl. 2):99-120.
- Anders, P., and R. Westerhof. 1996a. Conservation Aquaculture of Endangered White Sturgeon (*Acipenser transmontanus*) from the Kootenai River, Idaho. Pages 51-62 in *Proceedings of the International Congress on the Biology of Fishes*, San Francisco State University, California. July 14-18, 1996.
- Anders, P., and R. Westerhof. 1996b. Natural spawning of white sturgeon (*Acipenser transmontanus*) in the Kootenai River, 1995. Preliminary Report of Research FY95. Kootenai Tribe of Idaho and Bonneville Power Administration. Portland, Oregon. 14 pp.
- Anders, P.J. 1991. White sturgeon (*Acipenser transmontanus*) movement patterns and habitat use in the Kootenai River system, Idaho, Montana, and British Columbia. Master's thesis, Eastern Washington University, Cheney, Washington. 153 pp.
- Anders, P.J. 1994. Kootenai River Fisheries Studies. Annual Progress Report FY93. Report A: Natural spawning of white sturgeon in the Kootenai River. Kootenai Tribe of Idaho. Prepared for U.S. Department of Energy, Bonneville Power Administration. Project No. 88-64. Portland, Oregon.

- Anders, P.J. 1996. Kootenai River Fisheries Studies. Annual Progress Report FY93. Report A: Natural spawning of white sturgeon in the Kootenai River. Kootenai Tribe of Idaho. Prepared for U.S. Department of Energy, Bonneville Power Administration. Project No. 88-64. Portland, Oregon.
- Anders, P.J. 2000. Ancient fish need modern protection. *Fisheries* 25 (9):30.
- Anders, P.J. and M.S. Powell 2002. Geographic and Frequency Distributions of Control Region Length Variation in the mtDNA Genome of White Sturgeon (*Acipenser transmontanus*) from the Columbia River Basin. Chapter 2 in: Anders, P.J. Conservation Biology of White Sturgeon. Doctoral dissertation, University of Idaho, Aquaculture Research Institute, Center for Salmonid and Freshwater Species at Risk. 221 pp.
- Anders, P.J., and L.G. Beckman. 1995. Comparison of white sturgeon egg mortality and juvenile deformity among four areas of the Columbia River. Report H in Status and Habitat Requirements of the White Sturgeon Populations in the Columbia River Downstream from McNary Dam. Project No. 86-50. Final Report of Research to the Bonneville Power Administration, U.S. Department of Energy, Portland, Oregon. Volume 2.
- Anders, P.J., C. Gelok, and M.S. Powell. 2002. Population structure and mitochondrial DNA diversity of North American white sturgeon (*Acipenser transmontanus*): An empirical expansive gene flow model Chapter 3 in: Anders, P.J. Conservation Biology of White Sturgeon. Doctoral dissertation, University of Idaho, Aquaculture Research Institute, Center for Salmonid and Freshwater Species at Risk. 221 pp.
- Anders, P.J., D.L. Richards, M.S. Powell. 2002. The First Endangered White Sturgeon Population (*Acipenser transmontanus*): Repercussions in an Altered Large River-floodplain Ecosystem. Pages 67-82 in W. VanWinkle, P. Anders, D. Dixon, and D. Secor, eds. Biology, Management and Protection of North American Sturgeons. American Fisheries Society Symposium 28.
- Anderson, J.J. 2000. Decadal climate cycles and declining Columbia River salmon. In Sustainable Fisheries Management: Pacific Salmon. E. Knudsen et al., eds. CRC Lewis Publishers, Boca Raton, Florida. p. 467-484.
- Arkoosh, M.R., E. Casillas, E. Clemons, A.N. Kagley, R. Olson, P. Reno, and J.E. Stein. 1998b. Effect of pollution on fish diseases: potential impacts on salmonid populations. *Journal of Aquatic Animal Health* 10:182-190.
- Arkoosh, M.R., E. Casillas, P. Huffman, E. Clemons, J. Evered, J.E. Stein, and U. Varanasi. 1998a. Increased susceptibility of juvenile Chinook salmon from a contaminated estuary to *Vibrio anguillarum*. *Transactions of the American Fisheries Society* 127:360-374.
- Armstrong, R.W., and A.W. Argue. 1977. Trapping and coded-wire tagging of wild coho and chinook juveniles from the Cowichan River system, 1975. *Fish. Mar. Serv. (Can.) Pac. Reg. Tech. Rep. Ser. PAC/T-77-14:58 p.*
- Artyukhin, E.N. and A.E. Andronov. 1990. A morphobiological study of the green sturgeon, *Acipenser medirostris* (*Chondrostei, Acipenseridae*), from the Tumnin (Datta) River and some aspects of the ecology and zoogeography of the Acipenseridae. *Journal of Ichthyology* 30:11-21.
- Bajkov, A. D. 1951. Migration of white sturgeon (*Acipenser transmontanus*) in the Columbia River. *Fish Commission of Oregon, Portland, Oregon, Department of Research* 3(2):8-21.
- Bakkala, R.G. 1970. Synopsis of biological data on the chum salmon, *Oncorhynchus keta* (Walbaum) 1792. *FAO Fish. Symposium* 41; U.S. Fish and Wildlife Service Circular 315. 89 pp.

- Baltz, D. M., B. Vondracek, L. R. Brown, and P. B. Moyle. 1987. Influence of temperature on microhabitat choice by fishes in a California stream. *Transactions of the American Fisheries Society* 116: 12-20.
- Bams, R.A. 1976. Survival and propensity for homing as affected by presence or absence of locally adapted paternal genes in two transplanted populations of pink salmon (*Oncorhynchus gorbuscha*). *Journal of the Fisheries Research Board of Canada* 33: 2716-2725.
- Barnhart, R.A. 1986. Species profiles: Life histories and environmental requirements of coastal fishes and invertebrates (Pacific Southwest) – Steelhead. Biological Report 82(11.60) U.S. Army Corps of Engineers, Coastal Ecology Group, Vicksburg, Mississippi.
- Bartley, D. M., G. A. E. Gall, and B. Bentley. 1985. Preliminary description of the genetic structure of white sturgeon, *Acipenser transmontanus* in the Pacific Northwest. in F.P. Binkowski and S.E. Dorshov, eds. *North American Sturgeons*. W. Junk Publishers, Dordrecht, The Netherlands.
- Bax, N. and C.J. Whitmus. 1981. Early marine survival and migratory behavior of juvenile salmon released from the Enetai Hatchery, Washington, in 1980. University of Washington Fisheries Research Institute FRI-UW-9809. 48 pp.
- Bax, N.J. 1982. Seasonal and annual variations in the movement of juvenile chum salmon through Hood Canal, Washington, Pages 208-218 in E.L. Brannon and E.O. Salo, eds. *Proceedings of the Salmon and Trout Migratory Behavior Symposium*. School of Fisheries, University of Washington, Seattle.
- Bax, N.J. 1982a. The early marine migration of juvenile chum salmon (*Oncorhynchus keta*) through Hood Canal – its variability and consequences. Doctoral dissertation. University of Washington, Seattle. 196 p.
- Bax, N.J. 1983b Early marine mortality of marked juvenile chum salmon (*Oncorhynchus keta*) released into the Hood Canal, Puget Sound, Washington in 1980. *Canadian Journal of Fisheries and Aquatic Sciences* 40:426-435.
- Bax, N.J., E.O. Salo, and B.P. Snyder. 1979. Salmonid outmigration studies in the Hood Canal. Final Report, Phase V. University of Washington, College of Fisheries, Seattle.
- Bax, N.J., E.O. Salo, and B.P. Snyder. 1980. Salmonid outmigration studies in Hood Canal. Final report, phase V, January to July 1979. FRI-UW-8010, 55 pp. Fisheries Research Institute, University of Washington, Seattle.
- Baxter, J. S. 1995. Chowade River bull trout studies 1995: habitat and population assessment. Report prepared for British Columbia Ministry of Environment, Lands and Parks, Fisheries Branch, Fort St. John, British Columbia, 108 p.
- Baxter, J.S. and J.D. McPhail. 1996. Bull trout spawning and rearing habitat requirements: summary of the literature. British Columbia Ministry of Environment, Lands and Parks, Fisheries Branch, Fisheries Technical Circular 98.
- Beacham, T. D., and C. B. Murray. 1990. Temperature, egg size, and development of embryos and alevins of five species of Pacific salmon: a comparative analysis. *Transactions of the American Fisheries Society* 119: 927-945.
- Beacham, T.D., and C.B. Murray. 1986. Comparative developmental biology of chum salmon (*Oncorhynchus keta*) from the Fraser River, British Columbia. *Canadian Journal of Fisheries and Aquatic Sciences* 43:252-262.

- Beacham, T.D., and C.B. Murray. 1987. Adaptive variation in body size, age, morphology, egg size, and developmental biology of chum salmon (*Oncorhynchus keta*) in British Columbia. Canadian Journal of Fisheries and Aquatic Sciences 44:244-261.
- Beacham, T.D., and P. Starr. 1982. Population biology of chum salmon, *Oncorhynchus keta*, from the Fraser River, British Columbia. Canadian Journal of Fisheries and Aquatic Sciences 43: 252-262.
- Beall, E.P. 1972. The use of predator-prey tests to assess the quality of chum salmon *Oncorhynchus keta* fry. Master's thesis. University of Washington, Seattle. 105 p.
- Beamesderfer, C.P., D.L. Ward, and A.A. Nigro. 1996. Evaluation of the biological basis for a predator control program on northern pikeminnow (*Ptychocheilus oregonensis*) in the Columbia and Snake rivers. Canadian Journal of Fisheries and Aquatic Sciences 53:2898-2908.
- Beamesderfer, R. C. P. and R. A. Farr. 1997. Alternatives for the protection and restoration of sturgeons and their habitat. Environmental Biology of Fishes 48:407-417.
- Beamesderfer, R. C. P., T. A. Rien, and A. A. Nigro. 1995. Differences in the dynamics and potential production of impounded and unimpounded white sturgeon populations in the lower Columbia River. Transactions of the American Fisheries Society 124:857-872.
- Beamesderfer, R.C. 1992. Reproduction and early life history of northern pikeminnow, *Ptychocheilus oregonensis*, in Idaho's St. Joe River. Environmental Biology of Fishes 35:231-241.
- Beamesderfer, R.C. and B.E. Rieman. 1991. Abundance and distribution of northern squaw fish, walleyes, and smallmouth bass in John Day Reservoir, Columbia River. Transactions of the American Fisheries Society 120:439-447.
- Beamesderfer, R.C., B.E. Rieman, L.J. Bledsoe, and S. Vigg. 1990. Management implications of a model of predation by a resident fish on juvenile salmonids migrating through a Columbia River reservoir. North American Journal of Fisheries Management 10:290-304.
- Beamesderfer, R.C., D.L. Ward, and A.A. Nigro. 1996. Evaluation of the biological basis for a predator control program on northern squawfish (*Ptychocheilus oregonensis*) in the Columbia and Snake rivers. Canadian Journal of Fisheries and Aquatic Sciences 53: 2898-2908.
- Beamesderfer, R.C.P., and M. Webb. 2002 Green sturgeon status review information. S.P. Cramer and Associates to State Water Contractors, 455 Capitol Mall, Suite 200, Sacramento, California, 95814.
- Beamish, R. J., C. Mahnken, and C.M. Neville. 2004. Evidence that reduced early marine growth is associated with lower marine survival of coho salmon. Transactions of the American Fisheries Society 133: 26–33.
- Beamish, R.J. 1980. Adult Biology of the River Lamprey (*Lampetra agresi*) and the Pacific Lamprey (*Lampetra tridentata*) from the Pacific Coast of Canada. Canadian Journal of Fisheries and Aquatic Sciences 37: 1906-1923.
- Beamish, R.J. and C.D. Levings. 1991. Abundance and freshwater migrations of the anadromous parasitic lamprey, *Lampetra tridentata*, in a tributary of the Fraser River, British Columbia. Canadian Journal of Fisheries and Aquatic Sciences 48: 1250-1263.
- Beamish, R.J. and D.R. Bouillon. 1993. Pacific salmon production trends in relation to climate. Canadian Journal of Fisheries and Aquatic Science 50: 1002-1016.
- Beamish, R.J. and J.H. Youson. 1987. Life history and abundance of young adult *Lampetra ayresi* in the Fraser River and their possible impact on salmon and herring stocks in the Strait of Georgia. Canadian Journal of Fisheries and Aquatic Science 44:525-537.

- Beamish, R.J., and T.G. Northcote. 1989. Extinction of a population of anadromous parasitic lamprey, *Lampetra tridentate*, upstream of an impassable dam. *Canadian Journal of Fisheries and Aquatic Sciences* 46: 420-425
- Beamish, R.J., M. Smith, R. Scarsbrook, and C. Wood. 1976. Hake and Pollock study, Strait of Georgia cruise G.B. Reed, June 16-27, 1975. *Fish. Mar. Serv. Data Rec.* 1:174 p.
- Beauchamp, D.A. and J.J. Van Tassell. 1999. Modeling seasonal trophic interactions of adfluvial bull trout in Lake Billy Chinook, Oregon. Portland General Electric, Pelton Round Butte Hydroelectric Project FERC No. 2030, Portland, Oregon.
- Becker, C.D., D.A. Neitzel, and C.S. Abernethy. 1983. Effects of dewatering on Chinook salmon redds: tolerance of four development phases to one-time dewatering. *North American Journal of Fisheries Management* 3:373-382.
- Becker, C.D., D.A. Neitzel, and D.H. Fickeisen. 1982. Effects of dewatering on Chinook salmon redds: tolerance of four development phases to daily dewaterings. *Transactions of the American Fisheries Society* 111: 624-637.
- Beecher, H.A., T.H. Johnson, and J.P. Carleton. 1993. Predicting microdistributions of steelhead (*Oncorhynchus mykiss*) parr from depth and velocity preference criteria: test of an assumption of the Instream Flow Incremental Methodology. *Canadian Journal of Fisheries and Aquatic Sciences* 50: 2380-2387.
- Beechie, T.J., G. Pess, P. Kennard, R.E. Bilby, and S. Bolton. 2000. Modeling recovery rates and pathways for woody debris recruitment in Northwestern Washington streams. *North American Journal of Fisheries Management* 20:436-452.
- Behnke, R.J. 1992. Native trout of western North America. American Fisheries Society, Bethesda, Maryland. 275 p.
- Behnke, R.J. 1997. Evolution, systematics, and structure of *Oncorhynchus clarki clarki*. Pages 3-6 in J.D. Hall, P.A. bison and R.E. Gresswell, eds., *Sea-run cutthroat trout: biology management, and nature conservation*. Oregon Chapter, American Fisheries Society Symposium 4.
- Behnke, R.J. 2002. *Tout and salmon of North America*. The Free Press, New York.
- Bell, M. C. 1986. Fisheries handbook of engineering requirements and biological criteria. U.S. Army Corps of Engineers, Portland, Oregon. Fish Passage Development and Evaluation Program, North Pacific Division.
- Bell, M.C. 1986. Fisheries handbook of engineering requirements and biological criteria. U.S. Army Corps of Engineers, Portland, Oregon. Fish Passage Development and Evaluation Program, North Pacific Division.
- Bemis, W. E., and B. Kynard. 1977. Sturgeon rivers: an introduction to Acipenseriform biogeography and life history. *Environmental Biology of Fishes* 48:167-183.
- Bemis, W.E., E.K. Findeis, and L. Grand. 1997b. An overview of *Acipenseriformes*. *Environmental Biology of Fishes* 48: 25-71.
- Bemis, W.E., V.J. Birstein, and J.R. Waldman. 1997a. Sturgeon biodiversity and conservation: an introduction. Pages 13-14 in V.J. Birstein, J.R. Waldman, and W.E. Bemis, eds. *Sturgeon Biodiversity and Conservation*. Kluwer Academic Publishers Dordrecht, Netherlands.
- Berejikian, B., and M. Ford. 2003. A review of relative fitness of hatchery and natural salmon. Preliminary Review Draft. NOAA, Seattle.

- Bergstedt, R.A., and J.G. Seelye. 1995. Evidence for lack of homing by sea lampreys. *Transactions of the American Fisheries Society* 124: 235-239.
- Beschta, R.L. 1991. Stream habitat management for fish in the Northwestern United States: the role of riparian vegetation. *American Fisheries Society Symposium* 10:53-58.
- Beschta, R.L. 1997a. Restoration of riparian and aquatic systems for improved fisheries habitat in the upper Columbia Basin. Pages 475-489 in D.J. Stouder, P.A. Bisson, and R.J. Naiman, eds. *Pacific salmon and their ecosystems: status and future options*. Chapman & Hall, New York.
- Beschta, R.L., R.E. Bilby, G.W. Brown, L.B. Holtby, and T.D. Hofstra. 1987. Stream temperature and aquatic habitat: fisheries and forestry interactions. Pages 191-232 in E.O. Salo and T.W. Cundy, eds. *Streamside management: forestry and fishery interactions*. College of Forest Resources, University of Washington, Seattle.
- Bevelhimer, M. S. 2001. A bioenergetics model for white sturgeon, *Acipenser transmontanus*: Assessing differences in growth and reproduction among Snake River reaches. in *Proceedings of the Fourth International Symposium on Sturgeons, Oshkosh, Wisconsin, 8-13 July, 2001*
- Bigler, B.S., and J.H. Helle. 1994. Decreasing size of north Pacific salmon (*Oncorhynchus* spp.): Possible causes and consequences. Unpublished manuscript. Wards Cove Packing Co., P.O. Box c-5030, Seattle, WA, 98105. 34 pp.
- Bilby, R.E., and J.W. Ward. 1991. Characteristics and function of large woody debris in streams draining old-growth, clear-cut, and second growth forests in southwestern Washington. *Canadian Journal of Fisheries and Aquatic Sciences* 12:2499-2508.
- Bilby, R.E., B.R. Fransen, and P.A. Bisson. 1996. Incorporation of nitrogen and carbon from spawning coho salmon into the trophic system of small streams. *Canadian Journal of Fisheries and Aquatic Sciences* 53: 164-173.
- Bilby, R.E., K. Sullivan, and S.H. Duncan. 1989. The generation and fate of road-surface sediment in forested watersheds in Southwestern Washington. *Forest Science* 35:453-468.
- BioAnalysts, Inc. 2000. A Status of Pacific Lamprey in the Mid-Columbia Region. Rocky Reach Hydroelectric Project FERC Project No. 2145. Prepared for Public Utility District No. 1 of Chelan County, Wentachee, Washington.
- Birstein, V.J. 1993. Is *Acipenser medirostris* one or two species? *The Sturgeon Quarterly* 1(2):8.
- Birstein, V.J. 1993. Sturgeons and paddlefishes: threatened fishes in need of conservation. *Conservation Biology* 7:773-787.
- Birstein, V.J. and W.E. Bemis. 1997. How many species are within the genus *Acipenser*? *Environmental Biology of Fishes* 48:157-163.
- Birstein, V.J., J.R. Waldman, and W.E. Bemis. 1997. Sturgeon biodiversity and conservation. Kluwer Academic Publishers. Dordrecht, The Netherlands.
- Birstein, V.J., R. Hanner, and R. LaSalle. 1997. Phylogeny of the *Acipenseriformes*: cytogenetic and molecular approaches. *Environmental Biology of Fishes* 48:127-155.
- Birstein, V.J., W.E. Bemis, and J.R. Waldman. 1997b. The threatened status of *acipenseriform* fishes: a summary. 1997. *Environmental Biology of Fishes* 48:427-435.
- Bishop, F.G. 1975. Observations on the fish fauna of the Peace River in Alberta. *Canadian Field-Naturalist* 89:423-430.

- Bisson, P.A., G. H. Reeves, R. E. Bilby, and R. J. Naiman. 1997. Watershed management and Pacific salmon: desired future conditions. Pages 447-474 In D.J. Stouder, P.A. Bisson, and R. J. Naiman, editors. Pacific salmon and their ecosystems: status and future options. Chapman and Hall, New York.
- Bisson, P.A., K. Sullivan, and J.L. Nielsen. 1988. Channel hydraulics, habitat use, and body form of juvenile coho salmon, steelhead, and cutthroat trout in streams. Transactions of the American Fisheries Society 117:262-273.
- Bjerselius R., L. Weiming, J.H. Teeter, J.G. Seelye, P.B. Johnsen, P.J. Maniak, G.C. Grant, C.N. Polkinghorne, and P.W. Sorenson. 2000. Direct behavioral evidence that unique bile acids released by larval sea Lamprey (*Petromyzon marinus*) function as a migratory pheromone. Canadian Journal of Fisheries and Aquatic Sciences 57: 557-569
- Bjornn, T. C. 1971. Trout and salmon movements in two Idaho streams as related to temperature, food, stream flow, cover, and population density. Transactions of the American Fisheries Society 100: 423-438.
- Bjornn, T. C. 1978. Survival, production, and yield of trout and salmon in the Lemhi River, Idaho. Idaho Department of Fish and Game Bulletin 27: 57.
- Bjornn, T. C., and D. W. Reiser. 1991. Habitat requirements of salmonids in streams. Pages 83-138 in W. R. Meehan, ed. Influences of forest and rangeland management on salmonid fishes and their habitats. American Fisheries Society, Bethesda, Maryland.
- Bjornn, T.C. and C.A. Peary. 1992. A review of literature related to movements of adult salmon and steelhead past dams and through reservoirs in the lower Snake River. U.S. Army Corps of Engineers, Technical Report 92-1. 80 pp.
- Bjornn, T.C., M.A. Brusven, M.P. Molnau, J.H. Milligan, R.A. Klmat, E. Chacho, and C. Schaye. 1977. Transport of granitic sediment in streams and its effects on insects and fish. University of Idaho, Forest, Wildlife, and Range Experiment Station Bulletin, Moscow 17.
- Bjornn, T.C., M.L. Keefer, C.A. Peery, K.R. Tolotti, R.R. Ringe, and L.C. Stuehrenburg. 1998. Adult chinook and sockeye salmon, and steelhead fall back rates at Bonneville Dam – 1996, 1997, and 1998. Report to U.S. Army Corps of Engineers and Bonneville Power Administration, Project MPE-P-95-1. 73 pp.
- Bjornn, T.C., M.L. Keefer, and L.C. Stuehrenburg. 1999. Behavior and survival of adult chinook salmon that migrate past dams and into tributaries in the Columbia River drainage as assessed with radio telemetry. In Proceedings of the Fifteenth International Symposium on Biotelemetry, Juneau, Alaska.
- Boag, T.D. 1987. Food habits of bull char, *Salvelinus confluentus*, and rainbow trout, *Salmo gairdneri*, coexisting in a foothills stream in northern Alberta. Canadian Field-Naturalist 101:56-62
- Boag, T.D. 1991. Round III of Delphi analysis for bull trout habitat requirements. Unpublished report by Environmental Management Associates, Calgary, Alberta, 7 p.
- Booth, D. 1990. Stream channel incision following drainage basin urbanization. Water Resources Bulletin 26(3):407-417.
- Booth, D.B. and C.R. Jackson. 1997. Urbanization of aquatic systems: degradation thresholds, stormwater detention, and the limits of mitigation. Journal of the American Water Resources Association 33 No. 5.

- Bostick, W.E. 1955. Duwamish River seining studies. Pages 5-6 in Puget Sound stream studies, Washington Department of Fisheries, Olympia. (Available Washington Department of Fish and Wildlife, 600 Capitol Way N., Olympia, WA 98501-1091).
- Bottom, D. and K. Jones. 1990. Species composition, distribution, and invertebrate prey assemblages in the Columbia River estuary. *Progress in Oceanography* 25:243-270.
- Bottom, D.L., P.J. Howell, and J.D. Rodgers. 1985. The effects of stream alterations on salmon and trout habitat in Oregon. Oregon Department of Fish and Wildlife, Portland. 70 pp.
- Bovee, K. D. 1978. Probability-of-use criteria for the family Salmonidae. U.S. Department of Interior, Fish and Wildlife Service, Washington, D.C.
- Bradford, M.J. 1992. Precision of recruitment predictions from early life stages of marine fishes. *Fishery Bulletin* 90:439-453.
- Brannon, E.L., K.P. Currens, D. Goodman, J.A. Lichatowich, B.E. Riddell, and R.N. Williams. W.E. McConaha, chair. 1999. Review of artificial anadromous and resident fish production in the Columbia River basin. Part I: A scientific basis for Columbia River production programs. NWPPC Document 99-4. Scientific Review team, Independent Scientific Advisory Board, Northwest Power Planning Council, Portland Oregon. 132 pp.
- Brannon E., M. Powell, T. Quinn, and A. Talbot. 2002. Population structure of Columbia River Basin Chinook salmon and steelhead trout. Center for Salmonid and Freshwater Species at Risk, University of Idaho. Final report to the Bonneville Power Administration, Portland, Oregon. 178 pp.
- Brannon, E. L., C. L. Melby, and S. D. Brewer. 1985b. Columbia River white sturgeon enhancement. Final report (Project Number 83-316) to Bonneville Power Administration. Portland, Oregon.
- Brannon, E., A. Setter, M. Miller, S. Brewer, G. Winans, F. Utter, L. Carpenter, and W. Hershberger. 1986. Columbia River white sturgeon population genetics and early life history study. Final report (Project 83-316) to Bonneville Power Administration. Portland, Oregon.
- Brannon, E., S. Brewer, A. Setter, M. Miller, F. Utter, and W. Hershberger. 1985a. Columbia River white sturgeon early life history and genetics study. Final report (Project Number 83-316) to Bonneville Power Administration. Portland, Oregon.
- Brege, D.A., S.J. Grabowski, W.D. Muir, S.R. Hirtzel, S.J. Mazur, and B.P. Sandford. 1992. Studies to determine the effectiveness of extended traveling screens and extended bar screens at McNary Dam, 1991. Report to U.S. Army Corps of Engineers, Contract E86970083.
- Brege, D.A., W.T. Norman, G.A. Swan, and J.G. Williams. 1988. Research at McNary Dam to improve fish guiding efficiency of yearling and subyearling Chinook salmon – 1987. Report to U.S. Army Corps of Engineers, Contract DACW68-84-H-0034, 22 pp. + App. (Available from Northwest Fisheries Science Center, 2725 Montlake Blvd. E., Seattle, WA 98112-2097.)
- Brett, J.R. 1952. Temperature tolerance in young Pacific salmon, genus *Oncorhynchus*. *Journal of the Fisheries Research Board of Canada* 9:265-323.
- Brewin, P.A. and M.K. Brewin. 1997. Distribution maps for bull trout in Alberta. Pages 206-216 in W.C. Mackay, M.K. Brewin, and M. Monita, eds. Friends of the Bull Trout Conference Proceedings. Bull Trout Task Force (Alberta). Trout Unlimited Calgary, Alberta.
- Briggs, J.C. 1953. The behaviour and reproduction of salmonids fishes in a small coastal stream. *California Department of Fish and Game Fisheries Bulletin* 94. 62 p.

- Brown, J.B., A.T. Beckenbach, and M.J. Smith. 1992. Influence of Pleistocene glaciations and human intervention upon mitochondrial DNA diversity in white sturgeon (*Acipenser transmontanus*) populations. *Canadian Journal of Fisheries and Aquatic Sciences* 49: 358-367.
- Brown, J.B., A.T. Beckenbach, and M.J. Smith. 1993. Intraspecific DNA sequence variation of the mitochondrial control region of white sturgeon (*Acipenser transmontanus*). *Molecular Biology and Evolution* 10(2):326-341.
- Brown, L.R., and P.B. Moyle. 1991. Changes in habitat and microhabitat partitioning within an assemblage of stream fishes in response to predation by Sacramento squawfish (*Ptychocheilus grandis*). *Canadian Journal of Fisheries and Aquatic Sciences* 48:849-856.
- Bruch, R., F. P. Binkowski, and S. I. Doroshov, eds. 2001. Proceedings of the Fourth International Symposium on Sturgeons, Oshkosh, Wisconsin, USA, 8-13 July 2001.
- Bryant, F.G. 1949. A survey of the Columbia River and its tributaries with special reference to its fishery resources – Part II. Washington streams from the mouth of the Columbia to and including the Klickitat River (Area I). U.S. Fish and Wildlife Service, Special Service Report 63, 110 pp.
- Buchanan, D. V., M. L. Hanson and R. M. Hooten. 1997. 1996 Status of Oregon's bull trout. Draft report. Oregon Department of Fish and Wildlife, Portland.
- Buchanan, D.V., R.M. Hooten, and J.R. Moring. 1981. Northern pikeminnow (*Ptychocheilus oregonensis*) predation on juvenile salmonids in sections of the Willamette River basin, Oregon. *Canadian Journal of Fisheries and Aquatic Sciences* 38:360-364.
- Buckman, R., W. Hosford and P. Dupee. 1992. Malheur River bull trout investigations. Pages 45-57 in P.J. Howell and D.V. Buchanan, eds. Proceedings of the Gearhart Mountain bull trout workshop. Oregon Chapter of the American Fisheries Society, Corvallis.
- Buell, J.W. 1992. Fish entrainment monitoring of the Western-Pacific dredge RW Lofgren during operations outside the preferred work period. Buell and Associates, Inc. Portland, Oregon.
- Bulkley, R.V. 1966. Catch of the 1965 tidewater cutthroat sport fishery and notes on the life history of the coastal cutthroat trout in the Siuslaw River, Oregon. Research Division, Oregon Game Commission, Fisheries Research Report 4, Oregon State University, Corvallis.
- Burner, C.J. 1951. Characteristics of spawning nests of Columbia River salmon. *Fish. Bull. Fish Wildlife Service* 61:97-110.
- Busby, P.J., O.W. Johnson, T.C. Wainwright, F.W. Waknitz, and R.S. Waples. 1993. Status review for Oregon's Illinois River winter steelhead. NOAA Technical Memorandum NMFS-NMFSC-10. 85 pp.
- Busby, P.J., T.C. Wainwright, and R.S. Waples. 1994. Status review for Klamath Mountains Province steelhead. NOAA Technical Memorandum NMFS-NMFSC-19. 130 pp.
- Busby, P.J., T.C. Wainwright, G.J. Bryant, L. Leirheimer, R.S. Waples, F.W. Waknitz, and I.V. Lagomarsino. 1996. Status review of west coast steelhead from Washington, Idaho, Oregon, and California. NOAA Technical Memorandum. NMFS-NWFSC-27. 281 pp.
- Busch, W. N., R. L. Scholl, and W. L. Hartman. 1975. Environmental factors affecting the strength of walleye (*Stizostedion vitreum vitreum*) year-classes in western Lake Erie, 1969-1970. *Journal of the Fisheries Research Board of Canada* 32:1733-1743.
- Bustard, D. R., and D. W. Narver. 1975. Aspects of the winter ecology of juvenile coho salmon (*Oncorhynchus kisutch*) and steelhead trout (*Salmo gairdneri*). *Journal of the Fisheries Research Board of Canada* 32: 667-680.

- California Department of Fish and Game (CDFG). 1994. Documents submitted to the ESA Administrative Record for coastal steelhead: Annual reports for salmon and steelhead hatcheries.
- Campton, D.E. 1995. Genetic effects of hatchery fish on wild populations of Pacific salmon and steelhead: what do we really know? Pages 337-353 in H. L. Schramm, Jr. and R. G. Piper, editors, Uses and Effects of Cultured Fishes in Aquatic Ecosystems. American Fisheries Society Symposium 15, Bethesda, Maryland.
- Carl, G.C., W.A. Clemens, and C.C. Lindsey. 1977. The freshwater fishes of British Columbia. British Columbia Province Museum Handbook 5. 192 pp.
- Carl, L.M., and M.C. Healey. 1984. Differences in enzyme frequency and body morphology among three juvenile life history types of Chinook salmon (*Oncorhynchus tshawytscha*) in the Nanaimo River, British Columbia. Canadian Journal of Fisheries and Aquatic Sciences 41:1070-1077.
- Carlson, J.Y., C.W. Andrus, and H.A. Froehlich. 1990. Woody debris, channel features, and macroinvertebrates of streams with logged and undisturbed riparian timber in northeastern Oregon, USA. Canadian Journal of Fisheries and Aquatic Sciences 47.
- Casell, H. 2001. Matrix population models, 2nd edition. Sinauer Associates, Sunderland, Massachusetts.
- Casillas, E., B.B. McCain, M. Arkoosh, and J.E. Stein. 1996. Estuarine pollution and juvenile salmon health: potential impact on survival. Pages 169-178 in Estuarine and Ocean Survival of northeastern Pacific Salmon. NOAA Technical Memorandum NMFS-NWFSC-29.
- Cavender, T. M. 1978. Taxonomy and distribution of the bull trout *Salvelinus confluentus* (Suckley). American Northwest. California Fish and Game 64:139-174.
- Cech, J.J., Jr., Doroshov, S.I., Moberg, G.P., May, B.P., Schaffter, R.G., and D.M. Kohlhorst. 2000. Biological assessment of green sturgeon in the Sacramento-San Joaquin watershed (phase 1). Final Report to CALFED Bay-Delta Program (Project 98-C-15, Contract B-81738).
- Cederholm, C.J., and W.J. Scarlett. 1981. Seasonal immigrations of juvenile salmonids into four small tributaries of the Clearwater River, Washington, 1977-1981, Pages 98-110 in E.L. Brannon and E.O. Salo, eds. Proceedings of the Salmon and Trout Migratory Behavior Symposium. School of Fisheries, University of Washington, Seattle.
- Cederholm, C.J., D.B. Houston, D.L. Cole, and W.J. Scarlett. 1989. Fate of coho salmon (*Oncorhynchus kisutch*) carcasses in spawning streams. Canadian Journal of Fisheries and Aquatic Sciences 46:1347-1355.
- Cederholm, C.J., M.D. Kunze, T. Murota, and A. Sibatani. 1999. Pacific salmon carcasses: Essential contributions of nutrients and energy for aquatic and terrestrial ecosystems. Fisheries 24 (10): 6-15.
- Cederholm, C.J., W.J. Scarlett, and N.P. Peterson. 1988. Low-cost enhancement technique for winter habitat of juvenile salmon. North American Journal of Fisheries Management 8:438-441.
- Chamberlain, T. W., R. D. Harr, and F. H. Everest. 1991. Timber Harvesting, Silviculture, and Watershed Processes. Chapter 6 in Meehan, W.R., ed. Influence of Forest and Rangeland Management on salmonid fishes and their habitats.
- Chapman, D et al. 1991. Status of Snake River Chinook salmon. Pacific Northwest Utilities Conference Committee, Don Chapaman Consultants. 275pp.

- Chapman, D. W., and T. C. Bjornn. 1969. Distribution of salmonids in streams, with special reference to food and feeding. Pages 153-176 in T.G. Northcote, ed. H.R. MacMillan Lectures in Fisheries. University of British Columbia, Institute of Fisheries, University of British Columbia, Vancouver.
- Chapman, D.W. 1962. Aggressive behaviour in juvenile coho salmon as a cause of emigration. Journal of the Fisheries Research Board of Canada 19:1047-1080.
- Chapman, D.W. 1965. Net production of juvenile coho salmon in three Oregon streams. Transactions of the American Fisheries Society 94:40-52.
- Chapman, D.W. and E. Knudsen. 1980. Channelization and livestock impacts on salmonid habitat and biomass in western Washington. Transactions of the American Fisheries Society 109:357-63,
- Chapman, D.W., A. Giorgi, T. Hillman, D. Deppert, M. Erho, S. Hays, C. Peven, B. Suzumoto, and R. Klinge. 1994. Status of summer/fall Chinook salmon in the mid-Columbia region. Report for Chelan, Douglas and Grant County PUDs. 412 pp.+ app. (Available from Don Chapman Consultants, 3653 Rickenbacker, Ste. 200, Boise, ID 83705).
- Chapman, D.W., D.E. Weitcamp, T.L. Welsh, M.B. Dell, and T.H. Schadt. 1986. Effects of river flow on the distribution of Chinook salmon redds. Transactions of the American Fisheries Society 115:537-547.
- Chapman, F. A., J. P. Van Eenennaam, and S. I. Doroshov. 1996. The reproductive condition of white sturgeon, *Acipenser transmontanus*, in San Francisco Bay, California. Fishery Bulletin 94:628-634.
- Cheng, K.M., I.M. McCallum, R.I. McKay, B.E. March. 1987. A comparison of survival and growth of two strains of Chinook salmon (*Oncorhynchus tshawytscha*) and their crosses reared in confinement. Aquaculture 67: 301-311.
- Chilcote, M., S. Leider, and J. Loch. 1986. Differential reproductive success of hatchery and wild summer-run steelhead under natural conditions. Transactions of the American Fisheries Society 115:726-735.
- Chilcote, M.C., B.A. Crawford, and S.A. Leider. 1980. A genetic comparison of sympatric populations of summer and winter steelhead. Transactions of the American Fisheries Society 109:203-208.
- Chilcote, M.W. S.A. Leider, and J.J. Loch. 1986. Differential reproductive success of hatchery and wild summer-run steelhead under natural conditions. Transactions of the American Fisheries Society 115: 726-735.
- Claire, C. W. 2003. Pacific lamprey larvae life history and habitat utilization in Red River subbasin, South Fork Clearwater River drainage, Idaho. Draft Master's thesis, University of Idaho, Moscow. 74 pp.
- Close, D.A., M. Fitzpatrick, H. Li, B. Parker, D. Hatch, and G. James. 1995. Status report of the Pacific lamprey (*Lampetra tridentate*) in the Columbia River Basin. Report (Contract 95BI39067) to Bonneville Power Administration, Portland, Oregon.
- Coffin, B.A. and R.D. Harr 1992. Effects of Forest Cover on Volume of Water Delivery to Soil During Rain-on-Snow, final report, Project SH1-92-001. Timber Fish and Wildlife Program, Olympia, Washington.
- Cone, J. and S. Ridlington. 1996. The Northwest Salmon Crisis. A Documentary History. Oregon State University Press. Corvallis.

- Congleton, J.L. 1979. Feeding patterns of juvenile chum in the Skagit River salt marsh, Pages 141-150 in S.J. Lipvosky and C.A. Simenstad, eds. Gutshop '78: fish food habits studies: Proceedings of the Second Northwest Technical Workshop. Washington Sea Grant Program WSG-WO-79-1.
- Connolly, P.J. 1996. Resident cutthroat trout in the central Coast Range of Oregon: logging effects, habitat associations, and sampling protocols. Doctoral dissertation. Oregon State University, Corvallis.
- Connolly, P.J. 1997. Influence of stream characteristics and age-class interactions on populations of coastal cutthroat trout. Pages 173-174 In J.D. Hall, P.A. Bisson and R.E. Gresswell, eds. Sea-run cutthroat trout: biology, management, and future conservation. American Fisheries Society, Corvallis.
- Connolly, P.J., and B.E. Rieman. 1988. Population dynamics of walleye and smallmouth bass and potential predation on juvenile salmonids in a mainstem Columbia River reservoir, Pages 307-348 in T.P. Poe and B.E. Rieman, eds. Predation by resident fish on juvenile salmonids in John Day Reservoir, 1983-1986. Final Report (Contracts DE-AI79-82BP34796 and DE-AI79-82BP35097) to Bonneville Power Administration, Portland, Oregon.
- Conte, F.S., S.I. Doroshov, P.B. Lutes, and M.E. Strange. 1988. Hatchery manual for the white sturgeon (*Acipenser transmontanus*) with application to other North American Acipenseridae. Publications Division, Agriculture and Natural Resources, University of California, Oakland. Publication 3322.
- Cooke, D.W., S.D. Leach, and J.J. Isely. 2002. Behavior and lack of upstream passage of shortnose sturgeon at a hydroelectric facility/navigation lock complex. Pages 101-110 in W. VanWinkle, P. Anders, D. Dixon, and D. Secor, eds. Biology, Management and Protection of North American Sturgeons. American Fisheries Society Press, Bethesda, Maryland.
- Cooney, R.T., D. Urquhart, R. Neve, J. Hilsinger, R. Clasby, and D. Barnard. 1978. Some aspects of the carrying capacity of Prince William Sound, Alaska, for hatchery released pink and chum salmon fry. Alaska Sea Grant Rep. 78-4; University of Alaska Inst. Mar. Resources. IMS R78-3:98 p.
- Cooper, R., and T. H. Johnson. 1992. Trends in steelhead (*Oncorhynchus mykiss*) abundance in Washington and along the Pacific coast of North America. Washington Department of Wildlife, Fisheries Management Division. 90pp.
- Coots, M. 1957. The spawning efficiency of king salmon (*Oncorhynchus tshawytscha*) in Fall Creek, Siskiyou County: 1954-55 investigations. California Department of Fish and Game, Inland Fish. Admin. Report 57-1:15.
- Coronado, C. and R. Hilborn. 1998. Spatial and temporal factors affecting survival in coho salmon (*Oncorhynchus kisutch*) in the Pacific Northwest. Canadian Journal of Fisheries and Aquatic Science 55(9): 2067-2077.
- Counihan, T. C., J. D. DeVore, and M. J. Parsley. In Review. The effect of discharge and water temperature on the year-class strength of Columbia River white sturgeon. Submitted to Transactions of the American Fisheries Society.
- Counihan, T.C., A.I. Miller, M.G. Mesa, and M.J. Parsley. 1998. The effects of dissolved gas supersaturation on white sturgeon larvae. Transactions of the American Fisheries Society 127:316-322.
- Courchamp, F.T. Clutton-Brock, and B. Grenfell. 1999. Inverse density-dependence and the Allee effect. Trends Ecol. Evolution 14:405-410.

- Craig, J. A. and R. L. Hacker. 1940. Sturgeon fisheries of the Columbia River Basin. Bulletin of the Bureau of Fisheries 49:204:208.
- Cramer, D.P., and S.P. Cramer. 1994. Status and Population Dynamics of coho salmon in the Clackamas River. Technical Report, Portland General Electric Company, Portland, Oregon.
- Cramer, F.K. 1940. Notes on the natural spawning of cutthroat trout *Salmo clarkii clarkii* in Oregon. Proceedings of the Sixth Pacific Science Conference of the Pacific Science Association 3:335-339.
- Cramer, S. P. 1986. The influence of river temperature and flow on the production and harvest of summer steelhead in the Rogue River. Portland General Electric and Portland Water Bureau, Portland, Oregon.
- Cramer, S. P., T. Satterwaite, R. Boyce, and B. McPherson. 1985. Impacts of Lost Creek Dam on the biology of anadromous salmonids in the Rogue River. Oregon Department of Fish and Wildlife, Submitted to U.S. Army Corps of Engineers, Portland, Oregon.
- Cramer, S.P. 1998. Risk of extinction for cutthroat trout in the Umpqua Basin. S.P. Cramer & Associates, Inc. Submitted to Ronald Yokum, Attorney. (Available from Environmental and Technical Services Division, NMFS, 525 NE Oregon St., Portland, OR 97232.)
- Cramer, S.P. 1999. Evidence for an optimum escapement of sockeye salmon in the Kenai River basin. S.P. Cramer and Associates contract report to the Kenai River Sportfishing Association.
- Cramer, S.P., and R.C.P. Beamesderfer. 2002. Population dynamics, habitat capacity, and a life history simulation model for steelhead in the Deschutes River, Oregon. Prepared for Portland General Electric, S.P. Cramer and Associates, Sandy, Oregon.
- Cramer, S.P., D.B. Lister, P.A. Monk, and K.L. Witty. 2003. A review of abundance trends, hatchery and wild fish interactions, and habitat features for the mid-Columbia steelhead ESU. Prepared for Mid-Columbia stakeholders.
- Crane, P.A., L.W. Seeb, and J.E. Seeb. 1994. Genetic relationships among *Salvelinus* species inferred from allozyme data. Canadian Journal of Fisheries and Aquatic Sciences 51(Supplement 1): 182-197.
- Crawford, B.A. 1979. The origin and history of the trout brood stocks of the Washington Department of Game. Washington State Game Dept., Fishery Research Report, 76 p.
- Crone, R.A., and C.E. Bond. 1976. Life history of coho salmon *Oncorhynchus kisutch*, in Sashin Creek, southeastern Alaska. Fishery Bulletin (U.S.) 74:897-923.
- Crouse, M.R., C.A. Callahan, K.W. Malveg, S.E. Dominquez. 1981. Effects of fine sediments on growth of juvenile coho salmon in laboratory streams. Transactions of the American Fisheries Society 110(2): 281-286.
- Cuenco, M.L., T.W. Backman, and P.R. Mundy. 1993. The use of supplementation to aid in natural stock restoration. Pages 269-293 in J.G. Cloud and G.H. Thorgaard, editors. Genetic conservation of salmonid fishes. Plenum Press, New York.
- Currens, K.P., C.B. Schreck, and H.W. Li. 1990. Allozyme and morphological divergence of rainbow trout (*Oncorhynchus mykiss*) above and below waterfalls in the Deschutes River, Oregon. Copeia 1990:730-746.
- Currens, K.P., S.L. Stone, and C.B. Schreck. 1987. A genetic comparison of rainbow trout (*Salmo gairdneri*) above and below Izee Falls in the John Day River, Oregon. Oregon Cooperative Fisheries Research Unit Genet. Laboratory Report 87(2). 34 pp.

- Dambacher, J.M. 1991. Distribution, abundance, and emigration of juvenile steelhead (*Oncorhynchus mykiss*), and analysis of stream habitat in the Steamboat Creek basin. Pages 129. Oregon State University, Corvallis.
- Dames and Moore. 1976. Environmental studies report, January-December 1975, Skagit nuclear power project, Northwestern Washington. Report to Puget Sound Power and Light Company, Seattle, Washington. 376 pp.
- Deng, X.J. P Van Eenennaam, and S.I. Doroshov. 2002. Comparison of early life stages and growth of green sturgeon and white sturgeon. Pages 217-248 in Van Winkle et al., eds. Biology, management, and Protection of North American Sturgeons. American Fisheries Society Symposium 28.
- Detlaff, T.A., A.S. Ginsburg, and O.J. Schmalhausen. 1993. Sturgeon fishes: developmental biology and aquaculture. Springer-Verlag, Berlin. 300 pp.
- DeVore, J. 1987. Cowlitz River salmon investigation program: analysis of the 1983-1985 Cowlitz River runs of fall chinook and coho salmon. Washington Department of Fisheries.
- DeVore, J., B. James, and R. Beamesderfer. 1999. Lower Columbia River white sturgeon current stock status and management implications. Washington Department of Fish and Wildlife Report Number 99-08, Olympia, Washington.
- DeVore, J.D., B.W. James, C.A. Tracy, and D.A. Hale. 1995. Dynamics and potential production of white sturgeon in the Columbia River downstream from Bonneville Dam. Transactions of the American Fisheries Society 124:845-856.
- DeWitt, J.W. 1954. A survey of the coastal cutthroat trout, *Salmo clarki clarki* Richardson in California. California Fish and Game 40:329-335.
- Dill, L.M. 1969. The sub-gravel behaviour of Pacific salmon larvae, Pages 89-99 in T.G. Northcote, ed. Symposium on Salmon and Trout in Streams. H.R. MacMillan Lectures in Fisheries. Institute of Fisheries, University of British Columbia, Vancouver.
- Dill, L.M. 1983. Adaptive flexibility in the foraging behavior of fishes. Canadian Journal of Fisheries and Aquatic Sciences 40:398-408.
- Disler, N.N. 1954. Development of autumn chum salmon in the Amur River, Pages 129-143 in Trudy Soveshchaniia po voprosam lososevogo khozyaistva dal'nego vostoka, 1953. Tr. Soveshch. Ikhtiol. Kom. 4. (Transl. from Russian; Israel Program for Scientific Translations, Jerusalem, 1963).
- Don Chapman Consultants. 1989. Summer and winter ecology of juvenile chinook salmon and steelhead trout in the Wenatchee River, Washington. Don Chapman Consultants, Boise, Idaho, contract report submitted to Chelan County Public Utility District, Wenatchee, Washington. 301p.
- Dorcey, A.H.J., T.G. Northcote, and D.V. Ward. 1978. Are the Fraser River marshes essential to salmon? University of British Columbia Press, Vancouver. 29 pp.
- Downton, M.W. and K.A. Miller. 1998. Relationships between Alaskan salmon catch and North Pacific climate on interannual and interdecadal time scales. Canadian Journal of Fisheries and Aquatic Sciences 54: 2255-2265.
- Drucker, B. 1972. Some life history characteristics of coho salmon of Karluk River system, Kodiak Island, Alaska. Fisheries Bulletin (U.S.) 70:79-94.

- Duke, S., P. Anders, G. Ennis, R. Hallock, J. Hammond, S. Ireland, J. Laufle, L. Lockard, B. Marotz, V. Paragamian, and R. Westerhof. 1999. Recovery plan for Kootenai River white sturgeon (*Acipenser transmontanus*). *Journal of Applied Ichthyology* 15:157-163.
- Dunford, W.E. 1975. Space and food utilization by salmonids in marsh habitats of the Fraser River estuary. Master's thesis. University of British Columbia, Vancouver. 81 p.
- EA Engineering, Science, and Technology, 1998. Tilton watershed assessment. Prepared for the U.S. Forest Service.
- Easterbrooks, J.A. 1980. Salmon production potential evaluation for the Cowlitz River system upstream of the Cowlitz Falls Dam Site. Washington Department of Fisheries.
- Ebel, W.J. 1969. Supersaturation of nitrogen in the Columbia River and its effect on salmon and steelhead trout. U.S. National Marine Fisheries Service, Fishery Bulletin 68: 1-11.
- Ebel, W.J., and H.L. Raymond. 1976. Effect of atmospheric gas supersaturation on salmon and steelhead trout of the Snake and Columbia rivers. *Marine Fisheries*.
- Ebel, W.J., H.L. Raymond, G.E. Monan, W.E. Farr, and G.K. Tanonaka. 1975. Effects of atmospheric gas supersaturation caused by dams on salmon and steelhead trout of the Snake and Columbia Rivers. Northwest Fisheries Science Center. Processed Report, 111 pp.
- Eddie, B.G. 1975. A census of the juvenile salmonids of the Clearwater River basin, Jefferson County, Washington, in relation to logging. Master's thesis, University of Washington, Seattle. 86 p.
- Edwards, R.T. 1998. The Hyporheic Zone. Chapter 16 in *River Ecology and Management – Lessons from the Pacific Coastal Ecoregion*. Springer-Verlag, New York.
- Enk, M. 1985. Modeling the effects of forest sediment on bull trout. Page 5 in D.D. MacDonald, ed. *Proceedings of the Flathead River basin bull trout biology and population dynamics modeling exchange*. British Columbia Ministry of the Environment, Fisheries Branch, Cranbrook, British Columbia.
- Espenson, B. 2003. More sea lions at Bonneville Dam enjoying salmon cuisine. *Columbia Basin Bulletin* 5/16/2003.
- Everest, F. E., R.L. Beschta, J.C. Scrivener, K.V. Koski, J.R. Sedell, and C.J. Cederholm. 1987. Fine sediment and salmonid production: a paradox. Pages 98-142 in E.O. Salo and T.W. Cundy, eds. *Streamside management: forestry and fishery interactions*. University of Washington, Institute of Forest Resources, Seattle.
- Everest, F.H., and D.W. Chapman. 1972. Habitat selection and spatial interaction by juvenile chinook salmon and steelhead trout in two Idaho streams. *Journal of the Fisheries Research Board of Canada* 29: 91-100.
- Falter, C.M. 1969. Digestive rates and daily rations of northern pikeminnow in the St. Joe River, Idaho. Doctoral dissertation, University of Idaho, Moscow.
- Farley, E.V. Jr. and J.M. Murphy. 1997. Time series outlier analysis: evidence for management and environmental influences on sockeye salmon catches in Alaska and Northern British Columbia. *Alaska Fishery Research Bulletin* 4(1): 36-53.
- Fast, D.E., and Q.J. Stober. 1984. Intragravel behavior of salmonids alevins in response to environmental changes. Final report for Washington Water Research Center and City of Seattle. University of Washington Fish Research Institute FRI-UW-8414.103 pp.

- Fausch, K.D. 1993. Experimental analysis of microhabitat selection by juvenile steelhead (*Oncorhynchus mykiss*) and coho salmon (*O. kisutch*) in a British Columbia stream. *Canadian Journal of Fisheries and Aquatic Sciences* 50: 1198-1207.
- Fedorenko, A.Y., F.J. Fraser, and D.T. Lightly. 1979. A limnological and salmonids resource study of Nitinat Lake: 1975-1977. *Fish. Mar. Serv. (Can.) Technical Report* 839. 86 pp.
- Fidler, L.E. and S.B. Miller. 1993. British Columbia water quality guidelines for dissolved gas supersaturation. Report to British Columbia Ministry of Environment, Canada Department of Environment, Water quality Branch, Water Management Division.
- Findeis, E. K. 1997. Osteology and phylogenetic interrelationships of sturgeons (*Acipenseridae*). *Environmental Biology of Fishes* 48:73-126.
- Flagg, T.A., F.W. Waknitz, D.J. Maynard, G.B. Milner, and C.V. Mahnken. 1995. The effect of hatcheries on native coho salmon populations in the lower Columbia River. In: Schramm, H.I. and Piper, R.G., eds. *Uses and effects of cultured fishes in aquatic ecosystems*. American Fisheries Society Symposium 15. Bethesda, Maryland.
- Foerster, R.E., and W.E. Ricker. 1941. The effect of reduction of predaceous fishes on survival of young sockeye salmon at Cultus Lake. *Journal of the Fisheries Research Board of Canada* 5:315-336.
- Foerster, R.E., and W.E. Ricker. 1953. The coho salmon of Cultus Lake and Sweltzer Creek. *Journal of the Fisheries Research Board of Canada* 10:293-319.
- Fraley, J.J., and B.B. Shepard. 1989. Life history, ecology and population status of migratory bull trout (*Salvelinus confluentus*) in the Flathead Lake and River system, Montana. *Northwest Science* 63:133-143.
- Franklin, I.R. 1980. Evolutionary change in small populations. Pages 135-149 in M.E. Soule and B.A. Wilcox, eds. *Conservation biology: an evolutionary-ecological perspective*. Sinauer Associates, Sunderland, Massachusetts.
- Fraser, F. J., D.D. Bailey, and M.J. Wood. 1978. Big Qualicum River salmon development project. Vol. III: Experimental rearing of chum salmon juveniles (*Oncorhynchus keta*) in fresh water (1968-70). *Can. Fish. and Mar. Service Technical Report* 752. 22 pp.
- Fraser, F.J., E.A. Perry, and D.T. Lightly. 1983. Big Qualicum River salmon development project. Volume 1: a biological assessment 1959-1972. *Can. Technical Report of Fisheries and Aquatic Sciences* 1189. 198 pp.
- Fredin, R.A., R.L. Major, R.G. Bakkala, and G. Tanonaka. 1977. Pacific salmon and the high seas salmon fisheries of Japan. (Processed report.) Northwest and Alaska Fisheries Center, National Marine Fisheries Service, Seattle, Washington. 324 pp.
- Fresh, K.L. 1997. The role of competition and predation in the decline of anadromous salmonids in the Pacific Northwest. Pages 245-276 in D. J. Stouder//P. A. Bisson//R. J. Naiman, editors. *Pacific salmon and their ecosystems: status and future options*. Chapman and Hall, New York.
- Fresh, K.L., and G. Luchetti. 2000. Protecting and restoring the habitats of anadromous salmonids in the Lake Washington watershed, an urbanizing ecosystem. Pages 525-544 in E.E. Knudsen, C. S. Steward, D. D. MacDonald, J. E. Williams, and D. W. Reiser, editors. *Sustainable fisheries management: Pacific salmon*, Lewis Publishers, Boca Raton, Florida.
- Fresh, K.L., R.D. Cardwell, B.P. Snyder, and E.O. Salo. 1980. Some hatchery strategies for reducing predation upon juvenile chum salmon (*Onchorhynchus keta*) in freshwater. Pages 79-89 in Melteff, B.R., and R.A. Neve, eds. *Proceedings of the North Pacific Aquaculture Symposium*,

- Anchorage Alaska, August 18-21, 1980 [and] Newport, Oregon, August 25- 27, 1980. Publication 82-2, Alaska Sea Grant Program, University of Alaska, Fairbanks.
- Friesen, T.A., and D.L. Ward. 1999. Management of northern pikeminnow and implications for juvenile salmonid survival in the lower Columbia and Snake rivers. *North American Journal of Fisheries Management* 19:406-420.
- Frissell, C. A. 1992. Cumulative effects of land use on salmon habitat in Southwest Oregon coastal streams. Doctoral dissertation, Oregon State University, Corvallis.
- Frissell, C.A. 1993. A new strategy for watershed restoration and recovery of Pacific salmon in the Pacific Northwest. Pacific Rivers Council, Eugene, Oregon.
- Fulton, L.A. 1968. Spawning areas and abundance of Chinook salmon, *Oncorhynchus tshawytscha*, in the Columbia River Basin – Past and present. U.S. Fish and Wildlife Service Special Sci. Report Fish. 571. 26pp.
- Fuss, H.J. 1982. Age, growth and instream movement of Olympic Peninsula coastal cutthroat trout (*Salmo clarki clarki*). Master's thesis, University of Washington, Seattle. 128 pp.
- Gabriel, W., and R. Borger. 1992. Survival of small populations under demographic Stochasticity. *Theoretical Population Biology* 41:44-71.
- Gadomski, D.M., M.J. Parsley, D.G. Gallion, P. Kofoot. 2000. Pages 48-113 in D.L. Ward, ed. White sturgeon mitigation and restoration in the Columbia and Snake rivers upstream from Bonneville Dam. Annual progress report submitted to Bonneville Power Administration (Project 86-50), Portland, Oregon.
- Gadomski, D.M., M.J. Parsley, D.G., P. Kofoot. 2001. Pages 92-110 in D.L. Ward, ed. White sturgeon mitigation and restoration in the Columbia and Snake rivers upstream from Bonneville Dam. Annual progress report submitted to Bonneville Power Administration (Project 86-50), Portland, Oregon.
- Gadomski, D.M., M.J. Parsley, D.G., P. Kofoot. 2002. In preparation for D.L. Ward, ed., White sturgeon mitigation and restoration in the Columbia and Snake rivers upstream from Bonneville Dam. Annual progress report submitted to Bonneville Power Administration (Project 86-50), Portland, Oregon.
- Galbreath, J.L. 1985. Status, life history, and management of Columbia River white sturgeon, *Acipenser transmontanus*. In: F. P. Binkowski and S. I. Doroshov eds. *North American Sturgeons*. Dr. W. Junk Publishers, The Netherlands.
- Galbreath, J.L., and R.L. Ridenhour. 1966. Fecundity of Columbia River Chinook salmon. *Research Briefs Fish Comm. Oregon* 10(1):16-27
- Gallagher, A.F., Jr. 1979. An analysis of factors affecting brood year returns I the wild stocks of Puget Sound chum (*Oncorhynchus keta*) and pin salmon (*Oncorhynchus gorbuscha*). Master's thesis. University of Washington, Seattle. 152 pp.
- Gallion, D., and M.J. Parsley. 2001. Seasonal habitat use and movements of subadult and adult white sturgeon *Acipenser transmontanus* in a free flowing and impounded reach of the Columbia River. in: *Proceedings of the Fourth International Symposium on Sturgeons*, Oshkosh, Wisconsin, USA, 8-13, July, 2001.
- Gangmark, H.A. and R.D. Broad. 1955. Experimental hatching of king salmon in Mill Creek, a tributary of the Sacramento River. *California Fish and Game* 41:233-242.

- Gangmark, H.A., and R.G. Bakkala. 1960. A comparative study of unstable and stable (artificial channel) spawning streams for incubating king salmon at Mill Creek. *California Fish and Game* 46:151-164.
- Giger, R.D. 1972. Ecology and management of coastal cutthroat trout in Oregon. Oregon State Game Commission, Fish. Research Report 6. 61 pp.
- Gilbert, C.H. 1913. Age at maturity of the Pacific coast salmon of the genus *Oncorhynchus*. *Bull. Bur. Fish. (U.S.)* 32:1-22.
- Gilbreath, L. and E. Prentice. 1999. Post construction evaluation of the modified Bonneville Dam Second Powerhouse juvenile bypass system. In Abstracts, anadromous fish evaluations program 1999 annual research review, November 1999. (Available from U.S. Army Corps of Engineers, Walla Walla District, Walla Walla, WA)
- Gill, R. E. and L.R. Mewaldt. 1983. Pacific coast Caspian terns: Dynamics of an expanding population. *The Auk* 100: 369-381.
- Glova, G.J. 1987. Comparison of allopatric cutthroat trout stocks with those sympatric with coho salmon and sculpins in small streams. *Environmental Biology of Fishes* 20:275-284.
- Glova, G.J., and J.C. Mason. 1976. Interactive ecology of juvenile salmon and trout in streams. *Fish. Res. Board Can., Pacific Biological Station, Nanaimo, British Columbia. Manuscript Rep. Ser. 1391*, 24 p.
- Godfrey, H. 1965. Coho salmon in offshore waters, Pages 1-39. in *Salmon of the North Pacific Ocean. Part IX. Coho, chinook, and masu salmon in offshore waters. International North Pacific Fisheries Comm. Bulletin* 16.
- Godfrey, H., K.A. Henry, and S. Machidori. 1975. Distribution and abundance of coho salmon in offshore waters of the North Pacific Ocean. *International North Pacific Fisheries Comm. Bulletin* 31. 80 pp.
- Goetz, F. 1989. Biology of the bull trout, *Salvelinus confluentus*, literature review. Willamette National Forest, Eugene, Oregon.
- Goetz, F.A. 1997. Distribution of bull trout in cascade mountain streams of Oregon and Washington. Pages 339-351 in W.C. McKay, M.K. Brewin and M. Monita, eds. Friends of the bull trout conference proceedings. Bull Trout Task Force (Alberta), Trout Unlimited Canada, Calgary.
- Golder Associates. 2003. White sturgeon investigations in the Priest Rapids and Wanapum reservoirs of the middle Columbia River, Washington, USA. 2003 Final report to Grant Co. (Washington) PUD.
- Goodman, D. 1975. A synthesis of the impacts of proposed expansion of the Vancouver International Airport and other developments on the fisheries resources of the Fraser River estuary. Volumes I and II, Section II. in Fisheries resources and food web components off the Fraser River estuary and an assessment of the impacts of proposed expansion of the Vancouver International Airport and other developments on these resources. Prepared by Department of Environment, Fisheries and Marine Service. Environment Canada, Vancouver.
- Gould, W.R. 1987. Features in the early development of bull trout *Salvelinus confluentus*. *Northwest Science* 61(4):264-268.
- Grande, L., and W. E. Bemis. 1991. Osteology and phylogenetic relationships of fossil and recent paddlefishes (*Polyodontidae*) with comments on the interrelationships of Acipenseriformes. *Journal of Vertebrate Paleontology*, II supplement 1:1-121.

- Grande, L., and W. E. Bemis. 1997. A comprehensive phylogenetic study of amiid fishes (*Amiidae*) based on comparative skeletal anatomy. *Journal of Vertebrate Paleontology*.
- Gregory, S.V., F.J. Swanson, W.A. McKee, and K.W. Cummins. 1991. An ecosystem perspective of riparian zones. *BioScience* 41:540-551.
- Gribanov, V.I. 1948. The coho salmon (*Onchorhynchus kisuts Walb.*)- a biological sketch. *Izv. Tikhookean. Nauchno-Issled. Inst. Rybn. Khoz. Okeanogr.* 28:43-101. (Translated from Russian; Fisheries Research Board of Canada Translation Ser. 370).
- Griffith, J.S. 1988. Review of competition between cutthroat trout and other salmonids. Pages 134-140. In R.E. Gresswell, ed. Status and management of interior stocks of cutthroat trout. American Fisheries Society Symposium 4.
- Groot, C. 1982. Modifications on a theme-a perspective on migratory behavior of Pacific salmon in E. L. Brannon and E. O. Salo, eds. Salmon and trout migratory behavior symposium, University of Washington, College of Fisheries, Seattle.
- Groot, C. and L. Margolis. 1991. Pacific salmon life histories. University of British Columbia Press, Vancouver.
- Gross, M.R., J. Repka, D.H. Secor, and W. Van Winkle. 2002. Conserving ancient fish: Life history and demography of sturgeons. Pages 13-30 in W. Van Winkle, P. Anders, D. Dixon, and D. Secor, eds. Biology, Management and Protection of North American Sturgeons. American Fisheries Society Press, Bethesda, Maryland.
- Gunsolus, R.T. 1978. Th status of Oregon coho and recommendations for managing the production, harvest, and escapement of wild and hatchery reared stocks. Oregon Department of Fish and Wildlife, Columbia Region, Portland, Oregon. 59 pp.
- Hammond, R.J. 1979. Larval biology of the Pacific lamprey, *Entosphenus tridentatus* (Gairdner) of the Potlach River, Idaho. Master's thesis. University of Idaho, Moscow.
- Hankin, D.G., and J. Richards. 2000. The northern pikeminnow management program: An independent review of program justification, performance, and cost-effectiveness. Final Report to the Northwest Power Planning Council.
- Hardisty, M.W., and I.C. Potter. 1971. The general biology of adult lampreys. Pages 127-206 in M.W. Hardisty and I.C. Potter, eds. The Biology of Lampreys, Vol. 1. Academic Press, Inc., London and New York.
- Hare, S.R., N.J. Mantua, R.C. Francis. 1999. Inverse production regimes: Alaska and West Coast Pacific salmon. *Fisheries* 24(1): 6-14.
- Hart, J.L. 1973. Pacific Fishes of Canada. *Bulletin of the Fisheries Research Board of Canada* 180:83-84.
- Hart, J.L. 1973. Pacific fishes of Canada. *Bulletin of the Fisheries Research Board of Canada* 180:740 p.
- Hartman, G. F., C. Groot, and T. G. Northcote. 2000. Science and management in sustainable salmonid fisheries: the ball is not in our court. Pages 31-50 in E.E. Knudsen, C. S. Steward, D. D. MacDonald, J. E. Williams, and D. W. Reiser, editors. Sustainable fisheries management: Pacific salmon, Lewis Publishers, Boca Raton, Florida.
- Hartman, G.F. 1965. The role of behavior in the ecology and interaction of underyearling coho salmon (*Oncorhynchus kisutch*) and steelhead trout (*Salmo gairdneri*). *Journal of the Fisheries Research Board of Canada* 22: 1035-1081.

- Hartt, A.C. 1980. Juvenile salmonids in the oceanic ecosystem – the critical first summer, Pages 25-57 in W.J. McNeil and D.C. Himsworth, eds. Salmonid ecosystems of the North Pacific. Oregon State University Press, Corvallis.
- Hartt, A.C., and M.B. Dell. 1986. Early oceanic migration and growth of juvenile Pacific salmon and steelhead trout. International North. Pacific Fish. Comm. Bulletin 46. 105 pp.
- Harvey, B.C., and R.J. Nakamoto. 1997. Habitat-dependent interactions between two size-classes of juvenile steelhead in a small stream. Canadian Journal of Fisheries and Aquatic Sciences 54: 27-31.
- Harza Northwest, Inc. 1999. Technical Study Reports. Harza Northwest, Inc.; Cowlitz River Hydroelectric Project, FERC No. 2016. 46pp.
- Hatch, K.M. 1990. A phenotypic comparison of thirty eight steelhead (*Oncorhynchus mykiss*) populations from coastal Oregon. Masters thesis. Oregon State University, Corvallis.
- Hawkins, C.P, M.L. Murphy, N.H. Anderson, and M.A. Wilzbach. 1983. Density of fish and salamanders in relation to riparian canopy and physical habitat in streams of the northwestern United States. Canadian Journal of Fisheries and Aquatic Sciences 40:1173-1185.
- Healey, M.C. 1979. Detritus and juvenile salmon production in the Nanaimo Estuary. I. Production and feeding rates of juvenile chum salmon (*Oncorhynchus keta*). Journal of the Fisheries Research Board of Canada 36:488-496.
- Healey, M.C. 1980. The ecology of juvenile salmon in Georgia Strait, British Columbia, Pages 203-229. in W.J. McNeil and D.C. Himsworth, eds. Salmonid ecosystems of the North Pacific. Oregon State University Press, Corvallis.
- Healey, M.C. 1982. Juvenile pacific salmon in the estuaries: the life support system. in Estuarine Comparisons. Academic Press, Inc., New York.
- Healey, M.C. 1983. Coastwide distribution and ocean migration patterns of stream- and ocean-type Chinook salmon, *Oncorhynchus tshawytscha*. Canadian Field Naturalist 97-427-433.
- Healey, M.C. 1991. Life History of Chinook Salmon. In: C. Groot, and L. Margolis. Pacific Salmon Life Histories. University of British Columbia Press, Vancouver.
- Healey, M.C., and A. Prince. 1995. Scales of variation in life history tactics of Pacific salmon and the conservation of phenotype and genotype. Pages 176-184 In J.L. Nielsen, editor. Evolution and the aquatic ecosystem: defining unique units in population conservation. American Fisheries Society Symposium 17. Bethesda, Maryland.
- Healey, M.C., and C. Groot. 1987. Marine migration and orientation of ocean-type chinook and sockeye salmon, Pages 298-312 in M.J. Dadswell, R.J. Klanda, C.M. Moffitt, R.L. Saunders, R.A. Rulifson, and J.E. Cooper, eds. Common strategies of anadromous and catadromous fishes. American Fisheries Society Symposium 1
- Healey, M.C., and W.R. Heard. 1984. Inter- and intra-population variation in the fecundity of chinook salmon (*Oncorhynchus tshawytscha*) and its relevance to life history theory. Canadian Journal of Fisheries and Aquatic Sciences 41: 476-483.
- Heggenes, J., T.G. Northcote, and A. Peter. 1991a. Spatial stability of cutthroat trout (*Oncorhynchus clarki*) in a small, coastal stream. Canadian Journal of Fisheries and Aquatic Sciences 48:757-762.
- Heggenes, J., T.G. Northcote, and A. Peter. 1991b. Seasonal habitat selection and preferences by cutthroat trout (*Oncorhynchus clarki*) in a small, coastal stream. Canadian Journal of Fisheries and Aquatic Sciences 48:1364-1370.

- Helle, J.H. 1979. Influence of marine environment on age and size at maturity, growth, and abundance of chum salmon, *Oncorhynchus keta* (Walbaum), from Olsen Creek, Prince William Sound, Alaska. Doctoral dissertation. Oregon State University, Corvallis. 118 pp.
- Helle, J.H., and M.S. Hoffman. 1995. Size decline and older age at maturity of two chum salmon (*Oncorhynchus keta*) stocks in western North America, 1972-92. Pages 245-260 in Beamish, R.J., ed. Climate change and northern fish populations. Can. Spec. Publ. of Fisheries and Aquatic Sciences 121.
- Henchman, T.R. 1986. Distribution and food habits of northern pikeminnow, *Ptychocheilus oregonensis* (Richardson), and estimates of their consumption of migrating juvenile salmonids in John Day Reservoir and tailrace during spring and summer, 1983. Master's thesis, University of Washington, Seattle.
- Henry, K.A. 1953. Analysis of factors affecting the production of chum salmon (*Oncorhynchus keta*) in Tillamook Bay. Fish Comm. Oreg. Contrib. 18. 37 pp.
- Heppell, S., H. Caswell, and L. B. Crowder. 2000. Life histories and elasticity patterns: perturbation analysis for species with minimal demographic data. Ecology 81:654-665.
- Hicks, B.J. 1989. The influence of geology and timber harvest of channel morphology and salmonid populations in Oregon Coast range streams. Oregon State University, Corvallis.
- Hilborn, R. and C.J. Walters. 1992. Quantitative Fisheries Stock Assessment. Choice, Dynamics and Uncertainty. Chapman and Hall. International Thomas Publishing, New York.
- Hildebrand and McKenzie 1994. Status of white sturgeon in British Columbia.
- Hildebrand, L, C. McLeod, and S. McKenzie. 1999. Status and management of white sturgeon in the Columbia River in British Columbia, Canada: an overview. Journal of Applied Ichthyology 15:164-172.
- Hillman, T. W., J. S. Griffith, and W. S. Platts. 1987. Summer and winter habitat selection by juvenile Chinook salmon in a highly sedimented Idaho stream. Transactions of the American Fisheries Society 116: 185-195.
- Hillson T., and J. Tipping. 2000. Lewis River Hatchery Complex Fish Biologist Annual Report for 1999. Washington Department of Fish and Wildlife.
- Hiyama, Y., Y. Nose, M. Shimizu, T. Ishihara, H. Abe, R. Sato, M. Takashi, and T. Kajihara. 1972. Predation of chum salmon fry during the course of its seaward migration: II: Otsuchi River investigation 1964 and 1965. Bulletin of the Japanese Soc. Sci. Fish. 38:223-229.
- Hjort, J. 1926. Fluctuations in the year class of important food fishes. Journal du Conseil 1:5-39.
- Hoar, W.S. 1958. The evolution of migratory behavior among juvenile salmon of the genus *Oncorhynchus*. Journal of the Fisheries Research Board of Canada 15:391-428.
- Hollis, G.E. 1975. The effect of urbanization on floods of different recurrence interval. Water Resources Research. 11(3).
- Holmes, H.B. 1952. Loss of salmon fingerlings in passing Bonneville Dam as determined by marking experiments. Unpublished manuscript, U.S. Bureau of Commercial Fisheries Report to U.S. Army Corps of Engineers, North Pacific Division, Portland, Oregon, 52 p. (Available from the U.S. Fish and Wildlife Service, Vancouver, Washington.)

- Holtby, L. B., B. C. Andersen, and R. K. Kadowaki. 1990. Importance of smolt size and early ocean growth to interannual variability in marine survival of coho salmon (*Oncorhynchus kisutch*). Canadian Journal of Fisheries and Aquatic Sciences 47: 2181-2194.
- Houde, E. D. 1987. Fish early life dynamics and recruitment variability. Pages 17-29 in R. D. Hoyt, ed. 10th Annual Larval Fish Conference. American Fisheries Society Symposium 2. Bethesda, Maryland.
- House, R. 1995. Temporal variation in abundance of an isolated population of cutthroat trout in western Oregon, 1981-1991. North American Journal of Fisheries Management 15: 33-41.
- House, R. 1996. Temporal variation in abundance of an isolated population of cutthroat trout in western Oregon, 1981-1991. North American Journal of Fisheries Management 15: 33-41.
- Houston, D.B. 1983. Anadromous fish in Olympic National Park: A status report. Natl. Park Serv. Pacific Northwest Region, 72 p. (Available from Olympic National Park, 600 East Park Avenue, Port Angeles, WA 98362).
- Houston, J.J. 1988. Status of the green sturgeon, *Acipenser medirostris*, in Canada. The Canadian Field Naturalist 102:286-290.
- Howell, P., D. Scarnecchia, et al. 1985. Stock assessment of Columbia River anadromous salmonids. Portland, Oregon, Oregon Department of Fish and Wildlife and Bonneville Power Administration.
- Hubbs, C.L. 1967. Occurrence of the Pacific lamprey, *Entosphenus tridentatus*, off Baja California and in streams of Southern California; with remarks on its nomenclature. Trans. Sand Diego Soc. Natural History 14: 301-312.
- Hubbs, C.L., and R.R. Miller. 1948. The Great Basin. II. The zoological evidence. Bulletin of the University of Utah 38(20): 17-144.
- Hunter, J.G. 1959. Survival and production of pink and chum salmon in a coastal stream. Journal of the Fisheries Research Board of Canada 16:835-886.
- Huzyk, L. and H. Tsuyuki. 1974. Distribution of LDH-B geen in resident and anadromous rainbow trout (*Salmo gairdneri*) from streams in British Columbia. Journal of the Fisheries Research Board of Canada 31:106-108.
- Hymer, J., R. Pettit, M. Wastel, P. Hahn, K. Hatch. 1992. Stock summary reports for Columbia River anadromous salmonids, Volume III: Washington subbasins below McNary Dam. Bonneville Power Administration, Portland, Oregon.
- Hynes, H. B. N. 1970. The ecology of running waters. Liverpool University Press, Liverpool, England.
- IPC (Idaho Power Company). 2003. Draft Snake River White Sturgeon Conservation Plan. Idaho Power Company, Boise. 260 pp.
- Ireland, S.C., P.J. Anders, and J.T. Siple. 2002. Conservation aquaculture: An adaptive approach to prevent extinction of an endangered white sturgeon population (*Acipenser transmontanus*). Pages 211-222 in W. VanWinkle, P. Anders, D. Dixon, and D. Secor, eds. Biology, Management and Protection of North American Sturgeons. American Fisheries Society Symposium 28.
- Irvine, J.R. and B.R. Ward (1989). Patterns of timing and size of wild coho salmon (*Oncorhynchus kisutch*) smolts migrating from the Keogh River Watershed on northern Vancouver Island. Canadian Journal of Fisheries and Aquatic Sciences 46(7): 1086-1094.

- ISAB (Independent Scientific Advisory Board). 2003. ISAB Review of Salmon and Steelhead Supplementation. ISAB 2003-3 Supplementation Report, Portland.
- IUCN (International Union for the Conservation of Nature). 1994. Red list categories and criteria. Cambridge, UK.
- Iwamoto, R.N. and J.G. Williams. 1993. Juvenile salmonids passage and survival through turbines. Report to U.S. Army Corps of Engineers, Portland, OR, Contract E86920049, 27 p. (Available from Northwest Fisheries Science Center, 2725 Montlake Blvd. E., Seattle, WA 98112-2097.)
- Iwata, M. 1982. Downstream migration and seawater adaptability of chum salmon (*Oncorhynchus keta*) fry, Pages 51-59 in B.R. Melteff and R.A. Neve, eds. Proceedings of the North Pacific Aquaculture Symposium. Alaska Sea Grant Report 82-2.
- Jackson, A. D., D. R. Hatch, B. L Parker, M. S. Fitzpatrick, and H. Li. 1997. Research and Restoration Annual Report. U.S. Department of Energy, Bonneville Power Administration 1997:91. Portland, Oregon.
- Jager, H. I., J. A Chandler, K. B. LePla, and W. Van Winkle. 2001 A theoretical study of river fragmentation by dams and its effects on white sturgeon. *Environmental Biology of Fishes* 60:347-361.
- Jager, H. I., K. LePla, J. Chandler, P. Bates, W. VanWinkle. 2000. Population viability analysis of white sturgeon and other riverine fishes. *Environmental Science and Policy* 3:S483-S489.
- Jager, H. I., VanWinkle, W., J. A. Chandler, K. B. LePla, P. Bates, and T. D. Counihan. 2002. A simulation study of factors controlling white sturgeon recruitment in the Snake River. Pages 127-150 in W. VanWinkle, P. Anders, D. Dixon, and D. Secor, eds. *Biology, Management and Protection of North American Sturgeons*. American Fisheries Society Symposium 28.
- Jameson, R.J., and K.W. Kenyon. 1977. Prey of sea lions in the Rogue River, Oregon. *Journal of Mammalogy* 58(4): 672.
- Jensen, H.M. 1956. Recoveries of immature chum salmon tagged in southern Puget Sound. *Wash. Dep. Fish. Fish. Res. Pap.* 1(4):32
- Jin, F. 1995. Late Mesozoic acipenseriformes (*Osteichthyes: Actinopterygii*) in Central Asia and their biogeographical implications. Pages 15-21 in A. Sun and Y. Yang, eds., *Sixth Symposium on Mesozoic Terrestrial Ecosystems and Biota*, Short papers, China Ocean Press, Beijing.
- Johnson, D.L., J.D. Rodgers, and M.F. Solazzi. 1991. Development and evaluation of techniques to rehabilitate Oregon's wild salmonids. Oregon Department of Fish and Wildlife, Portland.
- Johnson, O.W., M.H. Ruckelshaus, W.S. Grant, F.W. Waknitz, A.M. Garrett, G.J. Bryant, K. Neely, and J.J. Hard. 1999. Status review of the coastal cutthroat trout from Washington, Oregon, and California. NOAA Technical Memorandum. NMFS-NWFSC-37. 292 pp.
- Johnson, O.W., M.H. Ruckelshaus, W.S. Grant, F.W. Waknitz, A.M. Garrett, G.J. Bryant, K. Neely, and J.J. Hard. 1999. Status Review of Coastal Cutthroat Trout From Washington, Oregon, and California. National Marine Fisheries Service, Conservation Biology Division, NOAA Technical Memorandum NMFS-NWFSC-37. Seattle, Washington.
- Johnson, O.W., W.S. Grant, R.G. Kope, K. Neely, F.W. Waknitz, and F.S. Waples. 1997. Status review of chum salmon from Washington, Oregon and California. NOAA Technical Memorandum NMFS-NWFSC-32. 280pp.

- Johnson, R.C., R.J. Gerke, D.W. Heiser, R.F. Orrell, S.B. Mathews, and J.G. Olds. 1971. Pink and chum salmon investigations, 1969: supplementary progress report. Washington Department of Fisheries, Fisheries Management and Research Division, Olympia. 66 pp.
- Johnson, S.L., M.F. Solazzi, and J.D. Rodgers. 1993. Development and evaluation of techniques to rehabilitate Oregon's wild salmonids. Oregon Department of Fish and Wildlife, Portland.
- Johnson, T.H. 1985. Density of steelhead parr for mainstem rivers in western Washington during the low flow period, 1984. Washington Department of Game, Snow Creek Research Station, Port Townsend. 29 pp.
- Johnston, J.M. 1982. Life histories of anadromous cutthroat with emphasis on migratory behavior. Pages 13-127 in E.L. Brannon and E.O. Salo, eds. Proceedings of the salmon and trout migratory behavior symposium. University of Washington, Seattle.
- Johnston, S. and P. Nealson 1999. Hydroacoustic smolt passage and spill effectiveness studies at John Day Dam in 1999. In Abstracts, Anadromous Fish Evaluations Program 1999 Annual Research Review, November 1999. (Available from U.S. Army Corps of Engineers, Walla Walla District, Walla Walla, Washington.)
- Jones, D.E. 1973. Steelhead and sea-run cutthroat trout life history in Southeast Alaska. Alaska Department of Fish and Game. Annual Progress Report 14, Project AFS-42 (AFS-42-1), 18 pp.
- Jones, D.E. 1974. The study of cutthroat-steelhead in Alaska. Alaska Department of Fish and Game. Annual Progress Report 15, Project AFS-42 (AFS-42-2):15-31.
- Jones, D.E. 1975. Steelhead and sea-run cutthroat trout life history study in Southeast Alaska. Alaska Department of Fish and Game Annual. Progress Report 16, Project AFS-42 (AFS-42-3-B):23-55.
- Jonhson, T.H. 1985. Density of steelhead parr for mainstem rivers in western Washington during the low flow period, 1984. Washington State Game Department, Fisheries Management Division, No. 85-6.
- June, J.A. 1981. Life history and habitat utilization of cutthroat trout (*Salmo clarki*) in a headwater stream on the Olympic Peninsula, Washington. Master's thesis, University of Washington, Seattle. 112 pp.
- Kaeriyama, M. 1989. Aspects of salmon ranching in Japan. *Physiol. Ecol. Japan.* 1:625-638.
- Kan, T.T. 1975. Systematics, Variation, Distribution and Biology of Lampreys of the genus *Lamptera* in Oregon. Doctoral dissertation, Oregon State University, Corvallis.
- Kareiva, P., M. Marvier, and M. McClure. 2000. Recovery and management options for spring/summer chinook salmon in the Columiba River Basin. *Science* 290:977-979.
- Keefer, M.L., and T.C. Bjornn 1999. Evaluation of adult salmon and steelhead migrations past dams and through reservoirs in the Columbia River Basin. Handout presented at the U.S. Army Corps of Engineers Anadromous Fish Evaluation Program Annual Review, November 14-18, 1999, Walla Walla, Washington, 10 pp. (Available from U.S. Army Corps of Engineers, Walla Walla, WA.)
- Keller, E.A. and F.J. Swanson. 1979. Effects of large organic material on channel form and fluvial processes. *Earth Surface Processes* 4:361-380.
- Keller, K. 1999. 1998 Columbia River chum return. Washington Department of Fish and Wildlife, Columbia River Progress Report 99-8, 53 pp.
- Keller, K. 2001. 1999 Columbia River Chum Return. Columbia River Progress Report 2000-6. Washington Department of Fish and Wildlife.

- Kempinger, J.J. (1996). Habitat, growth, and food of young lake sturgeons in the Lake Winnebago system, Wisconsin. *North American Journal of Fisheries Management* 16(1): 102-114.
- Kincaid, H. 1993. Breeding plan to preserve the genetic variability of the Kootenai River white sturgeon. Final Report to Bonneville Power Administration, U.S. Fish and Wildlife Service. Project 93-27. Contract Number DE-A179-93B002886. Portland, Oregon.
- Kirn, R.A., R.D. Ledgerwood, and R.A. Nelson. 1986. Increased abundance and the food consumption of northern pikeminnow (*Ptychocheilus oregonensis*) at River Kilometer 75 in the Columbia River. *Northwest Science* 60:197-200.
- Kjelson, M.A., P.F. Raquel, and F.W. Fisher. 1982. Life history of fall-run juvenile chinook salmon, *Oncorhynchus tshawytscha*, in the Sacramento-San Joaquin estuary, California, p. 393-411. in V.S. Kennedy, ed. *Estuarine comparisons*. Academic Press, Inc. New York.
- Kline, T. J., J. J. Goering, O. A. Mathisen, and P. H. Poe. 1993. Recycling of elements transported upstream by runs of Pacific salmon: II. 15N and 13C evidence in the Kivchak River watershed, Bristol Bay, Southwestern Alaska. *Canadian Journal of Fisheries and Aquatic Sciences* 50: 2350-2365.
- Klyashtorin, L. B. 1976. The sensitivity of young sturgeons to oxygen deficiency. *Journal of Ichthyology* 16:677-682.
- Knighton, D. 1998. *Fluvial Forms and Processes*. John Wiley and Sons, New York.
- Knudsen, E.E. 2002. Ecological perspectives on Pacific salmon: can we sustain biodiversity and fisheries? Pages 277-320 In K.D. Lynch, M.L. Jones, and W.W. Taylor, editors. *Sustaining North American salmon: Perspectives across regions and disciplines*. American Fisheries Society, Bethesda, Maryland.
- Knudsen, E.E., and S.J. Dilley. 1987. Effects of riprap bank reinforcement on juvenile salmonids in four western Washington streams. *North American Journal of Fisheries Management* 7:351-356.
- Knutsen, C.J., and D.L. Ward. 1999. Biological characteristics of northern pikeminnow in the lower Columbia and Snake rivers before and after sustained exploitation. *Transactions of the American Fisheries Society* 128:1008-1019.
- Kohlhorst, D. W. 1976. Sturgeon spawning in the Sacramento River in 1973 as determined by distribution of larvae. *California Fish and Game* 62(1):32-40.
- Kondolf, M. G. 2000. Assessing salmonid spawning gravel quality *Transactions of the American Fisheries Society* 129: 262-281.
- Koonce, J.F., T.B. Bagenal, et al. 1977. Factors influencing year-class strength of percids: a summary and a model of temperature effects. *Journal of the Fisheries Research Board of Canada* 34(10): 1900-1909.
- Koski, K.V. 1966. The survival of coho salmon (*Oncorhynchus kisutch*) from egg deposition to emergence in three Oregon coastal streams. Master's thesis. Oregon State University, Corvallis. 84 pp.
- Koski, K.V. 1975. The survival and fitness of two stocks of chum salmon (*Oncorhynchus keta*) from egg deposition to emergence in a controlled-stream environment at Big Beef Creek. Doctoral dissertation. University of Washington, Seattle. 212 pp.
- Koslow, J.A., A.J. Hobday and G.W. Boehlert. 2002. Climate variability and marine survival of coho salmon (*Oncorhynchus kisutch*) in the Oregon production area. *Fisheries Oceanography* 11(2): 65-77

- Kostarev, V.L. 1970. Quantitative calculation of Okhotsk keta juveniles. Izv. Tikhookean. Nauchno-Issled. Inst. Rybn. Khoz. Okeanogr. 71:145-158. (Transl. from Russian; Fish. Res. Board Can. Transl. Ser. 2589)
- Kostow, K. 1995. Biennial Report on the Status of Wild Fish in Oregon. Oregon Dept. Fish and Wild., Portland. 217 pp. + app.
- Kostow, K. 2002. Oregon Lampreys: Natural history status and analysis management issues. Oregon Department of Fish and Wildlife.
- Kostow, K. E., A. R. Marshall, and S. R. Phelps. 2003. Naturally spawning hatchery steelhead contribute to smolt production but experience low reproductive success. Transactions of the American Fisheries Society 132:780-790.
- Kruse, G.H. 1998. Salmon run failures in 1997-1998: A link to anomalous ocean conditions? Alaska Fishery Research Bulletin 5(1): 55-63.
- Kwain, W.H. 1975. Embryonic development, early growth, and meristic variation in rainbow trout (*Salmo gairdneri*) exposed to combinations of light intensity and temperature. Journal of the Fisheries Research Board of Canada 32: 397-402.
- Kyle, G.B. 1992. Summary of Sockeye salmon (*Oncorhynchus nerka*). Pages 419-434 in H.D. Smith, L. Margolis, and C.C. Wood, eds. Sockeye salmon (*Oncorhynchus nerka*) populations biology and future management. Canadian Special Publications in Fisheries and Aquatic Sciences 96.
- Kynard, B. 1997. Life history, latitudinal patterns, and status of shortnose sturgeon, *Acipenser brevirostrum*. Environmental Biology of Fishes 48:319-334.
- Lande, R. 1993. Risks of population extinction from demographic and environmental stochasticity and random catastrophes. American Naturalist 142: 911-927.
- Lande, R., and G.F. Barrowclough. 1987. Effective population size, genetic variation, and their use in population management in M.E. Soule, eds. Viable Populations for Conservation. Cambridge University, New York.
- Lane, E. D. 1991. Status of white sturgeon, *Acipenser transmontanus*, in Canada. Canadian Field-Naturalist 105(2):161-168.
- LaPatra, S. E., G. R. Jones, K. A. Lauda, T. S. McDowell, R. Schneider, R. P. Hedrick. 1995. White sturgeon as a potential vector of infectious hematopoietic necrosis virus. Journal of Aquatic Animal Health 7:225-230.
- LaPatra, S.E., J.M Groff, G.R. Jones, B. Munn, T.L. Patterson, R.A. Holt, A.K. Hauck, and R.P. Hedrick. 1994. Occurrence of white sturgeon iridovirus infections among cultured white sturgeon in the Pacific Northwest. Aquaculture 126:201-210.
- Larkin, P.A. 1977. Pacific salmon, Pages 156-186 in J.A. Gulland, ed. Fish population dynamics. John Wiley and Sons, New York.
- Larkins, H.A. 1964. Some epipelagic fishes of the North Pacific Ocean, Bering Sea, and Gulf of Alaska. Transactions of the American Fisheries Society 93: 286-290.
- Larsson, P. O. 1980. Predation on migrating smolt as a regulating factor of Baltic salmon, *Salmo salar L.*, populations. Journal of Fish Biology 26: 391-397.
- Laufle, J.C., G.B. Pauley, and M.F. Shepard. 1986. Species profiles: Life histories and environmental requirements of coastal fishes and invertebrates (Pacific Northwest): coho salmon. U.S. Fish and Wildlife Service Biological Report 82(11.48), 18 pp.

- Lavier, D. 1959. Progress report for 1959. Washington Department of Game, Olympia.
- Lawson, P.W. 1993. Cycles in ocean productivity, trends in habitat quality, and the restoration of salmon runs in Oregon. *Fisheries* 18(8):6-10.
- LCFRB (Lower Columbia River Fish Recovery Board). 2003. Draft Technical Foundation. June 2003.
- LCSCI (Lower Columbia Steelhead Conservation Initiative). 1998. State of Washington, Olympia.
- Lee, D.S., C.R. Gilbert, C.H. Hocutt, R.E. Jenkins, D.E. McAllister, J.R. Stauffer, Jr. 1981. Atlas of North American Freshwater Fishes. North Carolina State Museum of Natural History.
- Leider, S.A. 1997. Status of sea-run cutthroat trout in Washington. Pages 68-76 in J.D. Hal, P.A. Bisson and R.E. Gresswell, eds. Sea-run cutthroat trout: biology, management, and future conservation. American Fisheries Society, Corvallis, Oregon.
- Leider, S.A., M.W. Chilcote, and J.J. Loch. 1986. Movement and survival of presmolt steelhead in a tributary and the main stem of a Washington River. *North American Journal of Fisheries Management* 6: 526-531.
- Leider, S.A., P.L. Hulett, J.J. Loch, and M.W. Chilcote. 1990. Electrophoretic comparison of the reproductive success of naturally spawning transplanted and wild steelhead trout through the returning adult stage. *Aquaculture* 88: 239-252.
- Leider, S.A., S.R. Phelps, and P.L. Hulett. 1995. Genetic analysis of Washington steelhead: Implications for revision of genetic conservation management units. Washington Department of Fish and Wildlife Program Report. 21 pp.
- Leidy, R.A., and G. R. Leidy. 1984. Life stage periodicities of anadromous salmonids in the Klamath River Basin, northwestern California. U.S. Fish and Wildlife Service, Sacramento, California, 38 p.
- Leopold, L.B. 1968. Hydrology for Urban Land Planning – A Guidebook on the Hydrologic Effects of Urban Land Use. Geological Survey Circular 554.
- Levanidov, V. Ya. 1964. Salmon population trends in the Amur basin and means of maintaining the stocks, Pages 35-40 in E.N. Pavlovskii, ed. Lososevoe khozyaistvo dal'nego vostoka. Izdatel'stvo Nauka, Moscow. (Transl. from Russian; Univ. Wash. Fish. Res. Inst. Circ. 227).
- Levings, C.D. 1982. Short term use of a low tide refuge in a sandflat by juvenile chinook, *Oncorhynchus tshawytscha*, Fraser River estuary. *Can. Tech. Rep. of Fisheries and Aquatic Sciences* 1111:33 p.
- Levings, C.D., C.D. McAllister, and B.D. Chang. 1986. Differential use of the Campbell River estuary, British Columbia, by wild and hatchery-reared juvenile Chinook salmon (*Oncorhynchus tshawytscha*). *Canadian Journal of Fisheries and Aquatic Sciences* 43:1386-1397.
- Levy, D.A., and T.G. Northcote. 1981. The distribution and abundance of juvenile salmon in marsh habitats of the Fraser River Estuary. Westwater Research Center University of British Columbia Technical Report 25. 117 pp.
- Levy, D.A., and T.G. Northcote. 1982. Juvenile salmon residency in a marsh area of the Fraser River estuary. *Canadian Journal of Fisheries and Aquatic Sciences* 39:270-276.
- Levy, D.A., T.G. Northcote, and G.J. Birch. 1979. Juvenile salmon utilization of tidal channels in the Fraser River estuary, British Columbia. Westwater Research Center University of British Columbia Technical Report 23. 70 pp.
- Lichatowich, J., L. Mobrand, L. Lestelle, and T. Vogel. 1995. An approach to the diagnosis and treatment of depleted Pacific salmon populations in Pacific Northwest watersheds. *Fisheries* 20(1):10-18.

- Lichatowich, J.A. and J.D. McIntyre 1987. Use of hatcheries in the management of Pacific anadromous salmonids. American Fisheries Society Symposium 1: 131-136.
- Lichatowich, J.A. 1999. Salmon without rivers. Island Press, Washington, D.C. 317 pp.
- Light, J.T. 1987. Coastwide abundance of North American steelhead trout. Fisheries Research Institute Report FRI-UW-8710. University of Washington, Seattle. 18 pp.
- Light, J.T., S. Fowler, M.L. Dahlberg, M.L. 1988. High seas distribution of North American steelhead as evidenced by recoveries of marked or tagged fish. University of Washington, Fisheries Research Institute, FRI-UW-8816.
- Liscom, K.L., G.E. Monan, and L. Stuehrenberg. 1979. Radio tracking studies relating to fallback at hydroelectric dams on the Columbia and Snake rivers. Pages 39-53 in Fifth Progress Report on Fisheries Engineering Research Program 1973-78. U.S. Army Corps of Engineers, North Pacific Division, Portland, Oregon.
- Lister, D.B., and C.E. Walker. 1966. The effect of flow control on freshwater survival of chum, coho, and chinook salmon in the Big Qualicum River. Canadian Fish. Cult. 37:3-25.
- Lister, D.B., C.E. Walker, and M.A. Giles. 1971. Cowichan River Chinook salmon escapements and juvenile production 1965-1967. Fish. Serv. (Can.) Pac. Reg. Tech. Rep. 1971-3. 8 pp.
- Lister, D.B., D.G. Hickey, and I. Wallace. 1981. Review of the effects of enhancement strategies on the homing, straying and survival of Pacific salmonids. Prepared for Department of Fisheries and Oceans, Salmonid Enhancement Program. D.B. Lister and Associates, West Vancouver, British Columbia. 51 pp.
- Loch, J.L., M.W. Chilcote, and S.A. Leider. 1985. Kalama River studies final report, Part II. Juvenile downstream migrant studies. Washington Dept. of Fish and Game, Fisheries Management Division, Report 85-12.
- Logerwell, L.E., N. Mantua, P. Lawson, R.C. Francis, V. Agostini. 2002. Tracking environmental processes in the coastal zone for understanding and predicting Oregon coho marine survival. In review: Fisheries Oceanography.
- Long, J.A. 1995. The rise of fishes. Johns Hopkins University Press, Baltimore, Maryland.
- Lowry, G.R. 1965. Movement of cutthroat trout (*Salmo clarki clarki* Richardson) in three Oregon coastal streams. Transactions of the American Fisheries Society 94(4):334-338.
- Lynch, M. 1990. Mutation load and the survival of small populations. Evolution 44:1725-1737.
- Lynch, M. 1996. A quantitative-genetic perspective on conservation issues. In Avise, J.C. and J.L. Hamrick, eds. Conservation Genetics. Case Histories from Nature. Chapman and Hall, New York.
- Lynch, M., and M. O'Hely. 2001. Captive breeding and the genetic fitness of natural populations. Conservation Genetics 2:363-378.
- Magee, J.P., T.E. McMahon, and R.F. Thurow. 1996. Spatial variation in spawning habitat of cutthroat trout in a sediment-rich stream basin. Transactions of the American Fisheries Society 125: 768-779.
- Major, R.L., J. Ito, S. Ito, and H. Godfrey, 1978. Distribution and abundance of chinook salmon (*Oncorhynchus tshawytscha*) in offshore waters of the north Pacific Ocean. North Pacific International Fisheries Commission Bulletin 38.
- Marcot, B. G., W. E. McConaha, P. H. Whitney, T. A. O'Neil, P. J. Paquet, L. Moberand, G. R. Blair, L. C. Lestelle, K. M. Malone, and K. I. Jenkins. 2002. A multi-species framework approach for the

- Columbia River Basin: integrating fish, wildlife, and ecological functions. Northwest Power Planning Council, Portland, Oregon. CD-ROM and Web (www.edthome.org/framework).
- Marr, J.C. 1943. Age, length and weight studies of three species of Columbia River salmon (*Oncorhynchus keta*, *O. gorbuscha* and *O. kisutch*). Stanford Ichthyo. Bulletin 2:157-197
- Marshall, A.R., C. Smith, et al. 1995. Genetic diversity units and major ancestral lineages for Chinook salmon in Washington. C. Busack and J.B. Shaklee. Washington Department of Fish and Wildlife, Olympia.
- Martin, D.J., D.R. Glass, C.J. Whitmus, C.A. Simenstad, D.A. Milward, E.C. Volk, M.L. Stevenson, P. Nunes, M. Savvoie, and R.A. Grotefendt. 1986. Distribution, seasonal abundance, and feeding dependencies of juvenile salmon and non-salmonid fishes in the Yukon River Delta. NOAA OCSEAP Final Rep. 55(1988):381-770.
- Martin, S.W., M.A. Schuck, K. Underwood and A.T. Scholtz. 1992. Investigations of bull trout (*Salvelinus confluentus*), steelhead trout (*Oncorhynchus mykiss*), and spring chinook (*O. tshawytscha*) interactions in southeast Washington streams. Project No. 90-53. Contract No. DE-B179-91BP17758 for U.S. Department of Energy, Bonneville Power Administration, Division of Fish and Wildlife, P.O. Box 3621, Portland, Oregon, 97208-3621.
- Martinelli, T.L., and R.S. Shively. 1997. Seasonal distribution, movements and habitat associations of northern pikeminnow in two lower Columbia River reservoirs. Regulated Rivers: Research and Management 13:543-556.
- Martinson, R.D., G.M. Kovalchuk, R.B. Mills, and J.W. Kamps. 1998. Monitoring of downstream salmon and steelhead at federal hydroelectric facilities – 1997. Report to Bonneville Power Administration, Portland, Oregon. Contract DE-AI79-85BP20733, 64 pp. + App.
- Mason, J.C. 1974. Behavioral ecology of chum salmon fry (*Oncorhynchus keta*) in a small estuary. Journal of the Fisheries Research Board of Canada 31:83-92
- Mathews, S.B., and H.B. Senn. 1975. Chum salmon hatchery rearing in Japan, in Washington. Washington Sea Grant Publication WSG-TA 75-3:24 p.
- Matthews, G. M. and R. Waples, S. 1991. Status review for Snake River spring and summer chinook salmon. Seattle, National Marine Fisheries Service Northwest Fisheries Science Center.
- Mattson, C.R. 1949. The lamprey fishery at Willamette Falls, Oregon. in Fish Commission of Oregon Research Briefs. 2(2): 23-27.
- May, R. C. 1974. Larval mortality in marine fishes and the critical period concept. in J.H.S. Blaxter, ed. The early life history of fish. Springer-Verlag, New York.
- Mazer, J.I., and M.P. Shepard. 1962. Marine survival, distribution and migration of pink salmon off the British Columbia coast. H.R. MacMillan Lectures in Fisheries, University of British Columbia, Vancouver. p. 113-121.
- McCabe, G. T., Jr. 1993. Prevalence of the parasite *Cystoopsis acipenseris* (Nematoda) in juvenile white sturgeons in the lower Columbia River. Journal of Aquatic Animal Health 5:313-316.
- McCabe, G. T., Jr., R. L. Emmett, and S. A. Hinton. 1993. Feeding ecology of juvenile white sturgeon (*Acipenser transmontanus*) in the lower Columbia River. Northwest Science 63:170-180.
- McCabe, G. T., Jr., W. D. Muir, et al. 1983. Interrelations between juvenile salmonids and nonsalmonid fish in the Columbia River estuary. Fish Bulletin 81.
- McCuen, R.H. 1998. Hydrologic Analysis and Design. Prentice-Hall. Upper Saddle River, New Jersey.

- McElhany, P., and 11 coauthors. 2003. Interim report on viability criteria for Willamette and Lower Columbia Basin Pacific Salmonids. Willamette/Lower Columbia Technical Recovery Team Interim Report. NOAA Fisheries, Portland.
- McElhany, P., M.H. Ruckelshaus, M.J. Ford, T.C. Wainwright, and E.P. Bjorkstedt. 2000. Viable salmonid populations and the recovery of evolutionarily significant units. NOAA Technical Memorandum NMFS-NWFSC-42.
- McHenry, E.T. 1981. Coho salmon studies in the Resurrection Bay area. Annual Progress Report, Alaska Department of Fish and Game.
- McKinnell, S.M., C.C. Wood, D.T. Rutherford, K.D. Hyatt, and D.W. Welch. 2001. The demise of Owikeno Lake sockeye salmon. North American Journal of Fisheries Management 21: 774-791.
- McMahon, T.E. 1983. Habitat suitability models: coho salmon. U.S. Fish and Wildlife Service FWS/OBS 82/10.49. 29 pp.
- McMichael, G. A., T. N. Pearsons, and S. A. Leider. 2000. Minimizing ecological impacts of hatchery-reared juvenile steelhead trout on wild salmonids in a Yakima Basin watershed. Pages 365-380 in E.E. Knudsen, C. S. Steward, D. D. MacDonald, J. E. Williams, and D. W. Reiser, editors. Sustainable fisheries management: Pacific salmon, Lewis Publishers, Boca Raton, Florida.
- McMichael, G.A., C.S. Sharpe, and T.N. Pearsons. 1997. Effects of residual hatchery-reared steelhead on growth of wild rainbow trout and spring Chinook salmon. Transactions of the American Fisheries Society 126:230-239.
- McNair, M. 1996. Alaska fisheries enhancement program. 1995 annual report. Reg. Info. Rep. 5J96-08, 43 p. Alaska Department of Fish and Game, Fairbanks.
- McNeil, W.J. 1969. Survival of pink and chum salmon eggs and alevins. Pages 101-117 in Northcote, T.G., ed. Salmon and trout in streams. H.R. McMillan Lectures in Fisheries, University of British Columbia, Vancouver.
- McPhail, J.D. and C. Murray. 1979. The early life history and ecology of Dolly Varden (*Salvelinus malmo*) in the upper Arrow Lakes. Report to the British Columbia Hydro and Power Authority and Kootney Department of Fish and Wildlife.
- McPhail, J.D. and C.C. Lindsey. 1970. Freshwater fishes of northwestern Canada and Alaska. Bulletin of the Fisheries Research Board of Canada 173. 381pp.
- McPhail, J.D., and J.S. Baxter. 1996. A review of bull trout (*Salvelinus confluentus*) life-history and habitat use in relation to compensation and improvement opportunities. Department of Zoology, University of British Columbia. Fisheries Management Report 104. Vancouver.
- Meehan, W.R., F.J. Swanson, and J.R. Sedell. 1977. Influences of riparian vegetation on aquatic ecosystems with particular reference to salmonids fishes and their food supply. Pages 137-145 in R.R. Johnson and D.A. Jones, eds. Importance, Preservation and Management of Riparian Habitat: A Symposium held at Tucson, Arizona, July 9, 1977. U.S. Forest Service General Technical Report RM-43.
- Meehan, William R. 1996. Influence of riparian canopy on macroinvertebrate composition and food habits of juvenile salmonids in several Oregon streams. Research Paper PNW-RP-496. Portland, Oregon: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 14 p.

- Meeuwig, M.J., Bayer, J., Seelye, R., Reiche. 2002. Identification of Larval Pacific Lampreys (*Lampetra tridentate*), River Lampreys (*L. ayresi*), and Western Brook Lampreys (*L. richardsoni*) and Thermal Requirements of Early Life History Stages of Lampreys.
- Mesa, M. G., T. P. Poe, A. G. Maule, and C.B. Schreck. 1998. Vulnerability to predation and physiological stress response in juvenile chinook salmon (*Oncorhynchus tshawytscha*) experimentally infected with *Renibacterium salmoninarum*. Canadian Journal of Fisheries and Aquatic Sciences 55:1599-1606
- Mesa, M.G., and T.M. Olson. 1993. Prolonged swimming performance of northern pikeminnow. Transactions of the American Fisheries Society 122:1104-1110.
- Michael, J.H. 1980. Repeat spawning of Pacific Lamprey. California Fish and Game Notes. 187 pp.
- Migdalski, E. C. 1962. Anglers guide to the freshwater sport fishes. Ronald Press, New York.
- Miller, A. I., and L. G. Beckman. 1992. Age and growth of juvenile white sturgeon in the Columbia River downstream from McNary Dam. in R.C. Beamesderfer and A.A. Nigro, eds. Status and habitat requirements of the white sturgeon populations in the Columbia River downstream from McNary Dam, Volume II. Final report (Contract DE-AI79-86BP63584) to Bonneville Power Administration, Portland, Oregon.
- Miller, A. I., and L. G. Beckman. 1996. First record of predation on white sturgeon eggs by sympatric fishes. Transactions of the American Fisheries Society 125:338-340.
- Miller, B.A. and S. Sadro. 2003. Residence time and seasonal movements of juvenile coho salmon in the ecotone and lower estuary of Winchester Creek, South Slough, Oregon. Transactions of the American Fisheries Society 132:546-559.
- Miller, R.J., and E.L. Brannon. 1982. The origin and development of life history patterns in Pacific salmonids, Pages 296-309 in E.L. Brannon and E.O. Salo, eds. Proceedings of the Salmon and Trout Migratory Behavior Symposium. School of Fisheries, University of Washington, Seattle.
- Miller, R.R. 1965. Quaternary freshwater fishes of North America. Pages 569-581 in The Quaternary of the United States. Princeton University Press, Princeton, New Jersey.
- Milne, D.J. 1964. The chinook and coho salmon fisheries of British Columbia; with appendix by H. Godfrey. Bulletin of the Fisheries Research Board of Canada 142. 46 pp.
- Mobrand Biometrics. 1999. Application of the ecosystem diagnostic and treatment method (EDT) to analyze fish resources in the Cowlitz watershed in support of FERC relicensing process. Draft report Vol. 1. Prepared for the Resource Planning Group of the Cowlitz River FERC Relicensing Process. June, 1999.
- Mobrand, L.E., J.A. Lichatowich, L.C. Lestelle, and T.S. Vogel. 1997. An approach to describing ecosystem performance "through the eyes of salmon". Canadian Journal of Fisheries and Aquatic Sciences 54: 2964-2973.
- Moffett, J.W., and S.H. Smith. 1950. Biological investigations of the fishery resource of Trinity River, California. U.S. Fish and Wildlife. Service. Special Scientific Report Fisheries 12: 70 pp.
- Montgomery, D.R, T.B. Abbe, J.M. Buffington, N.P. Peterson, K.M. Schmidt, and J.D. Stock. 1996. Distribution of bedrock and alluvial channels in forested mountain drainage basins. Letters To Nature. Nature vol. 381.
- Montgomery, D.R. and J.M. Buffington. 1998. Channel Processes, Classification, and Response. Chapter 2 in River Ecology and Management – Lessons from the Pacific Coastal Ecoregion. Springer-Verlag, New York.

- Montgomery, D.R., E.M. Beamer, G.R. Pess, T.P. Quinn. 1999. Channel type and salmonid spawning distribution and abundance. *Canadian Journal of Fisheries and Aquatic Sciences* 56: 377-387.
- Montgomery, D.R., J.M. Buffington, N.P. Peterson, D. Schuett-Hames, and T.P. Quinn. 1996. Stream-bed scour, egg burial depths, and the influence of salmonid spawning on bed surface mobility and embryo survival. *Canadian Journal of Fisheries and Aquatic Sciences* 53: 1061-1070.
- Moore, K.M.S., and S.V. Gregory. 1988. Summer Habitat Utilization and Ecology of Cutthroat (*Salmo clarki*) in Cascade Mountain Streams. *Canadian Journal of Fisheries and Aquatic Sciences* 45:1921-1930.
- Moring, J.R., and R.L. Youker. 1979. Oregon rainbow and cutthroat trout evaluation. Pages 65. Oregon Department of Fish and Wildlife.
- Moser, M.L., P.A. Ocker, L.C. Stuehrenberg, and T.C. Bjornn. 2002. Passage Efficiency of Adult Pacific Lampreys at Hydropower Dams on the Lower Columbia River, USA. *Transactions of the American Fisheries Society* 131: 956-965.
- Moyle, P.B. 1976. *Inland Fishes of California*. University of California Press, Berkeley. 405 pp.
- Moyle, P.B., and J.J. Cech, Jr. 1982. *Fishes: An Introduction to Ichthyology*. Prentice-Hall Publishers, Inc. Englewood Cliffs, New Jersey.
- Moyle, P.B., R.M. Yoshiyama, J.E. Williams, and E.D. Wikramanayake. 1995. Green sturgeon. Pages 26-34 in *Fish species of Special Concern in California, 2nd ed.* Final Report to the Department of Fish and Game (Contract 2128IF).
- Muir, W.D., G.T. McCabe, Jr., M.J. Parsley, and S.A. Hinton. 2000. Diet of first feeding larval and young-of-the-year white sturgeon in the lower Columbia River. *Northwest Science* 74:25-33.
- Mullan, J.W., K.R. Williams, G. Rhodus, T.W. Hillman, and J.D. McIntyre. 1992. Production and habitat of salmonids in mid-Columbia River tributary streams. Monograph I, U.S. Fish and Wildlife Service, Leavenworth, Washington.
- Mundie, J.H. 1969. Ecological implications of the diet of juvenile coho in streams, Pages 135-152. in T.G. Northcote, ed. *Symposium on Salmon and Trout in streams*. H.R. MacMillan Lectures in Fisheries. Institute of Fisheries, University of British Columbia, Vancouver.
- Murphy, M.L. and W.R. Meehan. 1991. Stream Ecosystems. Chapter 2 in W.R. Meehan, ed. *Influences of Forest and Rangeland Management on Salmonid Fishes and Their Habitats*, American Fisheries Society Special Publication 19:17-46.
- Murray, C.B. 1980. Some effects of temperature on zygote and alevin, rate of development of five species of Pacific salmon (*Oncorhynchus*) embryos and alevins. *Canadian Journal of Zoology* 66:266-273.
- Mushens, C.J. and J.R. Post. 2000. Population dynamics of the Lower Kananaskis lake bull trout: 1999 progress report. Alberta Conservation Association and transAlta Utilities, Calgary, Alberta.
- Musick, J.A., and 17 coauthors. 2000. Marine, estuarine, and diadromous fish stocks at risk of extinction in North America (exclusive of pacific salmonids). *Fisheries* 25(11)6-30.
- Myers, J. M., C. Busack, and D. Rawding. 2002. Identifying historical populations of Chinook and chum salmon and steelhead within the Lower Columbia River and Upper Willamette River Evolutionarily Significant Units. Lower Columbia River and Upper Willamette River Technical Recovery Document. 132 pp + appendices.

- Myers, J. M., C. Busack, D. Rawding, and A. Marshall. 2003. Historical population structure of Willamette and Lower Columbia River Basin Pacific Salmonids. Population Identification Subcommittee of the Willamette/Lower Columbia Technical Recovery Team. 183pp.
- Myers, J.M., R.G. Kope, et al. 1998. Status review of Chinook salmon from Washington, Idaho, Oregon, and California. NOAA Technical. Memorandum.
- Myers, K.W., and H.F. Horton. 1982. Temporal use of an Oregon estuary by hatchery and wild juvenile salmon, p. 377-392. In: V.S. Kennedy (ed.). Estuarine comparisons. Academic Press, Inc., New York.
- Myers, K.W.W. 1980. An investigation of the utilization of four study areas in Yaquina Bay, Oregon, by hatchery and wild juvenile salmonids. Master's thesis. Oregon State University, Corvallis. 233 p.
- Neal, V.T. 1972. Physical aspects of the Columbia River and its estuary. Pages 19-40 in A.T. Pruter and D.L. Alverson, eds. The Columbia River estuary and adjacent ocean waters. Bioenvironmental Studies. University of Washington Press, Seattle.
- Neave, F. 1961. Pacific salmon: ocean stocks and fishery developments. Proceedings of the Ninth Pacific Science Congress 1957(10):59-62.
- Neave, F. 1948. Fecundity and mortality of Pacific salmon. Proc. Trans. R. Soc. Can Ser. 3 42(5): 99-105.
- Neave, F. 1949. Game fish populations of the Cowichan River. Bulletin of the Fisheries Research Board of Canada 84: 1-32.
- Neave, F., and W.P. Wickett. 1953. Factors affecting the freshwater development of Pacific salmon in British Columbia. Proceedings of the Seventh Pacific Science Congress 1949(4): 548-556.
- Neave, F., T. Yonemori, and R. Bakkala. 1976. Distribution and origin of chum salmon in offshore waters of the North Pacific Ocean. Int. North Pac. Fish. Comm. Bull. 35:79 p.
- Nehlsen, W., J.E. Williams, and J.A. Lichatowich. 1991. Pacific salmon at the crossroads: Stocks at risk from California, Oregon, Idaho, and Washington. Fisheries 16(2):4-21.
- Nei, M. 1972. Genetic distance between populations. American Naturalist 106:283-292.
- Nei, M. 1978. Estimation of average heterozygosity and genetic distance from a small number of individuals. Genetics 89:583-590.
- Neilson, J.D., and C.E. Banford. 1983. Chinook salmon (*Oncorhynchus tshawytscha*) spawner characteristics in relation to redd physical features. Canadian Journal of Zoology 61:1524-1531.
- Neilson, J.D., G.H. Green, and D. Bottom. 1985. Estuarine growth of juvenile chinook salmon (*Oncorhynchus tshawytscha*) as inferred from otolith microstructure. Canadian Journal of Fisheries and Aquatic Sciences 42:899-908.
- Nelson, J.S., E.J. Crossman, H. Espinosa-Perez, L.T. Findley, C.R. Gilbert, R.N. Lea, J.D. Williams. 2003. The "Names of Fishes" list, including recommended changes in fish names: Chinook salmon for Chinook salmon, and Sander to replace Stizostedion for the sauger and walleye. Fisheries 28(7): 38-39.
- Nelson, K. and M. Soule. 1987. Genetical conservation of exploited fishes. Pages 345-368 in N. Ryman and F. Utter, eds. Population Genetics and Fishery Management. University of Washington Press, Seattle.
- Nelson, W.R. and J. Bodle. 1990. Ninety years of salmon culture at the Little White Salmon National Fish Hatchery. U.S. Fish and Wildlife Service Biological Report 90.

- Netboy, A. 1958. Salmon of the Pacific Northwest. Fish vs. Dams. Binforde and Mort, Portland, Oregon. 119 pp.
- Nickelson, T. E., M. F. Solazzi, and S.L. Johnson. 1986. Use of hatchery coho salmon (*Oncorhynchus kisutch*) psmolts to rebuild wild populations in Oregon coastal streams. Canadian Journal of Fisheries and Aquatic Sciences 43(12): 2443-2449.
- Nickelson, T.E. J.D. Rodgers, S.L. Johnson, and M.L. Solazzi. 1992. Seasonal changes in habitat use by juvenile coho salmon (*Oncorhynchus kisutch*) in Oregon coastal streams. Canadian Journal of Fisheries and Aquatic Sciences 49: 783-789.
- Nielsen, J.L. 1994. Molecular genetics and stock identification in Pacific salmon (*Oncorhynchus spp.*). Doctoral dissertation, University of California, Berkeley, 167 p.
- Nielsen, J.L. 1994. Invasive cohorts: impacts of hatchery-reared coho salmon on the trophic, developmental, and genetic ecology of wild stocks. Pages 361-385 in D. J. Stouder, K. L. Fresh, and R. J. Feller, editors. Theory and application in fish feeding ecology. University of South Carolina Press, Columbia.
- Nielson, J.L. 1992. Microhabitat-specific foraging behavior, diet and growth of juvenile coho salmon. Transactions of the American Fisheries Society 121:617-634.
- NMFS (National Marine Fisheries Service) Northwest Fisheries Science Center. 2000. Preliminary data on contaminant concentrations in salmonids and prey from the Columbia River estuary. EC Division. Interim Report, Sept. 2000.
- NMFS (National Marine Fisheries Service). 1995. Status Review of Coho Salmon from Washington, Oregon, and California. NMFS. Southwest Region. Protected Species Management Division, Long Beach, California.
- NMFS (National Marine Fisheries Service). 1995. Proposed recovery plan for Snake River salmon.
- NMFS (National Marine Fisheries Service). 1996. Factors for Decline, a supplement to the notice of determination for West Coast steelhead under the Endangered Species Act. National Marine Fisheries Service, Portland, Oregon.
- NMFS (National Marine Fisheries Service). 1996. Making Endangered Species Act Determinations of Effect for Individual or Grouped Actions at the Watershed Scale.
- NMFS (National Marine Fisheries Service). 1996. Making Endangered Species Act determinations of effect for individual or grouped actions at the watershed scale. Environmental and Technical Services Division, Habitat Conservation Branch.
- NMFS (National Marine Fisheries Service). 1997. Review of the Status of Chinook Salmon (*Oncorhynchus tshawytscha*) from Washington, Oregon, California and Idaho under the U.S. Endangered Species Act. Prepared by the West Coast Chinook Salmon Biological Review Team.
- NMFS (National Marine Fisheries Service). 1997. Status Review of Chum Salmon from Washington, Oregon, and California. National Marine Fisheries Service, Seattle, Washington.
- NMFS (National Marine Fisheries Service). 1999. Evaluation of the status of chinook and chum salmon and steelhead hatchery populations of ESUs identified in final listing determinations. Report of the Conservation Biology Division, Northwest Fisheries Science Center, Seattle, Washington.
- NMFS (National Marine Fisheries Service). 2000. White paper: passage of juvenile and adult salmonids past Columbia and Snake River dams. Northwest Fisheries Science Center, 2725 Montlake Blvd. E., Seattle, Washington, 98112-2097.

- NMFS. 2000. Predation on salmonids relative to the Federal Columbia River power system. White paper by NOAA Fisheries, Northwest Fisheries Science Center.
- NOAA Fisheries. 2003. Preliminary conclusions regarding the updated status of listed ESUs of West Coast salmon and steelhead. West Coast Salmon Biological Review Team, Co-manager Review Draft.
- Normandeau Associates Inc., J.R. Skalski, and Mid-Columbia Consulting Inc. 1995. Turbine passage survival of juvenile Chinook salmon (*Oncorhynchus tshawytscha*) at Lower Granite Dam, Snake River, Washington. Report to U.S. Army Corps of Engineers. Contract DACW68-95-C-0031, 78 p. (Available from U.S. Army Corps of Engineers, Walla Walla, WA 99362.)
- Normandeau Associates Inc., J.R. Skalski, and Mid-Columbia Consulting Inc. 1996. Draft report on potential effects of spillway flow deflectors on fish condition and survival at the Bonneville Dam, Columbia River. Report to the U.S. Army Corps of Engineers, Contract DACW57-95-C-0086, 51 p. plus App. (Available from U.S. Army Corps of Engineers, Portland, OR 97208.)
- Normandeau Associates Inc., J.R. Skalski, and Mid-Columbia Consulting Inc. 1999. Relative passage survival and injury mechanisms for Chinook salmon smolts within the turbine environment at McNary Dam, Columbia River. Draft Report to U.S. Army Corps of Engineers, Contract DACW68-96-D-003. (Available from U.S. Army Corps of Engineers, Walla Walla, WA 99362.)
- North, J.A., R.C. Beamesderfer, and T.A. Rien. 1993. Distribution and movements of white sturgeon in three lower Columbia River reservoirs. Northwest Science 67(2):105-111.
- Northcote, T.G. 1997. Why sea-run? An exploration into the migratory/residency spectrum of coastal cutthroat trout. Pages 20-26 in J.D. Hall, P.A. Bisson and R.E. Gresswell, eds. Sea-run cutthroat trout: biology, management, and future conservation. American Fisheries Society, Corvallis.
- Northcote, T.G., N.T. Johnston, and K. Tsumura. 1979. Feeding relationships and food web structure of lower Fraser River fishes. Westwater Research Center University British Columbia Technical Report 16. 73 pp.
- NRC (National Research Council). 1996. Upstream: Salmon and society in the Pacific Northwest. National Academy Press, Washington, D.C.
- Ocker, P.A., L.C. Stuehrenberg, M.L. Moser, A.L. Matter, J.J. Vella, B.P. Stanford. 2001. Monitoring Adult Pacific Lamprey (*Lampetra tridentate*) Migration Behavior in the Lower Columbia River using Radiotelemetry, 1998-1999. U.S. Army Corps of Engineers, Portland, Oregon.
- ODFW (Oregon Department of Fish and Wildlife). 1994. Effects of Lost Creek Dam on summer steelhead in the Rogue River. Phase II Completion Report. Rogue Basin Fisheries Evaluation Project. U.S. Army Corps of Engineers, Contract DACW57-77-C-0033, 235 pp. (Available from Oregon Department of Fish and Wildlife, P.O. Box 59, Portland, OR 97207.)
- ODFW (Oregon Department of Fish and Wildlife). 1994. Effects of Lost Creek Dam on summer steelhead in the Rogue River. Phase II completion report. Contract report submitted to U.S. Army Corps of Engineers, Portland, Oregon.
- ODFW (Oregon Department of Fish and Wildlife). 1996. K. Beiningen and T. Rien, eds. Effects of mitigative measures on productivity of white sturgeon populations in the Columbia River downstream from McNary Dam, and Determine status and habitat requirements of white sturgeon from the Columbia and Snake rivers upstream from McNary Dam. Annual Progress Report to the Bonneville Power Administration. Project 86-50, Portland, Oregon.

- ODFW and WDFW (Oregon Department of Fish and Wildlife and Washington Department of Fish and Wildlife). 1995. Status report: Columbia River fish runs and fisheries, 1938-94. Oregon Dep. Fish and Wildlife, P.O. Box 59, Portland, OR 97207, 291 pp.
- ODFW and WDFW (Oregon Department of Fish and Wildlife and Washington Department of Fish and Wildlife). 2001. Status report: Columbia River fish runs and fisheries, 1938-94. Oregon Dep. Fish and Wildlife, P.O. Box 59, Portland, OR 97207, 291 pp.
- ODFW and WDFW (Oregon Department of Fish and Wildlife and Washington Department of Fish and Wildlife). 2000. Status report – Columbia River fish runs and fisheries, 1938-1999. Oregon Department of Fish and Wildlife and Washington Department of Fish and Wildlife.
- OFC (Oregon Fish Commission). 1960. Results of a tagging program to enumerate the numbers and to determine the seasonal occurrence of anadromous fish in the Snake River and its tributaries. Report to U.S. Army Corps of Engineers, North Pacific Division, Portland, Oregon. Pages 20-22. in U.S. Army Corps of Engineers, Progress Report on the Fisheries Engineering Research Program, Portland, Oregon. (Available from U.S. Army Engineer District, Portland, OR.)
- Ogura, M. and S. Ito. 1994. Change in the known ocean distribution of Japanese chum salmon, *Oncorhynchus keta*, in relation to the progress of stock enhancement. Canadian Journal of Fisheries and Aquatic Sciences 51(3): 501-505.
- Okazaki, T. 1984. Genetic divergence and its zoogeographical implications in closely related species *Salmo gairdneri* and *Salmo mykiss*. Japanese Journal of Ichthyology 31:297-310.
- Olney, F.E. 1975. Life history and ecology of the northern pikeminnow *Ptychocheilus oregonensis* (Richardson) in Lake Washington. Master's thesis, University of Washington, Seattle.
- Olsen, E., P. Pierce, M. McLean, and K. Hatch. 1992. Stock Summary Reports for Columbia River Anadromous Salmonids Volume I: Oregon. U.S. Dep. Energy., Bonneville Power Administration. Project No. 88-108. (Available from BPA, Division of Fish and Wildlife, Public Information Officer, P.O. Box 3621, Portland, OR 97208).
- Oosterhout, G. R., and C. W. Huntington. 2003. A stochastic life-cycle model investigation of the potential benefits of a conservation hatchery program for supplementing Oregon Coast coho (*Oncorhynchus kisutch*). Report to Pacific Rivers Council.
- Otto, R. G. 1971. Effects of salinity on the survival and growth of pre-smolt coho salmon (*Oncorhynchus kisutch*). Journal of the Fisheries Research Board of Canada 28(3): 343-349.
- Otto, R.G. and J.E. McInerney. 1970. Development of salinity preference in pre-smolt coho salmon, *Oncorhynchus kisutch*. Journal of the Fisheries Research Board of Canada 27(4): 793-800.
- Pacific Fisherman. 1928. Record chum caught off Quadra. October 13.
- Paragamian, V. L., and G. Kruse. 2001. Kootenai River white sturgeon spawning migration behavior and a predictive model. North American Journal of Fisheries Management 21:10-21.
- Paragamian, V. L., G. Kruse, and V. Wakkinen. 1995. Kootenai River white sturgeon spawning and recruitment evaluation. Annual Progress Report to the Bonneville Power Administration, Portland, Oregon. 68 pp.
- Paragamian, V. L., G. Kruse, and V. Wakkinen. 1997. Kootenai River white sturgeon spawning and recruitment evaluation. Annual Progress Report to the Bonneville Power Administration, Portland, Oregon. 66 pp.

- Paragamian, V.L., R.C.P. Beamesderfer, and S.C. Ireland. In Review. Status, Population Dynamics, and Future Prospects of an Endangered Kootenai River Sturgeon Population with and without Hatchery Intervention.
- Paragamian, V.L., G. Kruse, and V. Wakkinen. 2001 Spawning habitat of Kootenai River white sturgeon, post-Libby Dam. *North American Journal of Fisheries Management* 21:22-33.
- Parker, R.M., M.P. Zimmerman, and D.L. Ward. 1995. Variability in biological characteristics of northern pikeminnow in the Lower Columbia and Snake rivers. *Transactions of the American Fisheries Society* 124:335-346.
- Parker, R.R. 1962. A concept of the dynamics of pink salmon populations, Pages 203-211 in N.J. Wilimovsky, ed. *Symposium on Pink Salmon*. H.R. MacMillan Lectures in Fisheries. Institute of Fisheries, University of British Columbia, Vancouver.
- Parker, R.R. 1965. Estimation of sea mortality rates for the 1961 brood-year pink salmon of the Bella Coola area, British Columbia. *Journal of the Fisheries Research Board of Canada* 22:1523-1554.
- Parkinson, E.A. 1984. Genetic variation in populations of steelhead (*Salmo gairdneri*) in British Columbia. *Canadian Journal of Fisheries and Aquatic Sciences* 41:1412-1420.
- Parsley, M.J., and K.M. Kappenman. 2000. White sturgeon spawning areas in the lower Snake River. *Northwest Science* 74:192-201.
- Parsley, M.J., and L.G. Beckman. 1994. White sturgeon spawning and rearing habitat in the lower Columbia River. *North American Journal of Fisheries Management* 14:812-827.
- Parsley, M.J., D. Gallion, and M.B. Sheer. 2001. Effect of proposed reservoir drawdowns on productivity of white sturgeon *Acipenser transmontanus* populations. in *Proceedings of the Fourth International Symposium on Sturgeons*, Oshkosh, Wisconsin, USA. 8-13 July, 2001.
- Parsley, M.J., L.G. Beckman, and G.T. McCabe, Jr. 1993. Spawning and rearing habitat use by white sturgeons in the Columbia River downstream from McNary Dam. *Transactions of the American Fisheries Society* 122(2):217-227.
- Parsley, M.J., P.J. Anders, A.I. Miller, L.G. Beckman, and G.T. McCabe, Jr. 2002. Recovery of white sturgeon populations through natural production: Understanding the influence of abiotic and biotic factors on spawning and subsequent recruitment. Pages 55-66 in W. VanWinkle, P. Anders, D. Dixon, and D. Secor, eds. *Biology, Management and Protection of North American Sturgeons*. American Fisheries Society Symposium 28.
- Patten, B.G. 1971. Predation by sculpins on fall Chinook salmon, *Oncorhynchus tshawytscha*, fry of hatchery origin. *U.S. Nat. Mar. Fish. Serv. Spec. Sci. Rep. Fish.* 621. 14 pp.
- Paul, A.J. 2000. Recruitment dynamics in bull trout (*Salvelinus confluentus*): linking theory and data to specie management. Doctoral dissertation, University of Calgary, Department of Biological Science, Calgary, Alberta.
- Pearcy, W.G. 1997. The sea-run and the sea. Pages 29-36 in J.D. Hall, P.A. Bisson and R.E. Gresswell, eds. *Sea-run cutthroat trout: biology, management, and future conservation*, p. 29-36. American Fisheries Society, Corvallis, Oregon.
- Pearsons, T.N., H.W. Li, and G.A. Lamberti. 1992. Influence of habitat complexity on resistance to flooding and resilience of stream fish assemblages. *Transactions of the American Fisheries Society* 121: 427-436.
- Percy, W.G. 1992. What have we learned in the last decade? What are research priorities? College of Oceanic and Atmospheric Sciences, Oregon State University, Corvallis.

- Perkins, S.J. 1989. Interactions of landslide-supplied sediment with channel morphology in forested watersheds. Unpublished Master's thesis. University of Washington, Seattle.
- Perrin, C.J., A. Heaton, and M.A. Laynes. 1999. White sturgeon (*Acipenser transmontanus*) spawning habitat in the Lower Fraser River, 1988. Limnotek Research and Development Inc., Vancouver, British Columbia. Report to British Columbia Ministry of Fisheries, Victoria.
- Peterman, R.M. 1978. Testing for density dependent marine survival in Pacific salmonids. *Journal of the Fisheries Research Board of Canada* 35:1434-1450.
- Petersen, J.H. 2001. Density, aggregation, and body size of northern pikeminnow preying on juvenile salmonids in a large river. *Journal of Fish Biology* 58:1137-1148.
- Petersen, J.H., and D. L. DeAngelis. 1992. Functional response and capture timing in an individual-based model: predation by northern pikeminnow (*Ptychocheilus oregonensis*) on juvenile salmonids in the Columbia River. *Canadian Journal of Fisheries and Aquatic Sciences* 49:2551-2565.
- Peterson, N.P. 1980. The role of spring ponds in the winter ecology and natural production of coho salmon (*Oncorhynchus kisutch*) on the Olympic Peninsula, Washington. Master's thesis. University of Washington, Seattle. 96 pp.
- Phelps, S., J. Uehara, D. Hendrick, J. Hymer, A. Blakley, and R. Brix. 1995. Genetic diversity units and major ancestral lineages for chum salmon in Washington. in Busack, C., and J.B. Shaklee, eds. Genetic diversity units and major ancestral lineages of salmonid fishes in Washington, p. C1-C55. Tech. Rep. RAD 95-02, Washington Department of Fish and Wildlife, 600 Capitol Street N., Olympia, WA 98501-1091.
- Phelps, S.R., B.M. Baker, P.L. Hulett, and S.A. Leider. 1994. Genetic analysis of Washington steelhead: Initial electrophoretic analysis of wild and hatchery steelhead and rainbow trout. Washington Department of Fish and Wildlife, Fisheries Management Program Report 94-9.
- Phillips, R.B., and K.A. Pleyte. 1991. Nuclear DNA and salmonid phylogenetics. *Journal of Fish Biology* 39 (Suppl A);259-275.
- Phillips, R.B., K.A. Pleyte, and P.E. Ihssen. 1989. Patterns of chromosomal nuclear organize region (NOR) variation in fishes of the genus *Salvelinus*. *Copeia*: 47-53.
- Phillips, R.W., R.L. Lantz, E.W. Claire, and J.R. Moring. 1975. Some effects of gravel mixtures on emergence of coho salmon and steelhead trout fry. *Transactions of the American Fisheries Society* 104:461-466.
- Pickett, P.J. and R. Harding. 2002. Total maximum daily load for Lower Columbia River total dissolved gas. Washington State Department of Ecology, No. 02-03-004.
- Pike, G.C. 1950. Stomach contents of whales caught off the coast of British Columbia. *Fish Res. Board Can. Prog. Rep. Pac. Coast Stations* 83: 27-28.
- Pike, G.C. 1951. Lamprey marks on whales. *Journal of the Fisheries Research Board of Canada* 8(4): 275-280.
- Pitcher, T.J. 1986. Functions of shoaling in teleosts. Pages 294-337 in Fisher, T.J., ed. *The behavior of teleost fishes*, Johns Hopkins University Press, Baltimore, Maryland.
- Platts, W.S. 1991. Livestock Grazing. Chapter 11 in *Influences of Forest and Rangeland Management on Salmonid Fishes and Their Habitats*, W.R. Meehan ed. American Fisheries Society Special Publication 19:389-423.

- Pletcher, F.T. 1963. The life history and distribution of lampreys in the Salmon and certain other rivers in British Columbia, Canada. Master's thesis, University of British Columbia, Vancouver. 195pp.
- Poe, T.P., H.C. Hansel, S.Vigg, D.E. Palmer, and L.A. Prendergast. 1991. Feeding of predaceous fishes on out-migrating juvenile salmonids in John Day Reservoir, Columbia River. Transactions of the American Fisheries Society 120:405-420.
- Poe, T.P., H.C. Hansel, S. Vigg, D.E. Palmer, and L.A. Prendergast. 1991. Feeding of predaceous fishes on out-migrating juvenile salmonids in the John Day Reservoir, Columbia River. Transactions of the American Fisheries Society 120:405-420.
- Poe, T.P., R.S. Shively, and R.A. Tabor. 1994. Ecological consequences of introduced piscivorous fishes in the lower Columbia and Snake Rivers, Pages 347-360 in D.J. Strouder, K.L. Fresh, and R.J. Feller ed. Theory and application in fish feeding ecology. University of South Carolina Press, Columbia.
- Poff, N.L., J.D. Allan, M.B. Bain and others. 1997. The Natural Flow Regime: A paradigm for river conservation and restoration: BioScience 47(11): 769-784.
- Post, J.R., E.A. Parkinson and N.T. Johnston. 1999. Density-dependent processes in structured fish populations: interaction strengths in whole lake experiments. Ecological Monographs 69:155-175.
- Potter, I.C. 1970. The life cycles and ecology of Australian lampreys of the genus *Mordacia*. Journal of Zoology (London) 161: 487-511.
- Potter, I.C., and F.W.H. Beamish. 1977. The freshwater biology of adult and anadromous sea lampreys *Petromyzon marinus*. Journal of Zoology (London):181:113-130.
- Pratt, K.L. 1984. Habitat selection and species interactions of juvenile westslope cutthroat trout (*Salmo clarki lewisi*) and bull trout (*Salvelinus confluentus*) in the upper Flathead River basin. Masters thesis, University of Idaho, Moscow.
- Pratt, K.L. 1992. A review of bull trout life history. Pages 5-9 in P.J. Howell, and D.V. Buchanan, eds. Proceedings of the Gearhart Mountain bull trout workshop. Oregon Chapter of the American Fisheries Society, Corvallis.
- Pravdin, I.F. 1940. A review of investigations on the far-eastern salmon. Izv. Tikhookean. Nauchno-Issled. Inst. Rybn. Khoz. Okeangr. 18:5-105. (Transl. from Russian; Fish. Res. Board Can. Transl. Ser. 371).
- PSMFC (Pacific States Marine Fisheries Commission) 1992. White sturgeon management framework plan. PSMFC, Portland, Oregon.
- PSMFC (Pacific States Marine Fisheries Commission). 1994. Regional Mark Information System (RMIS) Coded-wire tag on-line database. (Available from Pacific States Marine Fisheries Commission, 45 SE 82nd Dr., Suite 100, Gladstone, OR 97027).
- Ptolemy, R.A. 1979. Production and carrying capacity relative to the design of a rearing channel for rainbow trout on Hill creek near Galena Bay, West Kootenay. British Columbia Ministry of Environment, Fish and Wildlife Branch, Victoria, British Columbia.
- Ptolemy, R.A., J.C. Wightman and C.D. Tredger. 1977. A fisheries reconnaissance assessment of the Salmon River drainage, Vancouver, B.C. Island relative to enhancement opportunities. Memo. report. Fish and Wildlife Branch, Ministry of Recreation and Conservation. Victoria, British Columbia. 58 pp.
- Pycha, R. L. 1956. Progress report on white sturgeon studies. California Fish and Game 42:23-35.

- Pyper, B.J., F.J. Mueter, R.M. Peterman, D.J. Blackbourn, and C.C. Wood. 2001. spatial convariation in survival rates of Northeast Pacific pink salmon (*Oncorhynchus gorbuscha*). Canadian Journal of Fisheries and Aquatic Sciences 58: 1501-1515.
- Quinn, T.P. 1984. Homing and straying in Pacific Salmon. Mechanisms of Migration in Fishes. Pages 357-362 in J.D. McCleave, J.P. Arnold, J.J. Dodson and W.H. Neills. Plenum Press, New York.
- Quinn, T.P. 1993. A review of homing and straying of wild and hatchery-produced salmon. Fisheries Research (Amst.) 18:29-44.
- Quinn, T.P., S. Hodgson, and C. Peven. 1997. Temperature, flow and migration of adult sockeye salmon (*Oncorhynchus nerka*) in the Columbia River. Canadian Journal of Fisheries and Aquatic Sciences 54:1349-1360.
- Quinn, T.P., S. Hodgson, et al. 1997. Temperature, flow, and the migration of adult sockeye salmon (*Oncorhynchus nerka*) in the Columbia River. Canadian Journal of Fisheries and Aquatic Sciences 54(6): 1349-1360.
- Radtke, H.D., and S.W. Davis. 2000. Economic feasibility of salmon enhancement propagation programs. Pages 381-392 in E.E. Knudsen, C. S. Steward, D. D. MacDonald, J. E. Williams, and D. W. Reiser, editors. Sustainable fisheries management: Pacific salmon, Lewis Publishers, Boca Raton, Florida.
- Randall, R.G., M.C. Healey, and J.B. Dempson. 1987. Variability in length of freshwater residence of salmon, trout, and char. American Fisheries Society Symposium 1:27-41.
- RASP (Regional Assessment of Supplementation Project). 1992. Supplementation in the Columbia Basin: Summary report series. Final Report DOE/BP-01830-14, Bonneville Power Administration, Portland, Oregon.
- Ratliff, D., S. Thiesfield, W. Weber, A. Stuart, M. Riehle, D. Buchanan. 1996. Distribution, life history, abundance, harvest, habitat, and limiting factors of bull trout in the Metolius River and Lake Billy Chinook, Oregon, 1983-1994. Oregon Department of Fish and Wildlife, Information Report 96-7.
- Ratliff, D.E., and P.J. Howell. 1992. The status of bull trout populations in Oregon. Pages 10-17 in P.J. Howell. and D.V. Buchanan, eds. Proceedings of the Gearhart Mountain bull trout workshop. Oregon Chapter of the American Fisheries Society, Corvallis.
- Raymond, H.L. 1979. Effects of dams and impoundments on migrations of juvenile Chinook salmon and steelhead from the Snake River, 1966 to 1975. Transactions of the American Fisheries Society 108:505-529.
- Reeves, G.H., J.D. Hall, and S.V. Gregory. 1997. The impact of land-management activities on coastal cutthroat trout and their freshwater habitats. Pages 138-144 in J.D. Hall, P.A. Bisson and R.E. Gresswell, eds. Sea-run cutthroat trout: biology, management, and future conservation, American Fisheries Society, Corvallis, Oregon.
- Refalt, W. 1985 Wetland in extremis: A nationwide survey. Wilderness Winter. 1985:28-41.
- Reimers, P. E. 1973. The length of residence of juvenile fall Chinook salmon in Sixes River, Oregon, Oregon Fish Commission: 1-43.
- Reimers, P.E. 1971. The length of residence of juvenile fall chinook in the Sixes River, Oregon. Doctoral dissertation. Oregon State University, Corvallis. 99 pp.

- Reischel, T.S., and T.C. Bjornn. 2003. Influence of fishway placement on fallback of adult salmon at the Bonneville Dam on the Columbia River. *North American Journal of Fisheries Management* 23:1215-1224.
- Reisenbichler, R.R. and S.R. Phelps. 1989. Genetic variation in steelhead (*Salmo gairdneri*) from the north coast of Washington. *Canadian Journal of Fisheries and Aquatic Sciences* 46:66-73.
- Reisenbichler, R.R., J.D. McIntyre, M.F. Solazzi, and S.W. Landino. 1992. Genetic variation in steelhead of Oregon and Northern California. *Transactions of the American Fisheries Society* 121:158-169.
- Reisenbichler, R.R., and J.D. McIntyre. 1977. Genetic differences in growth and survival of juvenile hatchery and wild steelhead trout, *Salmo gairdneri*. *Journal of the Fisheries Research Board of Canada* 34: 123-128.
- Reiser, D. W., and T. C. Bjornn. 1979. Habitat requirements of anadromous salmonids. Page 54 in W.R. Mehan, ed. *Influence of forest and rangeland management on anadromous fish habitat in the western United States and Canada 1*. U.S. Forest Service General Technical Report.
- Resienbichler, R.R. 1997. Genetic factors contributing to the declines of anadromous salmonids in the Pacific Northwest. Pages 223-244 in D. J. Stouder, P. A. Bisson, and R. J. Naiman, editors. *Pacific salmon and their ecosystems: status and future options*. Chapman and Hall, New York.
- Rice, J. A., L. B. Crowder, and F. P. Binkowski. 1987. Evaluating potential sources of mortality for larval bloater: starvation and vulnerability to predation. *Canadian Journal of Fisheries and Aquatic Sciences* 44:467-472.
- Rich, W.H. 1920. Early history and seaward migration of Chinook salmon in the Columbia and Sacramento Rivers. *Bull. Bur. Fish. (U.S.)* 37. 74 pp.
- Rich, W.H. 1942. The salmon runs of the Columbia River in 1938. *Fisheries Bull., U.S.* 50(37):103-147.
- Richards, J.E. 1980. Freshwater biology of the anadromous Pacific lamprey *Lampetra tridentate*. Master's thesis, University of Guelph, Ontario, Canada. 99pp.
- Ricker, W. E. 1975. Computation and interpretation of biological statistics of fish populations. *Bulletin of the Fisheries Research Board of Canada* 191.
- Ricker, W.E. 1941. The consumption of young sockeye salmon by predaceous fish. *Journal of the Fisheries Research Board of Canada* 5:293-313.
- Ricker, W.E. 1981. Changes in the average size and average age of Pacific salmon. *Canadian Journal of Fisheries and Aquatic Sciences* 38(12): 1636-1656.
- Riddell, B.E. 1993. Spatial organization of Pacific salmon: What to conserve? Pages 23-41 In J.G. Cloud, and G.H. Thorgaard, editors. *Genetic conservation of salmonid fishes*. Plenum Press, New York.
- Rieman, B.E. and J.D. McIntyre. 1996. Spatial and temporal variability in bull trout redd counts. *North American Journal of Fisheries Management* 16: 132-146.
- Rieman, B.E., and J.D. McIntyre. 1993. Demographic and habitat requirements for conservation of bull trout. U.S. Forest Service, Intermountain Research Station General Technical Report INT-302.
- Rieman, B.E., and R. C. Beamesderfer. 1990. White sturgeon in the Lower Columbia River: Is the stock overexploited? *North American Journal of Fisheries Management* 10:388-396.
- Rieman, B.E., and R.C. Beamesderfer. 1990. Dynamics of a northern pikeminnow population and the potential to reduce predation on juvenile salmonids in a Columbia River reservoir. *North American Journal of Fisheries Management* 10:228-241.

- Rieman, B.E., R.C. Beamesderfer, S. Vigg, and T.P. Poe. 1991. Estimated loss of juvenile salmonids to predation by northern pikeminnow, walleyes, and smallmouth bass in John Day Reservoir, Columbia River. *Transactions of the American Fisheries Society* 120:448-458.
- RL&L Environmental Services Ltd. 1994. Status of white sturgeon in the Columbia River, British Columbia. Report prepared for B.C. Hydro, Environmental Affairs, Vancouver, British Columbia by RL&L Environmental Services Ltd, Vancouver. Report No. 377F: 101 pp.
- RL&L Environmental Services Ltd. 1996. Columbia River white sturgeon investigations. 1995 study results. Report prepared for B.C. Hydro, Kootenay Generation, Vancouver, British Columbia, and British Columbia Ministry of Environment, Lands, and Parks, Nelson Region. RL&L Report No. 96-377F. 94pp.
- Roby, D.D., D.P. Craig, K. Collis, and S.L. Adamany. 1998. Avian Predation on Juvenile Salmonids in the Lower Columbia River 1997 Annual Report. Bonneville Power Administration Contract 97BI33475 and U.S. Army Corps of Engineers Contract E96970049. 70 pp.
- Rochard, E. G., G. Castelnaud, and M. Lepage. 1990. Sturgeons (Pisces: Acipenseridae): threats and prospects. *Journal of Fish Biology* 37 (Supplement A): 123-132.
- Rodgers, J.D., S.L. Johnson, T.E. Nickelson, and M.F. Solazzi. 1993. The seasonal use of natural and constructed habitat by juvenile coho salmon (*Oncorhynchus kisutch*) and preliminary results from two habitat improvement projects on smolt production in Oregon coastal streams. Pages 344-351 in L. Berg and P.W. Delaney, eds. *Proceedings of the Coho Workshop*, Nanaimo, British Columbia, May 26-28, 1992.
- Roelofs, T. D. 1983. Current status of California summer steelhead (*Salmo gairdneri*) stocks and habitat, and recommendations for their management. Submitted to USDA, National Marine Fisheries Service, Portland, Oregon.
- Roffe, T.J., and B.R. Mate. 1984. Abundances and feeding habits of pinnipeds in the Rogue River, Oregon. *Journal of Wildlife Management* 48(4): 1262-1274.
- Roper, B.B. 1995. Ecology of anadromous salmonids within the upper south Umpqua River basin, Oregon. Dissertation, University of Idaho, Moscow.
- Rucklelshaus, M., K. Currens, R. Fuerstenberg, W. Graeber, K. Rawson, N. Sands, and J. Scott. 2002. Planning ranges and preliminary guidelines for delisting and recovery of the Puget Sound chinook salmon evolutionarily significant unit. Puget Sound Technical Recovery Team.
- Ruggerone, G. T. 1986. Consumption of Migrating Juvenile Salmonids by Gulls Foraging below a Columbia River Dam. *Transactions of the American Fisheries Society* 115: 736-742.
- Ruggles, C.P. 1966. Depth and velocity as a factor in a stream rearing and production of juvenile coho salmon. *Canadian Fish. Cult.* 38:37-53.
- Russell, J.E., F.W.H. Beamish, and R.J. Beamish. 1987. Lentic spawning by the Pacific lamprey, *Lampetra tridentata* *Canadian Journal of Fisheries and Aquatic Sciences* 44: 476-478.
- Rutherford, J.C., S. Blackett, C. Blackett, L. Saito, and R.J. Davies-Colley. 1997. Predicting the effects of shade on water temperature in small streams. *New Zealand Journal of Marine and Freshwater Research* 31:707-721.
- Rutter, C. 1904. Natural history of the Quinnat salmon. *Investigations on the Sacramento River, 1896-1901.* *Bull. U.S. Fish Comm.* 22:65-141

- Sakuramoto, K., and S. Yamada. 1980. A study on the planting effect of salmon. 1. A mathematical model for the derivation of their rate of return and its applications. Bull. Japanese Soc. Sci. Fish. 46(6):653-661.
- Salo, E. O. 1991. Life history of chum salmon. Pages 231-309 in C. Groot and L. Margolis, eds. Pacific salmon life histories, University of British Columbia Press, Vancouver.
- Salo, E. O. and W. H. Bayliff 1958. Artificial and natural production of silver salmon (*Oncorhynchus kisutch*) at Minter Creek, Washington. Research Bulletins Washington Department, Dep. Fish 4, 76.
- Salo, E.O., and R.E. Noble. 1953. Chum salmon upstream migration, Pages 1-9. In: Minter Creek Biological Station progress report, September through October 1953. Washington Department of Fisheries, Olympia. 14 p.
- Salo, E.O., N.J. Bax, T.E. Prinslow, C.J. Whitmus, B.P. Snyder, and C.A. Simenstad. 1980. The effects of construction of naval facilities on the outmigration of juvenile salmonids from Hood Canal, Washington. Final Report, FRI-UW-8006, 159 p. Fish. Res. Inst., University of Washington, Seattle.
- Sandercock, F. K. 1991. Life history of coho salmon (*Oncorhynchus kisutch*). Pacific salmon life histories. C. Groot and L. Margolis. University of British Columbia Press, Vancouver. 395-446.
- Sano, S. 1966. Chum salmon in the East, pages 4-58. In: Salmon of the North Pacific Ocean. Part III. A review of the life history of North Pacific salmon. Int. North Pac. Fish. Comm. Bull. 18.
- Sano, S., and A. Nagasawa. 1958. Natural propagation of chum salmon, *Oncorhynchus keta*, in Memu River, Tokachi. Sci. Rep. Hokkaido Salmon Hatchery 12:1-19. (Partial transl. from Japanese; Fish. Res. Board Can. Transl. Ser. 198.)
- Scarlett, W.J., and C.J. Cederholm. 1984. Juvenile coho salmon fall-winter utilization of two small tributaries of the Clearwater River, Jefferson County, Washington, Pages 227-242 in J.M. Walton and D.B. Houston, eds. Proceedings of the Olympic Wild Fish Conference, March 23-25, 1983. Fisheries Technology Program, Peninsula College, Port Angeles, Washington.
- Schluchter, M.D., and J.A. Lichatowich. 1977. Juvenile life histories of Rogue River spring Chinook salmon, *Oncorhynchus tshawytscha* (Walbaum), as determined by scale analysis. Oregon Department Fish Wildlife. Reports. Ser. Fish. 77-5. 24 pp.
- Schoonmaker, P.K., T. Gresh, J. Lichatowich, and H.D. Radtke. 2003. Past and present Pacific salmon abundance: bioregional estimates for key life history stages. Pages 33- 40 In J.D. Stockner, editor. Nutrients in the freshwater salmonid ecosystem: Sustaining production and biodiversity. American Fisheries Society Special Publication 34. Bethesda, Maryland.
- Schreck, C.B., H.W. Li, R.C. Hjort, and C.S. Sharpe. 1986. Stock identification of Columbia River chinook salmon and steelhead trout. Final Report 1986 to the Department of Energy, Bonneville Power Administration, Division of Fish and Wildlife, Portland, Oregon.
- Schreiner, J.V. 1977. Salmonid outmigration studies in Hood Canal, Washington. Master's thesis. University of Washington, Seattle. 91 pp.
- Schroder, S.L., K.V. Koski, B.P. Snyder, K.J. Bruya, G.W. George, and E.O. Salo. 1974. Big Beef Creek studies, Pages 26-27 in Research in fisheries 1973. University of Washington, College of Fisheries Contribution 390.
- Schuck, M. and H. Kruse. 1982. South Fork Toutle River fish trap operation and salmonid investigations, 1981-82. Washington Department of Game (WDG) 82-11.

- Scott, W.B. and E.J. Crossman. 1973. Freshwater Fishes of Canada. Fisheries Research Board of Canada, Ottawa.
- Scott, W.B. and E.J. Crossman. 1973. Freshwater Fishes of Canada. Bulletin 184. Fisheries Research Board of Canada.
- Scott, W.B., and E.J. Crossman. 1973. Freshwater fishes of Canada. Bulletin of the Fisheries Research Board of Canada. 966 p.
- Scrivener, J.C., and B.C. Andersen. 1982. Logging impacts and some mechanisms which determine the size of spring and summer populations of coho salmon fry in Carnation Creek, Pages 257-272 in G.F. Hartman, ed. Proceedings of the Carnation Creek Workshop: a ten year review. Pacific Biological Station, Nanaimo, BC.
- Scuett-Hames, D, A.E. Pleus, and D. Smith. 1999. Method Manual for the Salmonid Spawning Habitat Availability Survey. TFW Monitoring Program. Washington Department of Natural Resources. Olympia.
- Sea Resources. 2001. Sea Resources website www.searesources.org
- Secor, D., H. and Gunderson. 1988. Effects of hypoxia and temperature on survival, growth, and respiration of juvenile Atlantic sturgeon *Acipenser oxyrinchus*. Fisheries Bulletin 96:603-613.
- Secor, D.H., and E.J. Niklitschek. 2001. Hypoxia and sturgeons: Report to the Chesapeake Bay Program dissolved oxygen criteria team. Technical Report Series No. TS-314-01-CBL; Chesapeake Biological Laboratory, Solomons, Maryland.
- Secor, D.H., P.J. Anders, W. Van Winkle, and D.A. Dixon. 2002. Can We Study Sturgeons to Extinction? What We Do and Don't Know about the Conservation of North American Sturgeons. Pages 3-12 In: W. VanWinkle, P. Anders, D. Dixon, and D. Secor, eds. Biology, Management and Protection of North American Sturgeons. American Fisheries Society Symposium 28.
- Secor, D.H., P.J. Anders, W. Van Winkle, and D.A. Dixon. 2002. Can We Study Sturgeons to Extinction? What We Do and Don't Know about the Conservation of North American Sturgeons. Pages 3-12 In: W. VanWinkle, P. Anders, D. Dixon, and D. Secor, eds. Biology, Management and Protection of North American Sturgeons. American Fisheries Society Symposium 28.
- Seiler, D., S. Neuhauser, and M. Ackley. 1981. Upstream/downstream salmonid trapping project, 1977-1980. Progress Report 144, Washington Department of Fisheries, Olympia, 197 pp. (Available from Washington Department Fish and Wildlife, 600 Capitol Way N., Olympia, WA 98504-1091.)
- Semakula, S. N. and P. A. Larkin. 1968. Age, growth, food, and yield of the white sturgeon (*Acipenser transmontanus*) of the Fraser River, British Columbia. Journal of the Fisheries Research Board of Canada 184.
- Semko, R.S. 1954. The stocks of West Kamchatka salmon and their commercial utilization . Izv. Tikhookean. Nauchno-Issled. Inst. Rybn. Khoz. Okeanogr. 41:3-109. (transl. from Russian; Fish. Res. Board. Can. Transl. Ser. 288).
- Serns, S. L. 1982. Influence of various factors on density and growth of age-0 walleye in Escanaba Lake, Wisconsin, 1958-1980. Transactions of the American Fisheries Society 111:299-306.
- Setter, A. and E. Brannon. 1992. A summary of stock identification research on white sturgeon of the Columbia River (1985-1990). Project No. 89-44. Final Report to the Bonneville Power Administration, Portland, Oregon.
- Sexauer, H.M. and P.W. James. 1997. Microhabitat use by juvenile trout in four streams located in the eastern Cascades, Washington. Pages 361-370 in W.C. Mackay, M.K. Brewin and M. Monita,

- eds. Friends of the Bull Trout Conference Proceedings. Bull Trout Task Force (Alberta), Trout Unlimited Calgary, Alberta.
- Sexaur, H.S. 1994. Life history aspects of bull trout, *Salvelinus confluentus*, in the eastern Cascades, Washington. Master's thesis, Central Washington University, Ellensburg.
- Shapovalov, L., and A. C. Taft. 1954 The life histories of the steelhead rainbow trout (*Salmo gairdneri gairdneri*) and silver salmon (*Oncorhynchus kisutch*) with special reference to Waddell Creek, California, and recommendations regarding their management. California Department of Fish and Game Fish Bulletin No. 98, Sacramento. 375pp.
- Shaw, P.A., and J.A. Maga. 1943 The effect of mining silt on yield of fry from salmon spawning beds. California Fish and Game 29:29-41.
- Shelton, J.M. 1955. The hatching of Chinook salmon eggs under simulated stream conditions. Prog. Fish-Cult. 17:20-35.
- Shelton, J.M., and R.D. Pollock. 1966. Siltation and egg survival in incubation channels. Transactions of the American Fisheries Society 95:183-187.
- Shepard, B., K. Pratt, and J. Graham. 1984. Life histories of westslope cutthroat and bull trout in the upper Flathead River Basin, Montana. Montana Department of Fish, Wildlife and Parks, Kalispell.
- Sheppard, D. 1972. The presents status of the steelhead trout stocks along the Pacific coast. Pages 519-556 in D. H. Rosenberd, ed. A review of the oceanography and renewable resources of the northern Gulf of Alaska. Institute of Marine Science, University of Alaska, Fairbanks.
- Shields, F. D., Jr. and Nunnally, N. R. 1984. Environmental Aspects of Clearing and Snagging Projects, Journal of Environmental Engineering, ASCE,110 (1):152-165.
- Shirvell, C.S. 1990. role of instream rootwads as juvenile coho salmon (*Oncorhynchus kisutch*) and steelhead trout (*O. mykiss*) cover habitat under varying stream flows. Canadian Journal of Fisheries and Aquatic Science 47:852-861.
- Sibert, J., and B. Kask. 1978. Do fish have diets? P. 48-57 in B.G. Shepherd and R.M. Ginetz (rapps.). Proceedings of the 1977 Northeast Pacific Chinook and Coho Salmon Workshop. Fish. Mar. Serv. (Can.) Tech. Rep. 759.
- Simenstad, C. A., K. L. Fresh, et al. 1982. The role of Puget Sound and Washington Coastal estuaries in the life history of Pacific salmon: An unappreciated function. Estuarine comparison. Pages 343-364 in V. Kennedy, ed. Academic Press, Inc. New York
- Simenstad, C.A., and E.O. Salo 1982. Foraging success as a determinant of estuarine and nearshore carrying capacity of juvenile chum salmon (*Oncorhynchus keta*) in Hood Canal, Washington, p. 21-37. In: B.R. Melteff and R.A. Neve (eds.). Proceedings of the North Pacific Aquaculture Symposium. Alaska Sea Grant Rep. 82-2.
- Simpson, J., and R. Wallace. 1982. Fishes of Idaho. University of Idaho Press, Moscow.
- Sinclair, M. 1988. Marine Populations: an essay on population regulation and speciation. University of Washington Press, Seattle and London. 252 pp.
- Sinokrot, B.A. and H.G. Stefan. 1993. Stream temperature dynamics: measurements and modeling. Water Resources Research 29(7):2299-2312.
- Sleeper, J. D. 1994. Seasonal changes in distribution and abundance of salmonids and habitat availability in a coastal Oregon Basin. Oregon State University, Corvallis.

- Smirnov, A.I. 1975. The biology, reproduction, and development of the Pacific salmon. Izdatel'stvo Moskovogo Universiteta, Moscow, USSR. 335 p. (Transl. from Russian; Fish. Mar. Serv. (Can.) Transl. Ser. 3861)
- Smith, C., J. Nelson, S. Pollard, S. McKay, B. May, J. Rodzen, and B. Koop. 2001. Population genetic analysis of white sturgeon (*Acipenser transmontanus*) in the Fraser River. In Proceedings of the Fourth International Symposium on Sturgeons, Oshkosh, Wisconsin, USA. 8-13 July, 2001.
- Smith, H.A. 1974. Spillway redesign abates gas supersaturation in Columbia River. Civil Engineering-ASCE, Sept., 4 p.
- Smith, R.D., R.C. Sidle, PE. Porter, and J.R. Noel. 1993. Effects of experimental removal of large woody debris on the channel morphology of a forest, gravel-bed stream. Journal of Hydrology (Amsterdam) 152:153-178.
- Smith, S.B. 1969. Reproductive isolation in summer and winter races of steelhead trout. in T.G. Northcote, ed. Symposium on Salmon and Trout in Streams. MacMillan Lectures in Fisheries, University of British Columbia, 388 p.
- Smoker, W.A. 1953. Stream flow and silver salmon production in Western Washington. Washington Department of Fisheries Research Paper 1:5-12.
- Smoker, W.W., A.J. Gharrett, and M.S. Stekoll. 1998. Genetic variation of return date in a population of pink salmon: A consequence of fluctuating environment and dispersive selection? Alaska Fishery Research Bulletin 5(1):46-54.
- Soin, S.G. 1954. Pattern of development of summer chum, masu, and pink salmon. Tr. Soveshch. Ikhyiol. Kom. Akad. Nauk SSSR 4:144-155. (Transl. from Russian; in: Pacific salmon: selected articles from Soviet periodicals, p. 42-54. Israel Program for Scientific Translations, Jerusalem, 1961)
- Solazzi, M.F., J.D. Rodgers, and S.L. Johnson. 1992. Development and evaluation of techniques to rehabilitate Oregon's wild salmonids. Oregon Department of Fish and Wildlife, Fish Research Project F-125-R, Annual Progress Report, Portland, Oregon.
- Solazzi, M.F., T.E. Nickelson, S.L. Johnson, and J.D. Rodgers. 2000. Effects of increasing winter rearing habitat on abundance of salmonids in two coastal Oregon streams. Canadian Journal of Fisheries and Aquatic Sciences 57: 906-914.
- Solazzi, M.F., T.E. Nickelson, S.L. Johnson, and S.V.D. Wetering. 1997. Juvenile sea-run cutthroat trout: habitat utilization, smolt production, and response to habitat modification. Pages 148-150 in J.D. Hall, P.A. Bisson, and R.E. Gresswell, eds. Sea-run cutthroat trout: biology, management, and future conservation. Oregon Chapter, American Fisheries Society, Corvallis.
- Solazzi, M.F., T.E. Nickelson, S.L. Johnson, J.D. Rodgers. 1998. Development and evaluation of techniques to rehabilitate Oregon's wild salmonids. Oregon Department of Fish and Wildlife, Fish Research Project F-125-R-13, Final Report, Portland.
- Soule, M. E. 1980. Thresholds for survival: maintaining fitness and evolutionary potential. Pages 151-170 in M.E. Soule and B.A. Wilcox, eds. Conservation biology. Sinauer Associates. Sunderland, Massachusetts.
- Sprague, C. R., L.G. Beckman, and S. D. Duke. 1993. Prey selection by juvenile white sturgeon in reservoirs of the Columbia River. Report N in: Status and Habitat Requirements of the White Sturgeon Populations in the Columbia River Downstream from McNary Dam. Project No. 86-50. Final Report of Research to the Bonneville Power Administration, U.S. Department of Energy, Portland, Oregon. Volume 2.

- Starke, G.M. and J.T. Dalen. 1995. Pacific Lamprey (*Lampetra tridentate*) Passage Patterns Past Bonneville Dam and Incidental Observations of Lamprey at the Portland District Columbia River Dams in 1993. U.S. Army Corps of Engineers, Bonneville Lock and Dam, Cascade Locks, Oregon.
- Stehr C.M., D.W. Brown, T. Hom, B.F. Anulacion, W.L. Reichert, and T.K. Collier. 2000. Exposure of juvenile chinook and chum salmon to chemical contaminants in the Hylebos Waterway of Commencement Bay, Tacoma Washington. *Journal of Aquatic Ecosystem Stress and Recovery* 7: 215–227.
- Stein, J.E., T. Hom, T.K. Collier, D.W. Brown, and U. Varanasi. 1995. Contaminant exposure and biochemical effects in outmigrant juvenile Chinook salmon from urban and nonurban estuaries of Puget Sound, Washington. *Environmental Toxicology and Chemistry* 14:1019-1029.
- Stein, R.A., P.E. Reimers, and J.D. Hall. 1972. Social interactions between juvenile coho (*Oncorhynchus kisutch*) and fall Chinook salmon (*O. tshawytscha*) in Sixes River, Oregon. *Journal of the Fisheries Research Board of Canada* 29:1737-1748.
- Stevens, D.E., and L.W. Miller. 1970. Distribution of sturgeon larvae in the Sacramento-San Joaquin river system. *California Fish and Game* 83:1-20.
- Stevens, D.G., A.V. Nebeker, et al. 1980. Avoidance responses of salmon and trout to air-supersaturated water. *Transactions of the American Fisheries Society* 109(6): 751-754.
- Stewart, D.J., D. Weininger, D.V. Rottiers and T.A. Edsall. 1983. An energetics model for lake trout, *Salvelinus namaycush*: application to the Lake Michigan population.
- Stockley, C.E. 1961. The migration of juvenile salmon past the Mayfield Dam site, Cowlitz River, 1955 and 1956. Washington Department of Fisheries.
- Stone, J., T. Sundlov, S. Barndt, T. Coley. – U.S. Fish and Wildlife Service. 2001. Evaluate Habitat Use and Population Dynamics of Lampreys in Cedar Creek. Annual Report. Project No. 2000-014-00. Contract No. 00000014.
- Stone, J., T. Sundlov, S. Barndt, T. Coley. – U.S. Fish and Wildlife Service. 2002. Evaluate Habitat Use and Population Dynamics of Lampreys in Cedar Creek. Annual Report. Project No. 2000-014-00. Contract No. 00000014.
- Stuehrenberg, L., K. Liscom, and G. Monan. 1978. A study of apparent losses of chinook salmon and steelhead based on count of discrepancies between dams on the Columbia and Snake Rivers, 1967-1968. 49 pp. (Available from the Northwest Fisheries Science Center, 2725 Montlake Blvd. E., Seattle, WA 98112-2097).
- Sullivan, K., D.J. Martin, R.D. Cardwell, J.E. Toll, and S. Duke. 2000. An analysis of the effects of temperature on salmonids of the Pacific Northwest with implications for selecting temperature criteria. Sustainable Ecosystems Institute. Portland, Oregon.
- Sullivan, K., J. Tooley, K. Doughty, J.E. Caldwell, and P. Knudsen. 1990. Evaluation of prediction models and characterization of stream temperature regimes in Washington. Timber, Fish, and Wildlife Report WQ3-90-006. Washington Department of Natural Resources, Olympia.
- Sumner, F.H. 1962. Migration and growth of coastal cutthroat trout in Tillamook County, Oregon. *Transactions of the American Fisheries Society* 91(1):77-83.
- Sumner, F.H. 1972. A contribution to the life history of the cutthroat trout in Oregon with emphasis on the coastal subspecies, *Salmo clarki clarki* Richardson. Oregon State Game Commission, Corvallis. 142 p.

- Swales, S., F. Caron, J.R. Irvine, and C.D. Levings. 1988. Overwintering habitats of coho salmon (*Oncorhynchus kisutch*) and other juvenile salmonids in the Keogh River system, British Columbia. *Canadian Journal of Zoology* 66:254-261.
- Swanston, D.N. 1991. Natural processes. Chapter 5 in Influences of forest and rangeland management on salmonid fishes and their habitats. W.R. Meehan, ed. American Fisheries Society Special Publication 19. Bethesda, Maryland.
- Tabor, R. A., R. S. Shively, and T. P. Poe. 1993. Predation of juvenile salmonids by smallmouth bass and northern pikeminnow in the Columbia River near Richland, Washington. *North American Journal of Fisheries Management* 13: 831-838.
- Tagart, J.V. 1984. Coho salmon survival from egg deposition to fry emergence, Pages 173-181. in J.M. Walton and D.B. Houston, eds. Proceedings of the Olympic Wild Fish Conference, March 23-25, 1983. Fisheries Technology Program, Peninsula College, Port Angeles, Washington.
- Technical Advisory Committee (TAC) to parties of U.S. v Oregon. 2002. Biological Assessment of Incidental Impacts on Salmon Species Listed under the Endangered Species Act in the 2002 Non-Indian and Treaty Indian Fall Season Fisheries in the Columbia River Basin.
- Thompson, G.G. 1991. Determining Minimum Viable Populations under the Endangered Species Act. NOAA Technical Memorandum NMFS F/NMC-198, NMFS, Seattle, Washington.
- Thompson, R.B. 1959. Food of the pikeminnow *Ptychocheilus oregonensis* (Richardson) of the Lower Columbia River. *Fisheries Bulletin* 60:43-58.
- Thompson, W.L., and D.C. Lee. 2000. Modeling relationships between landscape-level attributes and snorkel counts of Chinook salmon and steelhead parr in Idaho. *Canadian J. of Fisheries and Aquatic Sciences* 57: 1834-1842.
- Thorgaard, G.H. 1983. Chromosomal differences among rainbow trout populations. *Copeia* 1983(3):650-663.
- Timble, S.W. and A.C. Mendel. 1995. The cow as a geomorphic agent – a critical review. *Geomorphology* 13:233-253.
- Tipping, J., S. Springer, P. Buckley, and J. Danielson. 1979. Cowlitz River Steelhead Spawning, Fry Emergence and Stranding, 1977-79, and Adult Life History Study, 1977-79. Washington Department of Game.
- Tipping, J. 1984 A profile of Cowlitz River winter steelhead before and after hatchery propagation. Washington Department of Game 84-11.
- Tipping, J.M. 1981. Cowlitz sea-run cutthroat study 1980-1981. Washington Department of Game Fish. Management Division, Rep. 81-12, Olympia, Washington.
- Tipping, J.M. 1986. Effect of release size on return rates of hatchery sea-run cutthroat trout. *Prog. Fish-Cult.* 48(3):195-197.
- TOAST (Oregon Technical Outreach and Assistance Team). 2004. Understanding out-of-subbasin effects for Oregon subbasin planning with particular reference to Ecosystem Diagnosis and treatment assessments.
- Tomasson, T. 1978. Age and growth of cutthroat trout, *Salmo clarki clarki* Richardson, in the Rogue River, Oregon. Master's thesis, Oregon State University, Corvallis. 75 pp.

- Tredger, C. C. 1980. Carrying capacity and theoretical steelhead smolt yield from Nuaitch Creek, Nicola River system. British Columbia Ministry of Environment, Fish and Wildlife Branch, Victoria, British Columbia. 46 pp.
- Tredger, D. 1979. An evaluation of fish habitat and fish populations in Toboggan Creek near Smithers, relevant to steelhead enhancement opportunities. British Columbia Ministry of Environment, Fish and Wildlife Branch, Victoria.
- Tripp, D., and P. McCart. 1983. Effects of different coho stocking strategies on coho and cutthroat trout production in isolated headwater streams. Can. Tech. Rep. of Fisheries and Aquatic Sciences 1212:176 p.
- Trotter, P.C. 1989. Coastal cutthroat trout: A life history compendium. Transactions of the American Fisheries Society 118:463-473.
- Trotter, P.C. 1997. Sea-run cutthroat trout: life history profile. Pages 7-15 in J.D. Hall, P.A. Bisson and R.E. Gresswell, eds. Sea-run cutthroat trout: biology, management, and future conservation, American Fisheries Society, Corvallis, Oregon.
- Tschaplinski, P.J. 2000. The effects of forest harvesting, fishing, climate variation, and ocean conditions on salmonid populations of Carnation Creek, Vancouver Island, British Columbia. in Sustainable Fisheries Management: Pacific Salmon, CRC Press, Boca Raton, FL.
- Turman, D.L. 1972. Studies on spawning boxes for coastal cutthroat trout. Master's thesis, Humboldt State University, Arcata, California.
- Tyler, R.W. 1964. Distribution and migration of young salmon in Bellingham Bay, Washington. Circular 212, Fish. Research Institute, University of Washington, Seattle. 26 p.
- U.S. v. Oregon. Technical Advisory Committee. 1997. 1996 All species review. Columbia River Fish Management.
- USFWS (U.S. Department of Interior, Fish and Wildlife Service). 2002. 50 Code of Federal Regulations, Part 17. Endangered and threatened wildlife and plants; withdrawal of proposed rule to list the southwestern Washington/Columbia River distinct population segment of the coastal cutthroat trout as threatened; proposed rule.
- USFWS (U.S. Department of Interior, Fish and Wildlife Service). 1988. Trinity River flow evaluation. Annual Report, 1988, Sacramento, California.
- USFWS (U.S. Department of Interior, Fish and Wildlife Service). 1990. Trinity River flow evaluation. Annual Report, 1990, Sacramento, California.
- USFWS (U.S. Department of Interior, Fish and Wildlife Service). 1994. Determination of endangered status for the Kootenai River white sturgeon population. Federal Register 59(171) 45989.
- USFWS (U.S. Department of Interior, Fish and Wildlife Service). 1999. Recovery plan for the white sturgeon (*Acipenser transmontanus*): Kootenai River population. U.S. Fish and Wildlife Service, Portland, Oregon.
- USFWS (U.S. Department of Interior, Fish and Wildlife Service). 2002. Bull Trout (*Salvelinus confluentis*) Draft Recovery Plan. U.S. Fish and Wildlife Service, Portland, Oregon.
- USFWS (U.S. Department of Interior, Fish and Wildlife Service). 2002. Listing determination register notice
- Utter F.M. and F.W. Allendorf. 1977. Determination of the breeding structure of steelhead populations through gene frequency analysis. Pages 44-54 in T.J. Hassler and R.R. VanKirk, eds. Proceedings

- of the Genetic Implications of Steelhead Management Symposium, May 20-21, 1977, Arcata, California. California Cooperative Fish Research Unit Special Report 77-1.
- Utter, F., G. Milner, G. Stahl, and D. Teel. 1989. Genetic population structure of chinook salmon (*Oncorhynchus tshawytscha*), in the Pacific northwest. Fisheries Bulletin (U.S.) 87:239-264.
- Utter, F.M., D. Campton, S. Grant, G. Milner, J. Seeb, and L. Wishard. 1980. Population structures of indigenous salmonid species of the Pacific Northwest. Pages 285-304 in W.J. McNeil and D.C. Himsworth, eds. Salmonid ecosystems of the North Pacific. Oregon State University Press, Corvallis.
- Utter, F.M., D. Teel, G. Milner, and D. McIlsac. 1987. Genetic estimates of stock composition of 1983 Chinook salmon, *Oncorhynchus tshawytscha*, in the Pacific Northwest. Fish Bull. 87:239-264.
- Uusitalo, N. 2001. Evaluating factors limiting Columbia River Gorge chum salmon populations. Report to Bonneville Power Administration. DOE/BP – 000004669-2.
- Van Eenennaam, J.P., and S.I. Dorsohov. 2001. Reproductive conditions of Klamath River green sturgeon (*Acipenser medirostris*). Proceedings of International Symposium on Sturgeon, Wisconsin. (In review for Journal of Applied Ichthyology).
- Van Winkle, W., P.J. Anders, D.H. Secor, and D.A. Dixon, eds. 2002. Biology, Management, and Protection of Sturgeons. American Fisheries Society Symposium 28. 258 pp.
- Vella, J.J., L.C. Stuehrenberg, and T.C. Bjornn. 1997. Migration patterns of Pacific lamprey *Lampetra tridentata* in the lower Columbia River. Annual report of Research, U.S. Army Corps of Engineers.
- Vigg, S. 1988. Functional response of northern pikeminnow predation to salmonid prey density in McNary tailrace. Pages 174-207 in T.P. Poe and B.E. Rieman, eds. Predation by resident fish on juvenile salmonids in John Day Reservoir, 1983-1986. Final Report (Contracts DE-AI79-82BP34796) and DE-AI79-82BP35097) to Bonneville Power Administration, Portland, Oregon.
- Vigg, S., T.P. Poe, L.A. Prendergast, and H.C. Hansel. 1991. Rates of consumption of juvenile salmonids and alternative prey fish by northern pikeminnow, walleyes, smallmouth bass, and channel catfish in John Day Reservoir, Columbia River. Transactions of the American Fisheries Society 120:421-438.
- Vigg, S., T.P. Poe, L.A. Prendergast, and H.C. Hansel. 1991. Rates of consumption of juvenile salmonids and alternative prey fish by northern squawfish management program. Oregon Department of Fish and Wildlife, Contract number DE-B179-90BP07094 and 94BI24514. Final report of research, 1990-96, to the Bonneville Power Administration, Portland, Oregon.
- Vladykov, V.D. and E. Kott. 1979. A new parasitic species of the holarctic lamprey genus *Entosphenus* Gill, 1862 (Petroyzonidae) from Klamath River in California and Oregon. Canadian Journal of Zoology 57: 808-823.
- Vronskiy, B.B. 1972. Reproductive biology of the Kamchatka River Chinook salmon (*Oncorhynchus tshawytscha* (Walbaum)). Journal of Ichthyology 12:259-273.
- WAC (Washington Administrative Code). 2000. Forest Practice Rules. Washington Department of Natural Resources Forest Practices Board.
- Wade, G. 2000. Salmon and Steelhead Habitat Limiting Factors, WRIA 26 (Cowlitz). Washington Department of Ecology.
- Wade, G. 2000. Salmon and Steelhead Habitat Limiting Factors, WRIA 27 (Lewis). Washington Department of Ecology.

- Wade, G. 2001. Salmon and Steelhead Habitat Limiting Factors, WRIA 28 (Salmon-Washougal). Washington Department of Ecology.
- Wade, G. 2002. Salmon and Steelhead Habitat Limiting Factors, WRIA 25 (Grays-Elochoman). Washington Department of Ecology.
- Wahle, R.J. and R.Z. Smith. 1979. A historical and descriptive account of Pacific coast anadromous salmonid rearing facilities and a summary of their releases by region, 1960-76. National Marine Fisheries Service (NMFS), NOAA Technical Report NMFS SSRF-736.
- Waknitz, F.W., G.M. Matthews, T. Wainwright, and G.A. Winans. 1995. Status review for Mid-Columbia River summer chinook salmon. NOAA Technical Memorandum NMFN-NWFSC-22. 80 pp.
- Waldman, J.R. 1995. Sturgeon and paddlefishes: A convergence of biology, politics, and greed. *Fisheries* 20(9):20-49.
- Wales, J.H. and M. Coots. 1954. Efficiency of Chinook salmon spawning in Fall Creek, California. *Transactions of the American Fisheries Society* 84:137-149.
- Walters, C.J. and J.R. Post. 1993. Density –dependent growth and competitive asymmetries in size-structured fish populations: a theoretical model and recommendations for field experiments. *Transactions of the American Fisheries Society* 122: 34-45.
- Walters, C.J., J.S. Collie, and T. Webb. 1988. Experimental designs for estimating transient responses to management disturbances. *Canadian Journal of Fisheries and Aquatic Sciences* 45:530-538.
- Walters, C.J., R. Hilborn, R.M. Peterman, and M.J. Staley. 1978. Model for examining early ocean limitation of Pacific salmon production. *Journal of the Fisheries Research Board of Canada* 35:1303-1315.
- Walters, C.J./Cahoon, P. 1985. Evidence of decreasing spatial diversity in British Columbia salmon stocks. *Canadian Journal of Fisheries and Aquatic Sciences* 42:1033-1037.
- Wang, Y. L., F. P. Binkowski, and S. I Doroshov. 1985. Effect of temperature on early development of white and lake sturgeon *Acipenser transmontanus*, and *A. fulvescens*. *Environmental Biology of Fishes* 14:43-50.
- Wangaard, D.B., and C.B. Burger. 1983. Effects of various water temperature regimes on the egg and alevin incubation of Susitna River chum and sockeye salmon. U.S. Fish and Wildlife Service, Anchorage, Alaska. 43 p.
- Waples, R.S. 1990. Conservation genetics of Pacific salmon. II. Effective population size and the rate of loss of genetic variability. *Journal of Heredity* 81: 267-276.
- Waples, R.S. 1991a. Genetic interactions between hatchery and wild salmonids: lessons from the Pacific Northwest. *Canadian Journal of Fisheries and Aquatic Sciences* 48 (Suppl 1): 124-133.
- Waples, R.S. 1991b. Definition of "species" under the endangered species act: Application to Pacific salmon. National Marine Fisheries Service, Northwest Fisheries Center, Coastal Zone and Estuarine Studies Division, Seattle.
- Waples, R.S. and C. Do. 1994. Genetic risk associated with supplementation of Pacific salmonids: captive broodstock programs. *Canadian Journal of Fisheries and Aquatic Sciences* 51: 310-329.
- Waples, R.S., J. Robert, P. Jones, B.R. Beckman, and G.A. Swan. 1991. Status review for Snake River fall chinook salmon. NOAA Technical Memorandum NMFS N/NWC-201, Seattle, Washington. 73 pp.

- Waples, R.S., O.W. Johnson, and R.P. Jones Jr. 1991. Status review for Snake River sockeye salmon. NOAA Technical Memorandum F/NWC-195, U.S. Department of Commerce, National Marine Fisheries Service, Portland, Oregon.
- Ward, A.D, and W.J. Elliot. 1995. Environmental Hydrology. CRC Press, Boca Raton, Florida.
- Ward, B. R., and P. A. Slaney. 1988. Life history and smolt-to-adult survival of Keogh River steelhead trout (*Salmo gairdneri*) and the relationship to smolt size. Canadian Journal of Fisheries and Aquatic Sciences 45: 1110-1122.
- Ward, B. R., and P. A. Slaney. 1993. Egg-to-smolt survival and fry-to-smolt density dependence of Keogh River steelhead in R.J. Gibson and R.E. Cutting, eds. Production of juvenile Atlantic Salmon, *Salmo salar*, in natural waters. Canadian Journal of Fisheries and Aquatic Science.
- Ward, D. L. and M. P. Zimmerman. 1999. Response of smallmouth bass to sustained removals of northern pikeminnow in the lower Columbia and Snake rivers. Transactions of the American Fisheries Society 128:1020-1035.
- Ward, D. L. and M. P. Zimmerman. 1999. Response of smallmouth bass to sustained removals of northern pikeminnow in the lower Columbia and Snake rivers. Transactions of the American Fisheries Society 128:1020-1035.
- Ward, D.L. 2001. Lamprey Harvest at Willamette Falls, 2001. Oregon Department of Fish and Wildlife, Clackamas.
- Ward, D.L., J.H. Petersen, and J.J. Loch. 1995. Index of predation on juvenile salmonids by northern pikeminnow in the lower and middle Columbia River, and in the lower Snake River. Transactions of the American Fisheries Society 124:321-334.
- Ward, D.L., K. Collis, J.H. Petersen, D.D. Roby, and S.P. Barnes. 2002. Draft Predator Control Program Summary (Mainstem/Systemwide Province). Prepared for the Northwest Power Planning Council. October 24, 2002.
<http://www.cbfwa.org/files/province/systemwide/subsum/021024Predation.doc>
- Ward, R. D., N. Billington, and P. D. N. Hebert. 1989. Comparison of allozyme and mitochondrial DNA variation in populations of walleye, *Stizostedion vitreum*. Canadian Journal of Fisheries and Aquatic Sciences 46: 2074-2084.
- Warren, J.J. and L.G. Beckman. 1993. Fishway use by white sturgeon to bypass mainstem Columbia River dams. U.S. Fish and Wildlife Service Sea Grant Extension Project, Columbia River Series WSG-AG 93-02.
- Washington State Natural Resources Cabinet. 1999. State wide strategy to recover salmon: extinction is not an option. Washington Governor's Salmon Recovery Office. Olympia.
- WCSBRT (West Coast Salmon Biological Review Team). 2003. Preliminary conclusions regarding the updated status of listed ESUs of West Coast salmon and steelhead. NOAA Fisheries Northwest Fisheries Science Center, Seattle, WA, and Southwest Fisheries Science Center, Santa Cruz, CA. Co-manager review draft.
- WDF (Washington Department of Fisheries). 1990. Elochoman River Subbasin Salmon and Steelhead Production Plan. Olympia, Washington.
- WDF (Washington Department of Fisheries). 1990. Washougal River subbasin salmon and steelhead production plan. Columbia Basin System Planning. Northwest Power Planning Council, and the Agencies and Indian Tribes of the Columbia Basin Fish and Wildlife Authority. September 1990. 163 p.

- WDF and WDW (Washington Department of Fisheries and Washington Department of Wildlife). 1993. 1992 Washington State salmon and steelhead stock inventory. Appendix 3 – Columbia River stocks. Washington Department of Fisheries and Washington Department of Wildlife, Olympia.
- WDF, WDW, and WWTIT (Washington Department of Fisheries, Washington Department of Wildlife, and Western Washington Treaty Indian Tribes). 1993. 1992 Washington state salmon and steelhead stock inventory (SASSI). Washington Department of Fish and Wildlife, 212 pp. + appendices. Appendix 1: Hood Canal and Strait of Juan de Fuca (December 1994, 424 pp.), North Puget Sound (June 1994, 418 pp.), and South Puget Sound (September 1994, 371 pp.) volumes. Appendix 2: Coastal stocks (August 1994, 587 pp.). Appendix 3: Columbia River stocks (June 1993, 580 pp.). Washington Department of Fish and Wildlife, P.O. Box 43151, Olympia, WA 98504.
- WDFW (Washington Department of Fish and Wildlife). 1997. Final environmental impact statement for the wild salmonid policy. Washington Department of Fish and Wildlife, Olympia.
- WDFW (Washington Department of Fish and Wildlife). 2001. Lower Columbia River Fisheries Management and Evaluation Plan (updated January 9, 2003).
- WDFW (Washington Department of Fish and Wildlife). 1993. 1992 Washington State Salmon and Steelhead Stock Inventory.
- WDFW (Washington Department of Fish and Wildlife). 1998. Integrated landscape management plan for fish and wildlife in the Lewis-Kalama River watershed, Washington: a pilot project. Olympia, Washington.
- WDFW (Washington Department of Fish and Wildlife). 1998. Washington State Salmonid Stock Inventory- Bull Trout/Dolly Varden.
- WDFW (Washington Department of Fish and Wildlife). 2000. Future Brood Document. Hatcheries Division, Fish Program, Washington Department of Fish and Wildlife, Olympia.
- WDFW (Washington Department of Fish and Wildlife). 2001. Draft Lower Columbia Chum Stock Inventory. Washington Department of Fish and Wildlife, Olympia. 37
- WDOE (Washington Department of Ecology). 1998. 303(d) list of threatened and impaired water bodies. Washington Department of Ecology, Olympia.
- WDW (Washington Department of Wildlife). 1990. Salmon and Steelhead Production Plan – Kalama River Subbasin. Olympia, Washington.
- WDW (Washington Department of Wildlife). 1990. Salmon and Steelhead Production Plan - Little White Salmon River Subbasin. Olympia, Washington.
- WDW (Washington Department of Wildlife). 1990. Cowlitz River subbasin salmon and steelhead production plan. Columbia Basin System Planning. Northwest Power Planning Council, and the Agencies and Indian Tribes of the Columbia Basin Fish and Wildlife Authority. September 1990. 163 pp.
- Weaver, T.M. and R.G. White. 1985. Coal Creek fisheries monitoring study. No III. Final Report to U.S. Department of Agriculture, Forest Service, Flathead National Forest. Montana Cooperative Fisheries Research Unit, Bozeman.
- Welch, D.W., B.R. Ward, B.D. Smith, F. Whitney. 1997. Changes associated with the 1989-1990 ocean climate shift, and effects on British Columbia steelhead (*O. mykiss*) populations. Pacific Stock Assessment Review Committee (PSARC) Working Paper.

- Welch, E.B., J.M. Jacoby, and C.W. May. 1998. Stream Quality. Chapter 4 in River Ecology and Management – Lessons from the Pacific Coastal Ecoregion. R.J. Naiman and R.E. Bilby eds. Springer-Verlag. New York.
- WFPB (Washington Forest Practices Board). 2000. Washington Forest Practices Rules, Board Manual, and Forest Practices Act. Washington Department of Natural Resources. Olympia.
- Whitmus, C.J., and S. Olsen. 1979. The migratory behavior of juvenile chum salmon released in 1977 from the Hood Canal hatchery at Hoodsport, Washington. University of Washington. Fish. Research Institute FRI-UW-7916. 46 pp.
- Whitmus, C.J., Jr. 1985. The influence of size on the migration and mortality of early marine life history of juvenile chum salmon (*Oncorhynchus keta*). Master's thesis. University of Washington, Seattle. 69 pp.
- Whitney, R.R., L. Calvin, M. Erho, and C. Coutant. 1997. Downstream passage for salmon at hydroelectric projects in the Columbia River Basin: development, installation, and evaluation. U.S. Department of Energy, Northwest Power Planning Council, Portland, Oregon. Report 97-15. 101 pp.
- Whyte, J.N.C., R.J. Beamish, N.G. Ginther and C.E. Neville. 1993. Nutritional condition of the Pacific lamprey (*Lampetra tridentate*) deprived of food for periods of up to two years. Canadian Journal of Fisheries and Aquatic Sciences 50: 591-599.
- Wickett, W.P. 1954. The oxygen supply to salmon eggs in spawning beds. Journal of the Fisheries Research Board of Canada 11:933-953.
- Williams, I.V., and P. Gilhousen. 1968. Lamprey parasitism on Fraser River sockeye and pink salmon during 1967. International Pacific Salmon. Fish. Comm. Progress Report 18. 22 pp.
- Williams, R.N and 12 co-authors. 1999. Return to the river: Scientific issues in the restoration of salmonid fishes in the Columbia River. Fisheries 24(3):10-25.
- Willis, R. 1962. Gnat Creek Weir Studies. Oregon Fish Commission, Portland.
- Wilson, G.M., W.K. Thomas, and A.T. Beckenbach. 1985. Intra- and inter-specific mitochondrial DNA sequence divergence in *Salmo*: rainbow, steelhead, and cutthroat trouts. Canadian Journal of Zoology 63:2088-2094.
- Winans, G.A. 1989. Genetic variability in Chinook salmon stocks from the Columbia River basin. North American Journal of Fisheries Management 9:47-52.
- Wipfli, M.S., J.P. Hudson, D.T. Chaloner, and J.P. Caouette. 1999. Influence of salmon spawner densities on stream productivity in Southeast Alaska. Canadian Journal of Fisheries and Aquatic Sciences 56:1600-1611.
- Withler, I. L. 1966. Variability in life history characteristics of steelhead trout (*Salmo gairdneri*) along the Pacific Coast of North America. Journal of the Fisheries Research Board of Canada 23: 365-393.
- WJNRC (Washington Joint Natural Resources Cabinet). 2001 guidance on watershed assessment for salmon). Governor's Salmon Recovery Office, Olympia, Washington.
- Wolcott R.S.C., Jr. 1978. The chum salmon run at Walcott Slough. Special report to U.S. Department of Interior, Fish and Wildlife Service, Reno, Nevada. 41 p.
- Wood, C.C., B.E. Riddell, and D.T. Rutherford. 1987. Alternative juvenile life histories of sockeye salmon (*Oncorhynchus nerka*) and their contribution to production in the Stikine River, northern British Columbia. Pages 12-24 In H.D. Smith, L. Margolis, and C.C. Wood, editors. Sockeye salmon

- (*Oncorhynchus nerka*) population biology and future management. Canadian Special Publication of Fisheries and Aquatic Sciences 96, Ottawa.
- Woody, C.A. 1998. Ecological, morphological, genetic, and life history comparison of two sockeye salmon populations, Tustumena Lake, Alaska. Ph.D. Dissertation, University of Washington. 117 p.
- Woody, C.A., J. Olsen, J. Reynolds, and P. Bentzen. 2000. Temporal variation in phenotypic and genotypic traits in two sockeye salmon populations, Tustumena Lake, Alaska. Transactions of the American Fisheries Society 129: 1031-1043.
- Wright, S. 1931. Evolution in Mendelian populations. Genetics 16:97-159.
- Wydowski, R. S., and R. R. Whitney. 1979. Inland Fishes of Washington. University of Washington Press, Seattle.
- Wyzga, B. 1993. River response to channel regulation: Case study of the Raba River, Carpathians, Poland. Earth Surface Processes and Landforms 18: 541-556.
- Yakima Indian Nation (YIN), Washington Department of Fisheries and Washington Department of Wildlife. 1990. Yakima River sub-basin salmon and steelhead production plan. Columbia Basin Fish and Wildlife Authority, Portland, Oregon.
- YIN, WDW, WDF. 1990. Yakima River Subbasin Salmon and Steelhead Plan. Northwest Power Planning Council. Portland, Oregon. 282 pages.
- Zabel, R. W. 2003. Use of age-structured population projection matrices to develop “out-of-subbasin” survival estimates. Decision draft. NOAA Fisheries, Northwest Fisheries Science Center.
- Zanadrea, G. 1961. Studies on European lampreys. Evolution. 15:523-534.
- Ziemer, R.R. and T.E. Lisle. 1998. Hydrology. Chapter 3 in River Ecology and Management. Naiman and Bilby eds.
- Zimmerman, C. E., and G. H. Reeves. 1999. Steelhead and Resident Rainbow Trout: Early Life History and Habitat Use in the Deschutes River, Oregon. Prepared for Portland General Electric Co., Oregon State University, Department of Fish and Wildlife, Corvallis.
- Zimmerman, C.E. 1995. Population structure of coastal cutthroat trout (*Oncorhynchus clarki clarki*) in the Muck Creek Basin, Washington. Master’s thesis, Oregon State University, Corvallis.
- Zimmerman, M.P., and D.L. Ward. 1999. Index of predation on juvenile salmonids by northern pikeminnow in the lower Columbia River basin from 1994-96. Transactions of the American Fisheries Society 128:995-1007.